

**CHAPTER-3**  
**TRIGONOMETRIC FUNCTIONS**  
**02 MARK TYPE QUESTIONS**

Q. NO	QUESTION	MARK
1.	If $\tan \tan x = \frac{b}{a}$ , then find the value of $\sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}}$	2
2.	A circular wire of radius 3cm is cut and bent so as to lie along the circumference of a loop whose radius is 48cm. Find the angle in degrees which is subtended at the centre of the loop.	2
3.	Find the radian measure corresponding $37^\circ 30'$ .	2
4.	The minute hand of a clock is 10 cm long. How far does the tip of the hand move in 20 minutes?	2
5.	Find the values of $\sin \sin \left( -\frac{11\pi}{3} \right)$	2
6.	Prove that $\sin 5x - 2\sin 3x + \sin x / \cos 5x - \cos x = \tan x$	2
7.	The minute hand of a watch is 1.5 cm long. How far does its tip move in 40 minutes?	2
8.	Show that $\tan 3x \cdot \tan 2x \cdot \tan x = \tan 3x - \tan 2x - \tan x$	2
9.	Prove that $\cos 6x = 32\cos^2 2x - 48\cos^4 4x + 18\cos^2 2x - 1$	2
10.	A wheel makes 360 revolutions in 1 minute. How many radians does it turn in 1 second?	2
11.	Find the degree measure of the angle subtended at the center of a circle of radius 100cm by an arc of length 22cm.	2
12.	Prove that $\frac{\sin x - \sin 3x}{\sin^2 x - \cos^2 x} = 2 \sin x$ .	2
13.	Prove that $\cos 4x = 1 - 8 \sin^2 x \cos^2 x$ .	2
14.	Find the value of $2 \sin \frac{5\pi}{4} + 2 \cos^2 \frac{\pi}{4} + 2 \sec^2 \frac{\pi}{3}$ .	2
15.	Find the value of $\tan 15^\circ$	2

**ANSWERS:**

Q. NO	ANSWER	MARKS
1.	<p>Given, <math>\tan \tan x = \frac{b}{a}</math></p> $\therefore \sqrt{\frac{a+b}{a-b}} + \sqrt{\frac{a-b}{a+b}} = \frac{\sqrt{(a+b)^2} + \sqrt{(a-b)^2}}{\sqrt{(a-b)(a+b)}} = \frac{(a+b)+(a-b)}{\sqrt{a^2-b^2}} = \frac{2a}{a\sqrt{1-\left(\frac{b}{a}\right)^2}} = \frac{2}{\sqrt{1-\tan^2 x}}$ $\frac{2 \cos \cos x}{\sqrt{\cos^2 x - \sin^2 x}} = \frac{2 \cos \cos x}{\sqrt{\cos \cos 2x}}$	2
2.	<p>Solution:  Length of the wire <math>= 2\pi r = 2\pi \times 3 = 6\pi</math>  We know, <math>\theta = \frac{l}{r} = \frac{6\pi}{48} = \frac{\pi}{8}</math> radians <math>= \frac{\pi}{8} \times \frac{180^\circ}{\pi} = 22.5^\circ</math></p>	2
3.	<p>Solution:  <math>60' = 1^\circ</math>  <math>\therefore 30' = \left(\frac{1}{2}\right)^\circ</math>  <math>\therefore -37^\circ 30' = 37\frac{1}{2}^\circ = -\frac{75}{2} \times \frac{2\pi}{360}</math> radian <math>= -\frac{5\pi}{24}</math> radians</p>	2
4.	<p>Solution:  The minute hand moves through <math>120^\circ</math>  So <math>\theta = \frac{2\pi}{3}</math> radians  We know, <math>l = r\theta = 10 \times \frac{2\pi}{3} = \frac{20\pi}{3}</math></p>	2
5.	<p>Solution:  <math>\sin \sin \left(-\frac{11\pi}{3}\right) = -\sin \sin \left(4\pi - \frac{\pi}{3}\right) = \sin \sin \frac{\pi}{3} = \frac{\sqrt{3}}{2}</math></p>	2
6.	<p>Starting with the left-hand side and using the trigonometric difference identities for the sine function, we obtain</p> $\begin{aligned} \text{L.H.S.} &= \sin 5x + \sin x - 2 \sin 3x / \cos 5x - \cos x \\ &= 2 \sin 3x \cos 2x - 2 \sin 3x / -2 \sin 3x \sin 2x \\ &= 2 \sin 3x (\cos 2x - 1) / -2 \sin 3x \sin 2x \\ &= -(1 - \cos 2x) / -\sin 2x \\ &= 2 \sin 2x / 2 \sin x \cos x \\ &= \sin x / \cos x \\ &= \tan x \end{aligned}$	2
7.	Using the above given information, we have,	2

	<p><math>r=1.5\text{cm}</math></p> <p>Angle made in 60min=<math>360^\circ</math></p> <p>Angle made in 1min=<math>6^\circ</math></p> <p>Angle made in 40min=<math>6^\circ \times 40 = 240^\circ</math></p> <p>Calculating the arc distance</p> <p><math>\theta=l/r</math></p> <p><math>240 \times \pi/180 = l/1.5</math></p> <p><math>2 \times 3.14 = l</math></p> <p><math>6.28 = l</math></p> <p><math>l = 6.28\text{cm}</math></p>	
8.	<p>Let us take <math>\tan 3x</math> and we know that <math>3x=2x+x</math></p> <p><math>\tan 3x = \tan(2x+x)</math></p> <p><math>\tan 3x/1 = \tan 2x + \tan x/1 - \tan 2x \cdot \tan x</math></p> <p><math>\tan 3x(1 - \tan 2x \cdot \tan x) = \tan 2x + \tan x</math></p> <p><math>\tan 3x - \tan 3x \cdot \tan 2x \cdot \tan x = \tan 2x + \tan x</math></p> <p><math>\tan 3x \cdot \tan 2x \cdot \tan x = \tan 3x - \tan 2x - \tan x</math></p>	2
9.	<p>Starting with the left-hand side and using the trigonometric identities for the cosine function, we obtain</p> <p>L.H.S.</p> <p><math>= \cos 6x</math></p> <p><math>= \cos 2(3x) = 2\cos^2 3x - 1</math></p> <p><math>= \cos 2(3x)</math></p> <p><math>= 2(4\cos^3 x - 3\cos x)^2 - 1</math></p> <p><math>= 2[16\cos^6 x + 9\cos^2 x - 24\cos^4 x] - 1</math></p> <p><math>= 32\cos^6 x + 18\cos^2 x - 48\cos^4 x - 1</math></p>	2

	=32cos6x-48cos4x+18cos2x1  =R.H.S.	
10.	Given,  Number of revolutions made in 60s=360  Number of revolutions made in 1s=360/60  Angle moved in 6 revolutions = $2\pi \times 6 = 12\pi$	2
11.	$\theta = \frac{l}{r} = \frac{22}{100} rad = \frac{11}{50} rad$ $= \frac{11}{50} \times \frac{180}{\pi} = 12^\circ 36'$	2
12.	$LHS = \frac{\sin x - \sin 3x}{\sin^2 x - \cos^2 x} = \frac{2\cos 2x \sin(-x)}{-\cos 2x} = 2\sin x$	2
13.	$LHS = \cos(2 \times 2x) = 1 - 2 \sin^2 2x = 1 - 2(2 \sin x \cos x)^2 = 1 - 8 \sin^2 x \cos^2 x.$	2
14.	$2 \sin \frac{5\pi}{4} + 2 \cos^2 \frac{\pi}{4} + 2 \sec^2 \frac{\pi}{3} = -\sqrt{2} + 1 + 8 = 9 - \sqrt{2}$	2
15.	$\tan 15^\circ = \tan(60^\circ - 45^\circ) = \frac{\tan 60^\circ - \tan 45^\circ}{1 + \tan 60^\circ \tan 45^\circ} = \frac{\sqrt{3} - 1}{\sqrt{3} + 1} = 2 - \sqrt{3}$	2