



CHAPTER-3

TRIGONOMETRIC FUNCTIONS

05 MARK TYPE QUESTIONS

Q. NO	QUESTION	MARK
1.	<p>One day while sitting on the bridge across a river Arun observes angles of depression of the banks on opposite sides of the river are 30° and 60° respectively as shown in the figure. (Take $\sqrt{3} = 1.73$)</p>  <p>Based on the above information, answer the following questions.</p> <ol style="list-style-type: none"> If the bridge is at height of 6m , then $AD=$ <ol style="list-style-type: none"> 6m $\frac{\sqrt{3}}{6}$m $6\sqrt{3}$m $\frac{6}{\sqrt{3}}$m Width of the river is <ol style="list-style-type: none"> 10.85m 11.87m 13.85m 19.85m The angles of the elevation and depression are always <ol style="list-style-type: none"> Acute angles obtuse angles right angles straight angles If $BD=21m$, then height of the bridge is <ol style="list-style-type: none"> 7m 21m $7\sqrt{3}$m $\frac{7}{\sqrt{3}}$m $BD=$ <ol style="list-style-type: none"> 6m $6\sqrt{3}$m $\sqrt{3}$m $10\sqrt{3}$m 	5
2.	<p>A submarine is moving in such way that at a particular moment of time its angle of elevation for two ships, situated at different positions on the surface of water, is α and β respectively. If $\operatorname{cosec}\alpha = \sqrt{3}$ and $\sec\sec\beta = 2$, then answer the following.</p>	5



1. What will be the value of $\sec \alpha$?
 a) $\sqrt{\frac{2}{3}}$ b) $\sqrt{\frac{3}{2}}$ c) $\frac{1}{\sqrt{3}}$ d) $\frac{1}{\sqrt{6}}$
2. What will be the measure of the angle β ?
 a) $\frac{\pi}{3}$ b) $\frac{\pi}{6}$ c) $\frac{\pi}{4}$ d) $\frac{\pi}{12}$
3. What will be the value of $\tan \alpha$?
 a) $\sqrt{3}$ b) $\frac{1}{\sqrt{3}}$ c) $\frac{1}{\sqrt{2}}$ d) $\frac{1}{\sqrt{6}}$
4. What will be the value of $\tan \beta$?
 a) $\sqrt{3}$ b) $\sqrt{2}$ c) $\frac{1}{\sqrt{3}}$ d) $\frac{\sqrt{2}}{\sqrt{3}}$
5. What will be the value of $\tan(\alpha + \beta)$?
 a) $\sqrt{3} - \sqrt{2}$ b) $\sqrt{6} + 1$ c) $\frac{1+\sqrt{6}}{\sqrt{2}-\sqrt{3}}$ d) $-$

3.	Prove that $(1 + \cos \frac{\pi}{8})(1 + \cos \frac{3\pi}{8})(1 + \cos \frac{5\pi}{8})(1 + \cos \frac{7\pi}{8}) = \frac{1}{8}$.	5
4.	Prove that $\cos^2 x + \cos^2(x + \frac{2\pi}{3}) + \cos^2(x - \frac{2\pi}{3}) = \frac{3}{2}$.	5

ANSWERS:

Q. NO	ANSWER	MARKS
1.	i) d) ii) c) iii) a) iv) c) v) b)	
2.	i) b) ii) a) iii) c) iv) a) v) c)	
3.	$\cos \frac{7\pi}{8} = \cos \left(\pi - \frac{\pi}{8}\right) = -\cos \frac{\pi}{8}, \cos \frac{5\pi}{8} = \cos \left(\pi - \frac{3\pi}{8}\right) = -\cos \frac{3\pi}{8}$ $\begin{aligned} \text{LHS} &= \left(1 + \cos \frac{\pi}{8}\right) \left(1 + \cos \frac{3\pi}{8}\right) \left(1 + \cos \frac{5\pi}{8}\right) \left(1 + \cos \frac{7\pi}{8}\right) \\ &= \left(1 + \cos \frac{\pi}{8}\right) \left(1 + \cos \frac{3\pi}{8}\right) \left(1 - \cos \frac{3\pi}{8}\right) \left(1 - \cos \frac{\pi}{8}\right) \\ &= \left(1 - \cos^2 \frac{\pi}{8}\right) \left(1 - \cos^2 \frac{3\pi}{8}\right) \\ &= \sin^2 \frac{\pi}{8} \sin^2 \frac{3\pi}{8} = \frac{1}{4} (2\sin^2 \frac{\pi}{8}) (2\sin^2 \frac{3\pi}{8}) \\ &= \frac{1}{4} (1 - \cos \frac{\pi}{4}) (1 - \cos \frac{3\pi}{4}) = \frac{1}{4} \left(1 - \frac{1}{\sqrt{2}}\right) \left(1 - \frac{1}{\sqrt{2}}\right) = \frac{1}{8} \end{aligned}$	5
4.	$\begin{aligned} \text{LHS} &= \cos^2 x + \cos^2(x + \frac{2\pi}{3}) + \cos^2(x - \frac{2\pi}{3}) \\ &= \frac{1}{2} \left(2 \cos^2 x + 2\cos^2(x + \frac{2\pi}{3}) + 2\cos^2(x - \frac{2\pi}{3})\right) \\ &= \frac{1}{2} \left[1 + \cos 2x + 1 + \cos 2\left(x + \frac{2\pi}{3}\right) + 1 + \cos 2\left(x - \frac{2\pi}{3}\right)\right] \\ &= \frac{1}{2} \left[3 + \cos 2x + \cos 2\left(x + \frac{2\pi}{3}\right) + \cos 2\left(x - \frac{2\pi}{3}\right)\right] \\ &= \frac{1}{2} \left[3 + \cos 2x + \cos\left(2x + \frac{4\pi}{3}\right) + \cos\left(2x - \frac{4\pi}{3}\right)\right] \\ &= \frac{1}{2} \left[3 + \cos 2x + 2 \cos 2x \cos\left(\frac{4\pi}{3}\right)\right] \\ &= \frac{1}{2} \left[3 + \cos 2x + 2 \cos 2x \left(-\frac{1}{2}\right)\right] \text{ As, } \cos\left(\frac{4\pi}{3}\right) = -\frac{1}{2} \\ &= \frac{3}{2} \end{aligned}$	5