### **DIRECTORATE OF EDUCATION** Govt. of NCT, Delhi

### SUPPORT MATERIAL (2024 - 2025)

### **Class : XII**

# **MATHEMATICS**

Under the Guidance of

### Shri Ashok Kumar

Secretary (Education)

### Shri R.N. Sharma

Director (Education)

### Dr. Rita Sharma

Addl. DE (School & Exam.)

### **Coordinators**

Mr. Sanjay Subhas Kumar Mrs. Ritu Singhal Mr. Raj Kumar DDE (Exam)

OSD (Exam)

OSD (Exam)

Mr. Krishan Kumar OSD (Exam)

**Production Team** Anil Kumar Sharma

Published at Delhi Bureau of Text Books, 25/2, Institutional Area, Pankha Road, New Delhi-58 by **Rajesh Kumar**, Secretary, Delhi Bureau of Text Books and Printed at: Arihant Offset, Plot No. 212F & 229, Nangli Sakrawati Industrial Area, Najafgarh Road, New Delhi

#### ASHOK KUMAR IAS



सचिव ( शिक्षा ) राष्ट्रीय राजधानी क्षेत्र दिल्ली सरकार पुराना सचिवालय, दिल्ली-110054 दूरभाषः 23890187 टेलीफैक्स : 23890119

Secretary (Education) Government of National Capital Territory of Delhi Old Secretariat , Delhi-110054 Phone : 23890187, Telefax : 23890119 E-mail : secyedu@nic.in

DE.5/228/EXAM) Metsage/SM/2018/555 Dated: 01/07/2024

#### MESSAGE

### In the profound words of Dr. Sarvepalli Radhakrishnan, "The true teachers are those who help us think for ourselves."

Every year, our teams of subject experts shoulder the responsibility of updating the Support Material to synchronize it with the latest changes introduced by CBSE. This continuous effort is aimed at empowering students with innovative approaches and techniques, thereby fostering their problemsolving skills and critical thinking abilities.

I am confident that this year will be no exception, and the Support Material will greatly contribute to our students' academic success.

The development of the support material is a testament to the unwavering dedication of our team of subject experts. It has been designed with the firm belief that its thoughtful and intelligent utilization will undoubtedly elevate the standards of learning and continue to empower our students to excel in their examinations.

I wish to extend my heartfelt congratulations to the entire team for their invaluable contribution in creating this immensely helpful resource for our students.

Wishing all our students a promising and bright future brimming with success.

(ASHOK KUMAR)

#### **R.N. SHARMA, IAS** Director, Education & Sports



Directorate of Education Govt. of NCT of Delhi Room No. 12, Old Secretariat Near Vidhan Sabha, Delhi-110054 Ph.: 011-23890172 E-mail : diredu@nic.in D $\pounds \cdot 5/228 | \pounds \times m/Metropel Sm/$ 2018 | 576 $Dated <math>\cdot 04/07/2024$ 

#### **MESSAGE**

It brings me great pleasure to present the support material specifically designed for students of classes IX to XII by our dedicated team of subject experts. The Directorate of Education remains resolute in its commitment to empower educators and students alike, extending these invaluable resources at no cost to students attending Government and Government-Aided schools in Delhi.

The support material epitomizes a commendable endeavour towards harmonizing content with the latest CBSE patterns, serving as a facilitative tool for comprehending, acquiring and honing essential skills and competencies stipulated within the curriculum.

Embedded within this initiative is a structured framework conducive to nurturing an analytical approach to learning and problem-solving. It is intended to prompt educators to reflect upon their pedagogical methodologies, forging an interactive conduit between students and academic content.

### In the insightful words of Rabindranath Tagore, "Don't limit a child to your own learning, for he was born in another time."

Every child is unique, with their own interests, abilities and potential. By allowing children to learn beyond the scope of our own experiences, we support their individual growth and development, helping them to reach their full potential in their own right.

May every student embrace the joy of learning and be empowered with the tools and confidence to navigate and shape the future.

(R. N. SHARMA)

Dr. RITA SHARMA Additional Director of Education (School/Exam)



Govt. of NCT of Delhi Directorate of Education Old Secretariat, Delhi-110054 Ph.: 23890185

D.O. No DE 5/228 Gram Me ung SM/ 2018/570 Dated: ...02/07/2024

#### **MESSAGE**

"Children are not things to be molded, but are people to be unfolded." - Jess Lair

In line with this insightful quote, the Directorate of Education, Delhi, has always made persistent efforts to nurture and unfold the inherent potential within each student. This support material is a testimony to this commitment.

The support material serves as a comprehensive tool to facilitate a deeper understanding of the curriculum. It is crafted to help students not only grasp essential concepts but also apply them effectively in their examinations. We believe that the thoughtful and intelligent utilization of these resources will significantly enhance the learning experience and academic performance of our students.

Our expert faculty members have dedicated themselves to the support material to reflect the latest CBSE guidelines and changes. This continuous effort aims to empower students with innovative approaches, fostering their problem-solving skills and critical thinking abilities.

I extend my heartfelt congratulations to the entire team for their invaluable contribution to creating a highly beneficial and practical support material. Their commitment to excellence ensures that our students are well-prepared to meet the challenges of the CBSE examinations and beyond.

Wishing you all success and fulfilment in your educational journey.

(Dr. Rita Sharma)

### DIRECTORATE OF EDUCATION Govt. of NCT, Delhi

SUPPORT MATERIAL (2024-2025)

# MATHEMATICS

Class : XII (English Medium)

**NOT FOR SALE** 

**PUBLISHED BY : DELHI BUREAU OF TEXTBOOKS** 

### भारत का संविधान

#### भाग 4क

# नागरिकों के मूल कर्तव्य

#### अनुच्छेद 51 क

मूल कर्तव्य - भारत के प्रत्येक नागरिक का यह कर्तव्य होगा कि वह -

- (क) संविधान का पालन करे और उसके आदशाँ, संस्थाओं, राष्ट्रध्वज और राष्ट्रगान का आदर करे;
- (ख) स्वतंत्रता के लिए हमारे राष्ट्रीय आंदोलन को प्रेरित करने वाले उच्च आदर्शों को हृदय में संजोए रखे और उनका पालन करे;
- (ग) भारत की संप्रभुता, एकता और अखंडता की रक्षा करे और उसे अक्षुण्ण बनाए रखे;
- (घ) देश की रक्षा करे और आहवान किए जाने पर राष्ट्र की सेवा करे;
- (ङ) भारत के सभी लोगों में समरसता और समान भ्रातृत्व की भावना का निर्माण करे जो धर्म, भाषा और प्रदेश या वर्ग पर आधारित सभी भेदभावों से परे हो, ऐसी प्रथाओं का त्याग करे जो महिलाओं के सम्मान के विरुद्ध हों;
- (च) हमारी सामासिक संस्कृति की गौरवशाली परंपरा का महत्त्व समझे और उसका परिरक्षण करे;
- (छ) प्राकृतिक पर्यावरण की, जिसके अंतर्गत वन, झील, नदी और वन्य जीव हैं, रक्षा करे और उसका संवर्धन करे तथा प्राणिमात्र के प्रति दयाभाव रखे;
- (ज) वैज्ञानिक दृष्टिकोण, मानववाद और ज्ञानार्जन तथा सुधार की भावना का विकास करे;
- (झ) सार्वजनिक संपत्ति को सुरक्षित रखे और हिंसा से दूर रहे;
- (अ) व्यक्तिगत और सामूहिक गतिविधियों के सभी क्षेत्रों में उत्कर्ष की ओर बढ़ने का सतत् प्रयास करे, जिससे राष्ट्र निरंतर बढ़ते हुए प्रयत्न और उपलब्धि की नई ऊँचाइयों को छू सके; और
- (ट) यदि माता-पिता या संरक्षक है, छह वर्ष से चौदह वर्ष तक की आयु वाले अपने, यथास्थिति, बालक या प्रतिपाल्य को शिक्षा के अवसर प्रदान करे।

**0.** 

### **Constitution of India**

Part IV A (Article 51 A)

### **Fundamental Duties**

It shall be the duty of every citizen of India —

- (a) to abide by the Constitution and respect its ideals and institutions, the National Flag and the National Anthem;
- (b) to cherish and follow the noble ideals which inspired our national struggle for freedom;
- (c) to uphold and protect the sovereignty, unity and integrity of India;
- (d) to defend the country and render national service when called upon to do so;
- (e) to promote harmony and the spirit of common brotherhood amongst all the people of India transcending religious, linguistic and regional or sectional diversities; to renounce practices derogatory to the dignity of women;
- (f) to value and preserve the rich heritage of our composite culture;
- (g) to protect and improve the natural environment including forests, lakes, rivers, wildlife and to have compassion for living creatures;
- (h) to develop the scientific temper, humanism and the spirit of inquiry and reform;
- (i) to safeguard public property and to abjure violence;
- (j) to strive towards excellence in all spheres of individual and collective activity so that the nation constantly rises to higher levels of endeavour and achievement;
- \*(k) who is a parent or guardian, to provide opportunities for education to his child or, as the case may be, ward between the age of six and fourteen years.

**Note:** The Article 51A containing Fundamental Duties was inserted by the Constitution (42nd Amendment) Act, 1976 (with effect from 3 January 1977).

<sup>(</sup>k) was inserted by the Constitution (86th Amendment) Act, 2002 (with effect from 1 April 2010).



### THE CONSTITUTION OF INDIA

#### PREAMBLE

WE, THE PEOPLE OF INDIA, having solemnly resolved to constitute India into a <sup>1</sup>[SOVEREIGN SOCIALIST SECULAR DEMOCRATIC REPUBLIC] and to secure to all its citizens :

**JUSTICE**, social, economic and political;

**LIBERTY** of thought, expression, belief, faith and worship;

**EQUALITY** of status and of opportunity; and to promote among them all

**FRATERNITY** assuring the dignity of the individual and the <sup>2</sup>[unity and integrity of the Nation];

**IN OUR CONSTITUENT ASSEMBLY** this twenty-sixth day of November, 1949 do **HEREBY ADOPT, ENACT AND GIVE TO OURSELVES THIS CONSTITUTION.** 

Subs. by the Constitution (Forty-second Amendment) Act, 1976, Sec.2, for "Sovereign Democratic Republic" (w.e.f. 3.1.1977)
Subs. by the Constitution (Forty-second Amendment) Act, 1976, Sec.2, for "Unity of the Nation" (w.e.f. 3.1.1977)

#### Review Team Mathematics (Class XII) Session-(2024-25)

Name	Designation	School		
Team Leader				
Mr. Sanjeev Kumar	Principal	RPVV, Kishan Ganj		
Team Members				
Mr. Vidya Sagar Malik	Lecturer Mathematics	Core Academic Unit		
Mr. Shashank Vohra	Lecturer Mathematics	RPVV, Hari Nagar		
Smt. Suman Arora	Lecturer Mathematics	RPVV, Paschim Vihar		

### ANNUAL SYLLABUS MATHEMATICS (Code NO. 041) Class-XII Session (2024-25)

The Syllabus in the subject of Mathematics has undergone changes from time to time in accordance with growth of the subject and emerging needs of the society. Senior Secondary stage is a launching stage from where the students go either for higher academic education in Mathematics or for professional courses like Engineering, Physical and Biological science, Commerce or Computer /Applications. The present revised syllabus has been designed in accordance with National Curriculum Framework 2005 and as per guidelines given in Focus Group on Teaching of Mathematics 2005 which is to meet the emerging needs of all categories of students. Motivating the topics from real life situations and other subject areas, greater emphasis has been laid on application of various concepts.

#### Objectives

The broad objectives of teaching Mathematics at senior school stage intend to help the students:

- to acquire knowledge and critical understanding, particularly by way of motivation and visualization, of basic concepts, terms, principles, symbols and mastery of underlying processes and skills.
- to feel the flow of reasons while proving a result or solving a problem.
- to apply the knowledge and skills acquired to solve problems and wherever possible, by more than one method.
- to develop positive attitude to think, analyze and articulate logically.
- to develop interest in the subject by participating in related competitions.
- to acquaint students with different aspects of Mathematics used in daily life.

- to develop an interest in students to study Mathematics as a discipline.
- to develop awareness of the need for national integration, protection of environment, observance of small family norms, removal of social barriers, elimination of genderbiases.
- to develop reverence and respect towards great Mathematicians for their contributions to the field of Mathematics.

#### ANNUAL SYLLABUS CLASS XII SUBJECT: MATHEMATICS (041) SESSION (2024-25)

#### CONTENT

#### **Unit-I: Relations and Functions**

#### **Unit-I: Relations and Functions**

#### 1. Relations and Functions

Types of relations: reflexive, symmetric, transitive and equivalence relations. One to one and onto functions.

#### 2. Inverse Trigonometric Functions

Definition, range, domain, principal value branch. Graphs of inverse trigonometric functions.

#### Unit-II: Algebra

#### 1. Matrices

Concept, notation, order, equality, types of matrices, zero and identity matrix, transpose of a matrix, symmetric and skew symmetric matrices. Operation on matrices: Addition and multiplication and multiplication with a scalar. Simple properties of addition, multiplication and scalar multiplication. On-commutativity of multiplication of matrices and existence of non-zero matrices whose product is the zero matrix (restrict to square matrices of order 2). Invertible matrices and proof of the uniqueness of inverse, if it exists; (Here all matrices will have real entries).

#### 2. Determinants

Determinant of a square matrix (up to 3 x 3 matrices), minors, co-factors and applications of determinants in finding the area of a triangle. Adjoint and inverse of a square matrix. Consistency, inconsistency and number of solutions of system of linear equations by examples, solving system of linear equations in two or three variables (having unique solution) using inverse of a matrix.

#### Unit-III: Calculus

#### 1. Continuity and Differentiability

Continuity and differentiability, chainrule, derivative of inverse trigonometric functions, *like*  $\sin^{-1}x$ ,  $\cos^{-1}x$  and  $\tan^{-1}x$ , derivative of implicit functions. Concept of exponential and logarithmic functions. Derivatives of logarithmic and exponential functions. Logarithmic differentiation, derivative of functions expressed in parametric forms. Second order derivatives.

#### 2. Applications of Derivatives

Applications of derivatives: rate of change of quantities, increasing/decreasing functions, maxima and minima (first derivative test motivated geometrically and second derivative test given as a provable tool). Simple problems (that illustrate basic principles and understanding of the subject as well as real-life situations).

#### 3. Integrals

Integration as inverse process of differentiation. Integration of a variety of functions by substitution, by partial fractions and by parts, Evaluation of simple integrals of the following types and problems based on them

$$\int \frac{dx}{x^2 \pm a^2}, \int \frac{dx}{x^2 \pm a^2}, \int \frac{dx}{\sqrt{a^2 - x^2}}, \int \frac{dx}{ax^2 + bx + c}, \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$
$$\int \frac{px + q}{ax^2 + bx + c} dx, \int \frac{px + q}{\sqrt{ax^2 + bx + c}} dx, \int \sqrt{a^2 \pm x^2} dx, \int \sqrt{ax^2 + bx + c} dx$$

Fundamental Theorem of Calculus (without proof).Basic properties of definite integrals and evaluation of definite integrals

#### 4. Applications of the Integrals:

Applications in finding the area under simple curves, especially lines, circles/parabolas/ ellipses (in standard form only)

#### 5. Differential Equations

Definition, order and degree, general and particular solutions of a differential equation. Solution of differential equations by method of separation of variables, solutions of homogeneous differential equations of first order and first degree. Solutions of linear differential equation of the type Solutions of linear differential equation of the type:

 $\frac{dy}{dx}$  + py = q, where p and q are functions of x or constant.

 $\frac{dx}{dy} + px = q$ , where p and q are functions of y or constant.

#### COMPLETION OF MID TERM SYLLABUS BY 13th September 2024 REVISION

#### **Unit-IV: Vectors and Three-Dimensional Geometry**

#### 1. Vectors

Vectors and scalars, magnitude and direction of a vector. Direction cosines and direction ratios of a vector. Types of vectors (equal, unit, zero, parallel and collinear vectors), position vector of a point, negative of a vector, components of a vector, addition of vectors, multiplication of a vector by a scalar, position vector of a point dividing a line segment in a given ratio. Definition, Geometrical Interpretation, properties and application of scalar (dot) product of vectors, vector (cross) product of vectors.

#### 2. Three - dimensional Geometry

Direction cosines and direction ratios of a line joining two points. Cartesian equation and vector equation of a line, skew lines, shortest distance between two lines. Angle between two lines.

#### **Unit-V: Linear Programming**

#### 1. Linear Programming

Introduction, related terminology such as constraints, objective function, optimization, graphical method of solution for problems in two variables, feasible and infeasible regions (bounded or unbounded), feasible and infeasible solutions, optimal feasible solutions (up to three non-trivial constraints).

#### Unit-VI: Probability

#### 1. Probability

Conditional probability, multiplication theorem on probability, independent events, total probability, Bayes' theorem, Random variable and its probability distribution, mean of random variable.

#### Note-Syllabus must be completed by 13th September 2024

#### Preparation for Pre Board Examination (2024-25)

**Pre Board Examination** 

#### BOARD EXAM 2024-25

For further Information kindly refer to CBSE guidelines

https://cbseacademic.nic.in/



# Contents

S. No.	Chapter Name	Page No.
1.	Relations and Functions	19–36
2.	Inverse Trigonometric Functions	37–52
3.	Matrices	53–70
4.	Determinants	71–88
5.	Continuity and Differentiability	89–106
6.	Application of Derivatives	107–122
7.	Integrals	123–151
3.	Application of Integrals	152–161
).	Differential Equations	162–178
10.	Vectors	179–198
11.	Three-dimensional Geometry	199–215
12.	Linear Programming	216–232
13.	Probability	233–252
	Practice Papers	253–31











[Class XII : Maths]



#### **CONCEPT MAP OF CONTINUITY AND DIFFERENTIABILTY**

**Noteworthy Results on Continuous Functions** 

\* A constant Function f(x)=k is continuous everywhere.

\* Identity Function f(x)=x is continuous everywhere.

\* Polynomial Function  $f(x) = f(x) = a_0 + a_1 x + a_2 x^2 + \dots + a_n x^n, n \in N, x \in R$  is continuous everywhere.

\* The modulus function f(x)=|x| is continuous everywhere.

\* The logarithmic function f(x)=x is continuous in its domain

\* The exponential function  $f(x) = a^x, a > 0$  is continuous everywhere.

\* The sine function f(x)=sinx and cosine function f(x)=cosx are everywhere continuous .

\*The tangent function, cotangent function, secant function and cosecant function are continuous in their respective domains.

\*All the six inverse trigonometric functions are continuous in their respective domains.

\*A rational function f(x)=g(x)/h(x), h(x) not equal to zero is continuous at every point of its domain.

\* Sum , difference ,product and quotient of of two continuous function is a continuous function.

A function f may fail to **be** continuous at x=0 for any of the following reasons

(1) f is not defined at x=a, i.e, f(a) does not exist

- (2)Either  $\lim_{x\to a^-} f(x)$  does not exist or  $\lim_{x\to a^+} f(x)$  does not exist.
- (3)  $\lim_{x \to a^-} f(x) \neq \lim_{x \to a^+} f(x)$
- (4)  $\lim_{x \to a^-} f(x) = \lim_{x \to a^+} f(x) \neq f(a)$

## APPLICATION OF DERIVATIVE





#### INTEGRATION OF SOME SPECIAL FUNCTIONS

(i) 
$$\int \frac{dx}{x^2 - a^2} = \frac{1}{2} \log \left| \frac{x - a}{x + a} \right| + c$$
 (ii)  $\int \frac{dx}{a^2 - x^2} = \frac{1}{2a} \log \left| \frac{a + x}{a - x} \right| + c$   
(iii)  $\int \frac{dx}{x^2 - a^2} = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$  (iv)  $\int \frac{dx}{\sqrt{x^2 - a^2}} = \log \left| x + \sqrt{x^2 - a^2} \right| + c$   
(v)  $\int \frac{dx}{\sqrt{a^2 - x^2}} = \sin^{-1} \frac{x}{a} + c$  (v)  $\int \frac{dx}{\sqrt{x^2 - a^2}} = \log \left| x + \sqrt{x^2 + a^2} \right| + c$ 

#### SOME STANDARD INTEGRALS

(i) 
$$\int x^{n} dx = \frac{x^{n+1}}{n+1} + c, n \neq -1$$
 like,  $\int dx = x + c$   
(ii)  $\int \cos x \, dx = \sin x + c$  (iii)  $\int \sin x \, dx = -\cos x + c$   
(iv)  $\int \sec^{2} x \, dx = \tan x + c$  (v)  $\int \csc e^{2} x \, dx = -\cot x + c$   
(vi)  $\int \sec x \tan x \, dx = \sec x + c$  (vii)  $\int \cos \sec x \cot x \, dx = -\csc x + c$   
(viii)  $\int \frac{dx}{\sqrt{1-x^{2}}} = \sin^{-1}x + c$  (ix)  $\int \frac{dx}{\sqrt{1-x^{2}}} = \cos^{-1}x + c$   
(x)  $\int \frac{dx}{1+x^{2}} = \tan^{-1}x + c$  (xi)  $\int \frac{dx}{1+x^{2}} = \cos^{-1}x + c$   
(xii)  $\int e^{x} dx = e^{x} + c$  (xiii)  $\int a^{x} dx = \frac{a^{x}}{\log a} + c$   
(xiv)  $\int \frac{dx}{x\sqrt{x^{2}-1}} = \sec^{-1}x + c$  (xv)  $\int \frac{dx}{x\sqrt{x^{2}-1}} = -\csc^{-1}x + c$   
(xvi)  $\int \frac{1}{x} dx = \log |x| + c$ 

#### **INTEGRATION BY PARTS**

$$\int f_{1}(x) f_{2}(x) dx = f_{1}(x) \int f_{2}(x) dx - \int \frac{d}{dx} f_{1}(x) \int f_{2}(x) dx$$

#### INTEGRATION BY PARTIAL FRACTIONS

A rational function of the form  $\frac{P(x)}{Q(x)}$   $(Q(x) \neq 0) = T(x) + \frac{P_1(x)}{Q(x)}$ ,  $P_1(x)$  has degree less than that

of Q(x). We can integrate  $\frac{P_1(x)}{Q(x)}$  by expressing it in the following forms:

(i) 
$$\frac{px+q}{(x-a)(x-b)} = \frac{A}{x-a} + \frac{B}{x-b}, a \neq b$$

(ii) 
$$\frac{px+q}{(x+a)^2} = \frac{A}{x-a} + \frac{B}{(x-a)^2}$$

(iii) 
$$\frac{px+qx+r}{(x+a)(x-b)(x-c)} = \frac{A}{x-a} + \frac{B}{x-b} + \frac{C}{x-c}$$

(iv) 
$$\frac{px^2 + qx + r}{(x+a)^2 (x-b)} = \frac{A}{x-a} + \frac{B}{(x-a)^2} + \frac{C}{x-b}$$

(iv) 
$$\frac{px^2 + qx + r}{(x-a)(x^2 + bx + c)} = \frac{A}{x-a} + \frac{Bx + C}{x^2 + bx + c}$$

#### INTEGRATION BY PARTIAL FRACTIONS

(i) 
$$\int \sqrt{x^2 + a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log \left| x + \sqrt{x^2 - a^2} \right| + c$$

(ii) 
$$\int \sqrt{x^2 + a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} + \frac{a^2}{2} \log \left| x + \sqrt{x^2 + a^2} \right| + c$$

(iii) 
$$\int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + c$$

#### FIRST FUNDAMENTAL THEOREM OF INTEGRAL CALCULUS

Let the area functions be defined by  $A(x) = \int_{a}^{x} f(x) dx A \ge a$ , where *f* is continuous on [*a*, *b*] then A'(x) = f(x)  $\forall x \in [0, b]$ .

#### SECOND FUNDAMENTAL THEOREM OF INTEGRAL CALCULUS

Let f be a continuous functions of x defined on [a, b] and let F be another function such that

 $\frac{d}{dx}F(x) = f(x) \forall x \in \text{domain of } f_1 \text{ then } \int_a^b f(x) dx = [F(x) + c]_a^b = F(b) - F(a). \text{ This a called the definite integral of f over the range } [a, b] \text{ where } a \text{ and } b \text{ are called the limits of integration, a being the lower limit and } b \text{ be the upper limit.}$ 







	(1, 2) The angles made by $\overrightarrow{OP}$ with positive direction of x, y & respectively) are called is direction angles, and the cosine value of these angles is direction angles, and the cosine value of these angles i.e. $\cos \alpha$ , $\cos \beta$ & $\cos \gamma$ are called direction cosines of $\overrightarrow{OP}$ alled direction cosines of $\overrightarrow{OP}$ denoted by 1, m & n respectively. <i>Joining Two Point s</i> $(x_{1}, y_{1}, z_{1}) \& B(x_{2}, y_{2}, z_{2}) be any two point s$ $acc, then \overrightarrow{OA} = x_{1}^{2} + y_{1}^{2} + z_{1}^{2} \& \&$ $\frac{1}{2}(x_{2} - x_{1})^{2} + (y_{2} - y_{1})^{2} + (z_{2} - z_{1})^{2}$ <i>ECTION FORMULAE</i> <i>he position vectors are</i> $\vec{a} \& \vec{b}$ respectively, <i>the ratio m</i> : n <i>Internally is given by</i> $\frac{mb + n\vec{a}}{m - n}$ <i>in Externally is given by</i> $\frac{mb + n\vec{a}}{m - n}$ <i>in Externally is given by</i> $\frac{mb + n\vec{a}}{m - n}$ <i>in even by</i> $\frac{\vec{a} + \vec{b}}{2}$ .
<u>CONCEPT – MAP (CHAPTER – 10)</u> <u>VECTORS</u>	If a point P in space, having coordinates $(x, y)$ with respect to the origin $O(0, 0, 0)$ . Then, the vector $\overrightarrow{OP}$ having $O$ and P as its initial & terr points, respectively, is called the position vector $\overrightarrow{OP}$ the point P with respect to $O$ . Using distance formula, the magnitude of $\overrightarrow{OP}$ $(or \overrightarrow{r})$ is given by $ \overrightarrow{OP}  = \sqrt{x^2 + y^2 + z^2}$ $(or \overrightarrow{r})$ is given by $ \overrightarrow{OP}  = \sqrt{x^2 + y^2 + z^2}$ $(\overrightarrow{or} \overrightarrow{r})$ is given by $ \overrightarrow{OP}  = \sqrt{x^2 + y^2 + z^2}$ $(\overrightarrow{aB} = \frac{1}{2} + 4C)$ $(\overrightarrow{aB} = \frac{1}{2} + \frac{1}$
	A quantify that has magnitude as well as direction is called a vector.       Image: Complete the section is called a vector.         A directed line       Image: Complete the section is a vector denoted as AB or simply as a, and read as 'vector AB' or 'vector a'.         TYPES OF VECTORS       Image: Complete the section of the vector is a vector when the magnitude of the vector coincides with the termin al point.         A vector which has a magnitude of unit length is called a <u>unit vector</u> .       Image: Complete the same starting point of the vector same starting point of the vector same with the termin al point.         A vector which has a magnitude of unit length is called a <u>unit vector</u> .       Image: Complete the same starting point are called <u>complete</u> .         Two or more vectors are said to be <u>equal</u> when their magnitude is equal and also their magnitude is the same.       Image: the vector same the same starting bo int exector same the same their magnitude but exactly opposite in direction the vector same negative of a vector same negative of a vector same negative of each other.



[Class XII : Maths]




# CHAPTER-1

# **RELATIONS AND FUNCTIONS**



By looking at the two thermometers shown, you can make some general comparisons between the scales. For example, many people tend to be comfortable in outdoor temperatures between 50°F and 80°F (or between 10°C and 25°C). If a meteorologist predicts an average temperature of 0°C (or 32°F), then it is a safe bet that you will need a winter jacket.

Sometimes, it is necessary to convert a Celsius measurement to its exact Fahrenhelt measurement or vice versa.

For example, what if you want to know the temperature of your child in Fahrenheit, and the only thermometer you have measures temperature in Celsius measurement? Converting temperature between the systems is a straightforward process. Using the function

$$F = f(C) = \frac{9}{5}C + 32$$
, any temperature in Celsius can be converted into Fahrenheit scale.



If number of elements in set A and set B are p and q respectively, Means n(A) = p, n(B) = q, then

No. of Relation af A × A =  $2^{p^2}$ No. of Relation of B × B =  $2^{q^2}$ No. of Relation of A × B = No. of Relation of B × A =  $2^{pq}$ No. of **NON EMPTY** Relation of A × A =  $(2^{p^2} - 1)$ , No. of **NON EMPTY** Relation of B × B =  $(2^{q^2} - 1)$ .

No. of **NON-EMPTY** Relation of A × B = No. of Relation of B × A =  $(2^{pq} - 1)$ 

**Q.1** If A = {*a, b, c*} and B = {1, 2} find the number of Relation R on (i) A × A (ii) B × B (iii) A × B

**Ans.** As *n*(A) = 3, *n*(B) = 2, so

No. of Relation R on  $A \times A = 2^{3 \times 3} = 2^9 = 512$ 

No. of Relation R on B × B =  $2^{2 \times 2}$  =  $2^4$  = 16

No. of Relation R on  $A \times B = 2^{3 \times 2} = 2^8 = 64$ 

- **Q.2**  $A = \{d, o, e\}$  and  $B = \{22, 23\}$  find the number of Non-empty Relation R on (i)  $A \times A$  (ii)  $B \times B$
- **Ans.** As *n*(A) = 3, *n*(B) = 2, so

No. of Relation Non-empty relations R on A × A =  $2^{3\times3} - 1 = 2^9 - 1 = 511$ 

No. of Relation Non-empty R on  $B \times B = 2^{2 \times 2} - 1 = 2^4 - 1 = 15$ 

# **Different types of relations**

# Empty Relation Or Void Relation

A relation R in a set A is called an empty relation, if no element of A is related to any element of A and we denote such a relation by  $\phi$ .

**Example:** Let  $A = \{1, 2, 3, 4\}$  and let R be a relation in A, given by  $R = \{(a, b): a + b = 20\}$ .

# Universal Relation

A relation R in a set A is called an universal relation, if each element of A is related to every element of A.

**Example:** Let A =  $\{1, 2, 3, 4\}$  and let R be a relation in A, given by R =  $\{(a, b): a + b > 0\}$ .

# Identity Relation

A relation R in a set A is called an identity relation, where R = {(a, a),  $a \in A$ }.

Example : Let A =  $\{1, 2, 3, 4\}$  and let R be a relation in A, given by R =  $\{(1, 1), (2, 2), (3, 3), (4, 4)\}$ .

#### Reflexive Relation

A relation R in a set A is called a Reflexive relation, if  $(a, a) \in R$ , for all  $a \in A$ .

Example : Let A = {1, 2, 3, 4} and let R be a relation in A, given by

 $R = \{(1, 1), (2, 2), (3, 3), (4, 4)\}.$ 

 $\mathsf{R} = \{(1, 1), (2, 2), (3, 3), (4, 4), (1, 2)\}.$ 

 $\mathsf{R} = \{(1, 1), (2, 2), (3, 3), (4, 4), (2, 3), (1, 3), (3, 1)\}.$ 

#### Symmetric Relation

A relation R in a set A is called a symmetric relation, if  $(a, b) \in R$ , then  $(b, a) \in R$  for all  $a, b \in A$ .

Example : Let  $A = \{1, 2, 3, 4\}$  and let R be a relation in A, given by

 $R = \{(1, 1), (2, 2), (3, 3)\}.$ 

 $R = \{(1, 2), (2, 1), (3, 3)\}.$ 

 $R = \{(1, 1), (2, 2), (3, 3), (4, 4), (2, 3), (1, 3), (3, 1), (3, 2)\}.$ 

# Transitive Relation

A relation R in a set A is called a transitive relation,

if  $(a, b) \in \mathbb{R}$  and  $(b, c) \in \mathbb{R}$  then  $(a, c) \in \mathbb{R}$  for all  $a, b, c \in \mathbb{A}$ 

Or

 $(a, b) \in \mathbb{R}$  and  $(b, c) \notin \mathbb{R}$  for all  $a, b, c \in \mathbb{A}$ 

Example : Let {1, 2, 3, 4} and let R be a relation in A, given by

 $R = \{(1, 1), (2, 2), (3, 3)\}.$  (According to second condition)

 $R = \{(1, 2), (2, 1), (1, 1), (2, 2)\}.$  (According to first condition)

 $\mathsf{R} = \{(2, 3), (1, 3), (3, 1), (3, 2), (3, 3), (2, 2), (1, 1)\}.$ 

#### Equivalence Relation

A relation R in a set A is said to be an equivalence relation if it is reflexive, symmetric and transitive.

#### Illustration:

Let A be the set of all integers and let R be a relation in A, defined by  $R = \{a, b\}$ :  $a = b\}$ , Prove that R is Equivalence Relation.

**Solution:** Reflexivity : Let R be reflexive  $\Rightarrow$  (*a*, *a*)  $\in$  R  $\forall$  *a*  $\in$  A

```
\Rightarrow a = a, which is true
```

Thus, R is Reflexive Relation.

**Symmetricity :** Let  $(a, b) \in \mathbb{R} \forall a, b \in \mathbb{A}$ 

- $\Rightarrow$  a = b
- $\Rightarrow$  b = a

so  $(b, a) \in R$ . Thus R is symmetric Relation.

**Transitivity :** Let  $(a, b) \in \mathbb{R}$  and  $(b, c) \in \mathbb{R} \ \forall a, b, c \in \mathbb{A}$ 

- $\Rightarrow$  a = b and b = c
- $\Rightarrow$  a = b = c
- $\Rightarrow$  a = c

so  $(a, c) \in \mathbb{R}$ . Thus R is transitive Relation.

As, R is reflexive, Symmetric and transitive Relation

∴ R is an Equivalence Relation

#### **FUNCTIONS**

Functions can be easily defined with the help of concept mapping. Let X and Y be any two non-empty sets. "A function from X to Y is a rule or correspondence that assigns to each element of set X, one and only one element of set Y". Let the correspondence be '*f* then mathematically we write  $f: X \to Y$ .

where y = f(x), x  $\varepsilon$  X and  $y \varepsilon$  Y. We say that 'y' is the images of 'x' under f (or x is the pre image of y).

- A mapping *f* : X → Y is said to be a function if each element in the set X has its image in set Y. It is also possible that there are few elements in set Y which are not the images of any element in set X.
- Every element in set X should have one and only one image. That means it is impossible to have more than one image for a specific element in set X.
- Functions cannot be multi-valued (A mapping that is multi-valued is called a relation from X and Y) eg.



#### Testing for a function by Vertical line Test

A relation  $f : A \rightarrow B$  is a function or not, it can be checked by a graph of the relation. If it is possible to draw a vertical line which cuts the given curve at more that none point then the given relation is not a function and when this vertical line means line parallel to Y-axis cuts the curve at only one point then it is a function. Following figures represents which is not a function and which is a function.



NOT FUNCTION NOT A

NOT A FUNCTION

### **Number of Functions**

Let X and Y be two finite sets having m and n elements respectively. Thus each element of set X can be associated to any one of n elements of set Y. So, total number of functions from set X to set Y is  $n^m$ .

Real valued function: if R, be the set of real numbers and A, B are subsets of R, then the function f :  $A \rightarrow B$  is called a real function or real valued functions.

### Domain, Co-Domain And Range of Function

If a function f is defined from a set A to set B then (if :  $A \rightarrow B$ ) set A is called the domain of f and set B is called the co-domian of f.

The set of all *f*-images of the elements of A is called the range of *f*.

In other words, we can say

Domain = All possible values of x for which f(x) exists.

Range = For all values of , all possible values of f(x).



From the figure we observe that

Domain = A = {*a*, *b*, *c*, *d*}

Range = {*p*, *q*, *r*}, Co-Domain = {*p*, *q*, *r*, *s*} = B

# EQUAL FUNCTION

Two function f and g are said to be equal functions, if and only if

- (i) Domain of f = Domain of g
- (ii) Co-domain of f = Co-domain of g
- (iii) f(x) = g(x) for all  $x \in$  their common domain

#### **TYPES OF FUNCTION**

One-one function (injection): A function  $f : A \rightarrow B$  is said to be a one-one function or an injection, if different elements of A have different images in B.

e.g. Let  $f : A \rightarrow B$  and  $g : X \rightarrow Y$  be two functions represented by the following diagrams.



Clearly,  $f : A \rightarrow B$  is a one-one function. But  $g : X \rightarrow Y$  is not one-one function because two distinct elements  $x_1$  and  $x_2$  have the same image under function g.

#### Method to check the injectivity (One-One) of a function

- (i) Take two arbitrary elements x, y (say) in the domain of f.
- (ii) Solve f(x) = f(y). If f(x) = f(y) give x = y only, then  $f : A \to B$  is a one-one function (or an injection). Otherwise not.

If function is given in the form of ordered pairs and if two ordered pairs do not have same second element then function is one-one.

If the graph of the function y = f(x) is given and each line parallel to x-axis cuts the given curve at maximum one point then function is one-one. (Strictly increasing or Strictly Decreasing Function). E.g.



Number of one-one functions (injections) : If A and B are finite sets having *m* and *n* elements respectively, then number of one-one functions from A and B =  ${}^{n}P_{m}$  is  $n \ge m$  and 0 if n < m.

If f(x) is not one-one function, then its Many-one function.

**Onto function (surjection) :** A function  $f : A \rightarrow B$  is onto if each element of B has its preimage in A. In other words, Range of f = Co-domain of f. *e.g.* The following arrow-diagram shows onto function.



Number of onto function (surjection): If A and B are two sets having m and n elements

respectively such that  $1 \le n \le m$ , then number of onto functions from A to B is  $\sum_{r=1}^{n} (-1)^{n-r} C_r$ 

. **r**<sup>m</sup>

Into function: A function  $f: A \rightarrow B$  is an into function if there exists an element in B having no pre-image in A.

In other words,  $f: A \rightarrow B$  is an into function if it is not an onto function e.g, The following arrow diagram shows into function.



# Method to find onto or into function:

- (i) Solve f(x) = y by taking x as a function of y i.e., g(y)(say).
- (i) Now if g(y) is defined for each  $y \varepsilon$  co-domain and  $g(y) \varepsilon$  domain then f(x) is onto and if any one of the above requirements is not fulfilled, then f(x) is into.

**One-one onto function (bijection) :** A function  $f: A \rightarrow B$  is a bijection if it is one-one as well as onto.

In other words, a function  $f: A \rightarrow Bis$  a bijection if

(i) It is one-one ie.,  $f(x) = f(y) \Rightarrow x = y$  for all  $x, y \in A$ .

(ii) It is onto i.e., for all  $y \in B$ , there exists  $x \in A$  such that f(x) = y.

Clearly, *f* is a bijection since it is both injective as well as surjective.

# Illustration :

Let  $f: \mathbb{R} \to \mathbb{R}$  be defined as f(x) = 7x - 5, then show that function is one-one and onto Both.

**Solution** : Let  $f(x) = f(y) \forall x, y \in \mathbb{R}$ 

$$\Rightarrow$$
 7x-5=7y-5

$\Rightarrow$ )	x = y, so	f(x) is one-one function	

Now, As f(x) = 7x - 5, is a polynomial function.

so it is defined everywhere. Thus, Range = R

As, Range = co-domain, so f is onto function.

Alternative method : Graph of f(x) is a line which is strictly increasing for all values of x, so its one-one function and Range of f(x) is R which is equal to R so onto function.

# ILLUSTRATION:

If  $f: X \to Y$  is defined, then show that f is neither one-one nor onto function.



**Solution :** As for elements 3 and 4 from set X we have same image *c* in set Y, so *f* is not one-one function.

Further element d has no pre -image in set X,

so f is not onto function

# ILLUSTRATION:

Prove that the function  $f : \mathbb{N} \to \mathbb{N}$ , defined by  $f(x) = x^2 + x + 2022$  is one-one. **SOLUTION :** APPROACH-I

Let 
$$f(x_1) = f(x_1) \forall x_1, x_2 \in N \Rightarrow = x_1^2 + x_1 + 2022 = x_2^2 + x_2 + 2022$$
  
 $\Rightarrow x_1^2 + x_1 = x_2^2 + x_2$   
 $\Rightarrow (x_1^2 - x_2^2) + (x_1 - x_2) = 0$   
 $\Rightarrow (x_1 - x_2) + (x_1 + x_2 + 1) = 0$   
Thus, $(x_1 - x_2) = 0$  as  $(x_1 + x_2 + 1) \neq 0 \forall x_1, x_2 \in N$   
so, f is ONE-ONE function  
**APPROACH-II**  
 $f(x) = x^2 + x + 2022 \Rightarrow f'(x) = 2x + 1$   
As,  $x \in N$  so,  $2x + 1 > 0 \Rightarrow f'(x) = 0$  (Strictly Increasing function)  
so, f is ONE-ONE function

Name of Function	Definition	Domain	Range	Graph
1. Identify Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) = x \forall x \in \mathbb{R}$	R	R	X V O X
2. Constant Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) = c  \forall x \in \mathbb{R}$	R	(c)	y = c $c  f(x) = c$ $x = 0$ $x$
3. Polynomial Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) = p_0 + p_1 x + p_2 x^2 + + p_n x^n$ , where $n \in \mathbb{N}$ and $p_0, p_1, p_2, p_n$ $\in \mathbb{R} \ \forall \ x \in \mathbb{R}$			
4. Rational Function	The function <i>f</i> defined by $f(x) = \frac{P(x)}{Q(x)}$ , where P(x) and Q(x) are polynomial functions, Q(x) $\neq 0$			
5. Modulus Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) =  x  = \begin{cases} x, & x \ge 0 \\ -x, & x < 0 \end{cases}  \forall \ x \in \mathbb{R}$	R	<b>[0, ∞)</b>	The states
6. Signum Function	The function $f : \mathbb{R} \to \mathbb{R}$ defined by $f(x) = \begin{cases} \frac{ x }{x}, & x \neq 0\\ 0, & x = 0 \end{cases} = \begin{cases} -1, & x < 0\\ 1, & x > 0\\ 0, & x = 0 \end{cases}$	R	{1,0,1}	(0,1)
7. Greatest Integer Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) =  x  = \begin{cases} x, x \in \mathbb{Z} \\ \text{integer less than} \\ equal \text{ to } x, x \notin \mathbb{Z} \end{cases}$	R	Z	
8. Linear Function	The function $f: \mathbb{R} \to \mathbb{R}$ defined by $f(x) = mx + c, x \in \mathbb{R}$ where <i>m</i> and <i>c</i> are constants	R	R	

Type of Functions

#### **ONE-MARK QUESTIONS**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE

1.	Consider the se	et A =	= {1, 2, 3}, then wr	ite sn	nallest equivalence re	latio	n on A.
	(a) {}	(b)	{(1, 1)}	(c)	{(1, 1), (2, 2), (3, 3)}	(d)	{(3, 3)}
2.	Consider the s from A onto itse	et A elf are	containing 5 elem e	ients,	then the total numbe	r of i	njective functions
	(a) 5	(b)	25	(c)	120	(d)	125
3.	Let Z be the se divisible by 4, t	t of i hen l	ntegers and R be R partitions the se	the re et Z in	elation defined in Z su to how many Pairwise	ch th e disj	at <i>aRb</i> if ( <i>a</i> – <i>b</i> ) is oint subsets.
	(a) 2	(b)	3	(c)	4	(d)	5
4.	If A = {d, 0, e} t	nen t	he number of rela	itions	on A × A are		
	(a) 3	(b)	8	(c)	15	(d)	512
5.	If A = {2023, 20	24) t	hen the number o	f non	empty relations on A	×Aa	re
	(a) 1	(b)	4	(c)	15	(d)	16
6.	If A = {2023, 20	24) t	hen the number o	f Refl	exive relations on A × .	Aare	9
	(a) 2	(b)	4	(c)	8	(d)	16
7.	If A = {s, u, v}, t	hen	the number of Syr	nmet	ric relations on A × A a	are	
	(a) 8	(b)	9	(c)	32	(d)	64
8.	Let A be the se number of refle	et of xive	the Letters of the relations on A × A	nam	e of our country the '	ʻIND	IA". Then find the
	(a) 4096	(b)	2048	(c)	1024	(d)	16
9.	Let A = $\{x : x^2 < x^2$	: 3, x	$x \in W$ }, then the number of the matrix $x \in W$	umbe	r of Symmtric relation	is on	A × A are
	(a) 1	(b)	2	(c)	4	(d)	8
10.	If there are 'p' e then p =	leme	ents in set A, such t	hat n	umber of Reflexive rela	ation	on A × A are 4096,
	(a) 4	(b)	6	(c)	8	(d)	12
11.	Let A = {d, 0, e	}, the	en Find 'p' if the nu	ımbei	r of Symmetric relatio	ns oi	n A × A are 2º.
	(a) 4	(b)	6	(c)	8	(d)	12
12.	Find the maxim	num	number of equival	lence	relations on the set A	= {1	, 2, 3}.
	(a) 3	(b)	5	(c)	8	(d)	9
					× <sup>2</sup>		
13.	If the function f	: R-	$- \{1, -1\} \rightarrow A \text{ def}$	ined	by $f(x) = \frac{x}{1+x^2}$ is S	Surje	ctive, then A =
	(a) R	(b)	$R - \{1, -1\}$	(c)	[0, 1)	(d)	<b>[</b> 0, ∞ <b>]</b>
14.	The number of	inje	ctions possible fro	m A =	= {1, 2, 3, 4} to B = {5,	6, 7}	are
	(a) 0	(b)	3	(c)	6	(d)	12

15. If the number of one-one functions that can defined from A =  $\{4, 8, 12, 16\}$  to B is 5040, then n(B) =

16. If the function  $f : \mathbb{R} \to \mathbb{A}$  defined by  $f(x) = 3 \sin x + 4\cos x$  is Surjective, then  $\mathbb{A} =$ 

(a) [-7, 7] (b) [-1, 1] (c) [1, 7] (d) [-5, 5]

17. The Part of the graph of a Non-Injective function  $f: \mathbb{R} \to \mathbb{R}$  ange defined by  $f(x) = x^2 - 2x + a$  is given below. If the domain of f(x) is modified as either  $(-\infty, b]$  or  $[b, \infty]$  then f(x) becomes the Injective function. What must be the value of (b - a).

18. The graph of the function  $f: \mathbb{R} \to A$  defined by y = f(x) is given below, then find A such that function f(x) is onto function





# ASSERTION-REASON BASED QUESTIONS (Q.19 & Q.20)

In the following questions, a statement of assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

(a) Both A and R are true and R is the correct explanation of A.

- (b) Both A and R are ture but R is not the correct explanation of A.
- (c) A is true but R is false
- (d) A is false but R is true
- 19. ASSERTION (A) : A relation R = {(a, b) : |a b| < 1} defined on the set A = {1, 2, 3, 4} is Reflexive

Reason (R) : A realtion R on the set A is said to be reflexive if for  $(a, b) \in R$ 

&  $(b, c) \in \mathbb{R}$ , we have  $(a, c) \in \mathbb{R}$ .

20. Assertion (A) : A function  $f : \mathbb{R} \to \mathbb{R}$  given f(x) = |x| is one-one function.

Reason (R) : A function  $f : A \rightarrow B$  is said to be Injective if

$$f(a) = f(b) \Rightarrow a = b$$

#### **TWO MARKS QUESTIONS**

- 21. If A = {a, b, c, d} and f = {(a, b), (b, d), (c, a), (d, c)}, show that f is one-one from A to A.
- 22. Show that the relation R on the set of all real numbers defined as  $R = \{(a, b) : a \le b^3\}$  is not transitive.
- 23. If the function  $f: \mathbb{R} \{1, -1\} \rightarrow \mathbb{A}$  defined by  $f(x) = \frac{x^2}{1 x^2}$ , is Surjective, then find A.
- 24. Give an example to show that the union of two equivalence relations on a set A need not be an equivalence relation on A.
- 25. How many reflexive relations are possible in a set A whose (A) = 4. Also find How many symmetric relations are possible on a set B whose n(B) = 3.
- 26. Let W denote the set of words in the English dictionary. Define the relation R by R {(x, y)  $\varepsilon$  W × W such that x and y have at least one letter in common). Show that this relation R is reflexive and symmetric, but not transitive.
- 27. Show that the relation R in the set of all real numbers, defined as  $R = \{(a, b): a \le b^2\}$  is neither reflexive Nor symmetric.
- 28. Consider a function  $f: \mathbb{R}_+ \to (7, \infty)$  given by  $f(x) = 16x^2 + 24x + 7$ , where  $\mathbb{R}+$  is the set of all positive real numbers. Show that function is one-one and onto both.
- 29. Let L be the set of all lines in a plane. A relation R in Lis given by R {( $L_1$ ,  $L_2$ ):  $L_1$  and  $L_2$  intersect at exactly one point,  $L_1$ ,  $L_2 \in L$ }, then show that the relation R is symmetric Only.
- 30. Show that a relation R on set of Natural numbers is given by R = {(x, y): xy is a square of an integer} is Transitive.

#### **THREE MARKS QUESTIONS**

- 31. Are the following set of ordered pairs functions? If so, examine whether the mapping is injective or surjective.
  - (i)  $\{(x, y) : x \text{ is a person}, y \text{ is the mother of } x\}$ .

- (ii)  $\{(a, b): a \text{ is a person}, b \text{ is an ancestor of } a\}$ .
- 32. Show that the function  $f: \mathbb{R} \to \mathbb{R}$  defined by  $f(x) = \frac{x^2}{x^2 + 1}$ ;  $\forall x \in \mathbb{R}$ , is neither one-one nor

onto.

- 33. Let R be the set of real numbers and  $f: R \rightarrow R$  be the function defined by f(x) = 4x + 5. Show that *f* is One-one and onto both.
- 34. Show that the relation R in the set A =  $\{3, 4, 5, 6, 7\}$  given by R =  $\{(a, b) : |a b| \text{ is divisible by 2}\}$  is an equivalence relation. Show that all the elements of  $\{3, 5, 7\}$  are related to each other and all the ements of  $\{4, 6\}$  are related to each other, but no element of  $\{3, 5, 7\}$  is related to any element  $\{4, 6\}$ .
- 35. Check whether the relation R in the set Z of integers defined as  $R = ((a, b) : a + b is "divisible by 2"}$  is reflexive, symmetric, transitive or Equivalence.
- 36. Show that that following Relations R are equivalence relation in A.
  - (a) Let A be the set of all triangles in a plane and let R be a relation in A, defined by  $R = \{(T_1, T_2) : T_1, \text{ is congruent } T_2\}$
  - (b) Let A be the set of all triangles in a plane and let R be a relation in A, defined by  $R = \{(T_1, T_2) : T_1, \text{ is similar } T_2, \}$
  - (c) Let A be the set of all lines in xy-plane and let R be a relation in A, defined by  $R = \{(L_4, L_2) : L_4, \text{ is parallel to } L_2\}$
  - (d) Let A be the set of all integers and let R be a relation in A, defined by  $R = \{(a, b) : (a b) \text{ is even}\}$
  - (e) Let A be the set of all integers and let R be a relation in A, defined by R = {(a, b) : |a - b| is a multiple of 2}
  - (f) Let A be the set of all integers and let R be a relation in A, defined by  $R = \{(a, b) : |a b| \text{ is a divisible by 3}\}$
- 37. Check whether the following Relations are Reflexive, Symmetric or Transitive.
  - (a) Let A be the set of all lines in *xy*-plane and let R be a relation in A, defined by  $R = \{(L_1, L_2) : L_1 \text{ is perpendicular to } L_2\}$
  - (b) Let A be the set of all real numbers and let R be a relation in A defined by R = { $(a, b): a \le b$ }
  - (c) Let A be the set of all real numbers and let R be a relation in A defined by R = {(a, b):  $a \le b^2$ }
  - (d) Let A be the set of all real numbers and let R be a relation in A defined by R =  $\{(a, b) : a \le b^3\}$
  - (e) Let A be the set of all natural numbers and let R be a relation in A defined by

R = (a, b) : a is a factor b

R {(a.b): b is divisible by a}

- (f) Let A be the set of all real numbers and let R be a relation in A defined by R {(a.b): (1+ ab) > 0}
- 38. Let S be the set of all real numbers. Show that the relation  $R = \{a, b\}$ :  $a^2 + b^2 = 1\}$  is symmetric but neither reflexive nor transitive.

OR

- 39. Check whether relation R defined in R as R = {*a*, *b*):  $a^2 4ab + 3b^2 = 0$ , *a*, *b*  $\in$  R} is reflexive, symmetric and transitive.
- 40. Show that the function  $f: (-\infty, 0) \rightarrow (-1, 0)$  defined by  $f(x) = \frac{x}{1+|x|}$ ,  $x \in (-\infty, 0)$  is one-one and onto.

#### **FIVE MARKS QUESTIONS**

- 41. For real numbers x and y, define x R y if and only if  $x y + \sqrt{2}$  is an irrational number. Then check the reflexivity, Symmetricity and Transitivity of the relation R.
- 42. Determine whether the relation R defined on the set of all real numbers as

 $R = \{(a, b) : a, b \in R \text{ and } a - b + \sqrt{3} \in S\}$ 

(Where S is the set of all irrational Numbers) is reflexive, symmetric or transitive.

- 43. Let N be the set of all natural numbers and let R be a relation on N × N, defined by Show that R is an equivalence relation.
  - (i)  $(a, b) R (c, d) \Leftrightarrow a + d = b + c$
  - (ii)  $(a, b) R (c, d) \Leftrightarrow ad = bc$

(iii) (a, b) R (c, d) 
$$\Leftrightarrow \frac{1}{a} + \frac{1}{d} = \frac{1}{b} + \frac{1}{c}$$

(iv) 
$$(a, b) R (c, d) \Leftrightarrow ad (b + c) = bc (a + d)$$

- 44. Let A = R {1}, f: A  $\rightarrow$  A is a mapping defined by  $f(x) = \frac{x-2}{x-1}$ , show that f is one-one and onto.
- 45. Let  $f: \mathbb{N} \to \mathbb{R}$  be a function defined as  $f(x) = 4x^2 + 12x + 15$ . Show that  $f: \mathbb{N} \to \mathbb{S}$ , where S is the range of *f*, is One-One and Onto Function.

#### **CASE STUDIES**

A. A person without family is not complete in this world because family is an integral part of all of us Human deings are considered as the social animals living in group called as family. Family plays many important roles throughout the life.

Mr. D.N. Sharma is an Honest person who is living happily with his family. He has a son Vidya and a Daughter Madhulika. Mr. Vidya has 2 sons Tarun and Gajender and a daughter Suman while Mrs. Madhulika has 2 sons Shashank and Pradeep and 2 daughters Sweety and Anju. They all Lived together and everyone shares equal responsibilities

within the family. Every member of the family emotionally attaches to each other in their happiness and sadness. They help each other in their bad times which give the feeling of security.

A family provides love, warmth and security to its all members throughout the life which makes it a complete family. A good and healthy family makes a good society and ultimately a good society involves in making a good country.



On the basis of above information, answer the following questions:

Consider Relation R in the set A of members of Mr. D. N. Sharma and his family at a particular time

- (a) If  $R = \{(x, y) : x \text{ and } y \text{ live in the same locality}\}$ , then R show that R is reflexive Relation.
- (b) If R = {(x, y) : x is exactly 7 cm taller than y}, then R show that R is not Symmetric relation.
- (c) If  $R = \{(x, y) : x \text{ is wife of } y\}$ , then show that R is Transitive only.



B. Let A be the Set of Male members of a Family, A = (Grand father, Father, Son) and B be the set of their 3 Cars of different Models, B = {Model 1, Model 2, Model 3}



On the basis of The above Information, answer the following questions:

- (a) If m & n represents the total number of Relations & functions respectively on  $A \times B$ , then find the value of (m + n).
- (b) If p & q represents the total number of Injective function & total numbers of Surjective functions respectively on A × B, then find the value of |p q|.
- C. An organization conducted bike race under two different categories—Boys & Girls. There were 28 participants in all. Among all of them finally three from category 1 and two from category 2 were selected for the final race. Ravi forms two sets B and G with

these Participants for his college Project.

Let B = {b1, b2, b3} and G = {g1, g2}, represents the set of Boys selected & G the set of Girls selected for the final race.



- (a) How many relations are possible from B to G?
- (b) Among all possible relations form B to G, how many functions can be formed from B to G?
- (c) Let  $R:B \rightarrow B$  be defined by

 $R = \{(x, y) : x \& y \text{ are students of same sex}\}$ . Check R is equivalence Relation.

OR

A function f : B  $\rightarrow$  G be defined by  $f = \{(b_1, g_1), (b_2, g_2), (b_3, g_1)\}$ 

Check if f is bijective. Justify your answer.

#### **SELF ASSESSMENT-1**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE COR-**RECTALTERNATIVE:**

- 1. Consider the set A = {1. 2, 3} and R be the smallest equivalence relation on A.then R =
  - (b) {(1,1), (2,2)} (a) {(1.1)}
  - (c)  $\{(1,1),(2,2),(3,3)\}$
- 2. Consider the set A containing n elements. Then, the total number of injective functions from A onto itself is
  - (a) 2<sup>n</sup> (b) *n*
  - (c) n (d) n!
- 3. The total number of injective nappingsfrom a set with m elements to a set with n elements,  $m \le n$  is
  - (a) n! (b) *n*<sup>*m*</sup>

(d) (c)  $m^n$ [Class XII : Maths]

34

- 4. The number of injections possible from A =  $\{1,3,5,6\}$  to B =  $\{2,8,11\}$  is
  - (a) 12 (b) 22
  - (c) 3 (d) 0
- 5. The number of one-one functions that can defined from

A = {4,8,12,16} to B is 5040, then n(B) =

- (a) 7 (b) 8
- (c) 9 (d) 10

#### SELF ASSESSMENT-2

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT CHOOSE THE COR-RECT ALTERNATIVE.

- 1. A relation R in a set A is called ..... if  $(a_1, a_2) \in R$  implies  $(a_2, a_1) \in R$ , for all  $a_1, a_2 \in R$ 
  - Α.
  - (a) Reflexive (b) Symmetric
  - (c) Transitive (d) Equivalence

2. Let  $f: \mathbb{R} - \{0\} \to \mathbb{R} - \{0\}$  be defined by  $f(x) = \frac{1}{x} \forall x \in \mathbb{R}$ . Then *f* is

- (a) One-One (b) Many-One
- (c) Not defined (d) None of these
- 3. Let  $P = \{(x, y) | x^2 + y^2 = 1, x, y \in R\}$ . Then P is
  - (a) Reflexive (b) Symmetric
  - (c) Transitive (d) Equivalence
- 4. The function  $f: \mathbb{R} \to \mathbb{R}$  defined by f(x) = [x], where [.] is greatest integer function is
  - (a) One-One (b) Many-One
  - (c) Onto (d) None of these
- 5. The number of bijective functions (One-one and onto both) from set A to itself when A contains 2022 elements is
  - (a) 2022 (B) 2022!
  - (C) 2022<sup>2</sup> (D) 2022<sup>2022</sup>

# ANSWER

# **One Mark Questions**

1.	(c){	(1,1), (2,2),	(3,3	3)}	2.	(c) 120		3.	(c) 4
4.	(d) 5	512			5.	(c) 15		6.	(b)4
7.	(d)6	64			8.	(a) 4096		9.	(d) 8
10	. (a) 4	Ļ			11.	(b) 6		12.	(b) 5
13	. (c)[	0,1)			14.	(a) 0		15.	(d) 10
16	. (d)[	-5,5]			17.	(a) 6		18.	(a)[-1,5]
19	. (C)				20.	(d)			
Ai	s true	e but R is fa	alse		A is	false but R is	true		
					Two	Mark Quest	ions		
23	. A =	R – [–1, 0)							
25	. Ref	exive Rela	tions	s = 4096 Sy	/mme	etric Relation =	64		
					Thre	e Mark Ques	tions		
31	. (a)	Yes it's fur	nctic	on, Not Inje	ctive	but Surjective	(b) No, i	its n	ot a function
35	. EQI	JIVALENCE	ERE	LATION					
37	. (a)	Symmetrie	c		(b)	Reflexive and	Transitiv	/e	
	(c)	Neither Re	flex	ive, Symme	etric I	nor Transitive			
	(d)	Neither Re	flex	ive, Symme	etric I	nor Transitive			
	(e)	<b>Reflexive</b> a	and <sup>-</sup>	Transitive					
	(f)	Reflextive	and	Symmetric					
39	. Refl	exive only							
				F	our/l	ive Mark Que	estions		
41	. Refl	exive only			42.	Reflexive only	/		
				CASE	STU	DIES BASED	QUESTI	ON	
Β.	(a) 5	12 +27 =53	9		B. (	b) 0			
C.	(a) 6	4							
	(b) 8								
	(c) R	is an Equiv	/ale	nce Relatic	on OF	R (c) f is not Bij	ective		
					SEL	F ASSESSME	NT-1		
1.	(c)		2.	(d)		3. (d)	4.	(d	)
					SEL	F ASSESSME	NT-2		
1.	(b)		2.	(a)		3. (b)	4.	(b	)

5. (d)

5. (b)

# **CHAPTER-2**

# INVERSE TRIGONOMETRIC FUNCTIONS



An example of people using inverse trigonometric functions would be builders such as construction workers, architects, and many others.

An example of the use would be the creation of bike ramp. You will have to find the height and the length. Then find the angle by using the inverse of sine. Put the ength over the height to find the angle. Architects would have to calculate the angle of a bridge and the supports when drawing outlines. These calculations are then applied to find the safest angle. The workers would then uses these calculations to build the bridge.

# TOPIC TO BE COVERED AS PER CBSE LATEST CURRICULUM (2024-25)

- Definition, range, domain, principal value branch.
- Graphs of inverse trigonometric functions.



Function	Domain	Range
$y = \sin^{-1} x$	[–1, 1]	$\left[\frac{-\pi}{2},\frac{\pi}{2}\right]$
$y = \cos^{-1} x$	[–1, 1]	[ <b>0</b> , π]
$y = \tan^{-1} x$	R	$\left(\frac{-\pi}{2},\frac{\pi}{2}\right)$
$y = \cot^{-1} x$	R	(0, π)
$y = \sec^{-1} x$	R – (–1, 1)	$[0, \pi] - \left\{\frac{\pi}{2}\right\}$
$y = \operatorname{cosec}^{-1} x$	R – (–1, 1)	$\left[\frac{-\pi}{2},\frac{\pi}{2}\right]-\{0\}$

•	when $x \in [-1, 1]$	• $\sin^{-1}(\sin x) = x$ , when $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$
	$\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$ .	• $\cos^{-1}(\cos x) = x$ , when $x \in [0, \pi]$
•	when $x \in R$	• $\tan^{-1}(\tan x) = x$ , when $x \in \left(-\frac{\pi}{2}, \frac{\pi}{2}\right)$
	$\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$ .	• $\cot^{-1}(\cot x) = x$ , when $x \in (0, \pi)$
•	when $x \in \mathbb{R} - (-1, 1)$	• cosec <sup>-1</sup> (cosec x) = x, when $x \in \left[-\frac{\pi}{2}, \frac{\pi}{2}\right] - \{0\}$
	$\sec^{-1} x + \csc^{-1} x = \frac{\pi}{2}$ .	• sec <sup>-1</sup> (sec x) = x, when $x \in [0, \pi] - \left\{\frac{\pi}{2}\right\}$

- $\sin(\sin^{-1} x) = x$ , when  $x \in [-1, 1]$
- $\cos(\cos^{-1} x) = x$ , when  $x \in [-1, 1]$
- tan  $(\tan^{-1} x) = x$ , when  $x \in \mathbb{R}$
- $\cot(\cot^{-1} x) = x$ , when  $x \in \mathbb{R}$
- cosec (cosec<sup>-1</sup> x) = x, when  $x \in R (-1, 1)$
- sec (sec<sup>-1</sup> x) = x, when  $x \in R (-1, 1)$

•  $\sin^{-1}(-x) = -\sin^{-1}x$ , when  $x \in [-1, 1]$ •  $\cos^{-1}(-x) = \pi - \cos^{-1}x$ , when  $x \in [-1, 1]$ •  $\tan^{-1}(-x) = -\tan^{-1}x$ , when  $x \in \mathbb{R}$ •  $\cot^{-1}(-x) = \pi - \cot^{-1}x$ , when  $x \in \mathbb{R}$ •  $\csc^{-1}(-x) = -\csc^{-1}x$ , when  $x \in \mathbb{R}(-1, 1)$ •  $\sec^{-1}(-x) = \pi - \sec^{-1}x$ , when  $x \in \mathbb{R}(-1, 1)$ 

Illustration:  
Find the principal value of 
$$\sin^{-1}\left(\frac{1}{2}\right) + \cos^{-1}\left(\frac{-1}{2}\right)$$
.  
Solution: As,  $\sin^{-1}\left(\frac{1}{2}\right) = \sin^{-1}\left(\sin\frac{\pi}{6}\right) = \frac{\pi}{6}, \frac{\pi}{6} \in \left[\frac{-\pi}{2}, \frac{\pi}{2}\right]$   
 $\cos^{-1}\left(\frac{-1}{2}\right) = \pi - \cos^{-1}\left(\frac{1}{2}\right) = \pi - \frac{\pi}{3}, \frac{\pi}{3} \in [0, \pi]$   
so,  $\sin^{-1}\left(\frac{1}{2}\right) + \cos^{-1}\left(\frac{-1}{2}\right) = \frac{\pi}{6} + \pi - \frac{\pi}{3} = \frac{\pi}{6} + \frac{2\pi}{3} = \frac{5\pi}{6}$ 

Illustration:  
Find the principal value of sec<sup>-1</sup> (2) + sin<sup>-1</sup> 
$$\left(\frac{1}{2}\right)$$
 + tan<sup>-1</sup> ( $-\sqrt{3}$ ).  
Solution: As, sec<sup>-1</sup>(2) = cos<sup>-1</sup>  $\left(\frac{1}{2}\right)$   
tan<sup>-1</sup> ( $-\sqrt{3}$ ) =  $-\tan^{-1}(\sqrt{3})$  =  $-\tan^{-1}(\tan\frac{\pi}{3})$  =  $-\frac{\pi}{3}$ ,  $-\frac{\pi}{3} \in \left(\frac{-\pi}{2}, \frac{\pi}{2}\right)$   
cos<sup>-1</sup>  $\left(\frac{1}{2}\right)$  + sin<sup>-1</sup>  $\left(\frac{1}{2}\right)$  + tan<sup>-1</sup> ( $-\sqrt{3}$ ) =  $\frac{\pi}{2} - \frac{\pi}{3} = \frac{\pi}{6}$ 

**Illustration:** Find the range of the function  $f(x) = \tan^{-1} x + \cot^{-1} x$ . **Solution:** As,  $\tan^{-1} x + \cot^{-1} x = \frac{\pi}{2}$ so,  $f(x) = \frac{\pi}{2}$  (A constant function) Thus range of f(x) is  $\left\{\frac{\pi}{2}\right\}$ .

#### Illustration:

If  $\sin^{-1} x + \sin^{-1} y = \frac{2\pi}{3}$ , then find the value of  $\cos^{-1} x + \cos^{-1} y$ . **Solution:** As,  $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2} \Rightarrow \boxed{\cos^{-1} x = \frac{\pi}{2} - \sin^{-1} x}$  $\cos^{-1} x + \cos^{-1} y = \pi - (\sin^{-1} x + \sin^{-1} y) = \pi - \frac{2\pi}{3} = \boxed{\frac{\pi}{3}}$ 

Illustration: If  $a \le 2 \sin^{-1} x + \cos^{-1} x \le b$ , then find the value a and b. Solution: We know that,  $\sin^{-1} x + \cos^{-1} x = \frac{\pi}{2}$  and  $\frac{-\pi}{2} \le \sin^{-1} x \frac{\pi}{2}$ ,  $\Rightarrow 0 \le (\sin^{-1} x) + \frac{\pi}{2} \le \pi$   $\Rightarrow 0 \le (\sin^{-1} x) + \sin^{-1} x + \cos^{-1} x \le \pi$   $\Rightarrow 0 \le 2 \sin^{-1} x + \cos^{-1} x \le \pi$ , but given,  $a \le \sin^{-1}, x + \cos^{-1} x \le b$ Thus, a = 0 and  $b = \pi$ 

Illustration:  
If 
$$sin[cot^{-1} (1 + x)] = cos[tan^{-1} x]$$
, then find x.  
Solution: As,  $sin[cot^{-1} (1 + x)] = cos[tan^{-1} x]$   
 $\Rightarrow sin[sin^{-1} \frac{1}{\sqrt{x^2 + 2x + 2}}] = cos[cos^{-1} \frac{1}{\sqrt{1 + x^2}}]$   
 $\Rightarrow x^2 + 2x + 2 = 1 + x^2$   
 $\Rightarrow 2x = -1 \Rightarrow x = -0.5$ 

# Illustration:

If 
$$\tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \frac{\pi}{2}$$
, then prove that  $xy + yz + zx = 1$ .  
**Solution:** Let,  $\tan^{-1} x = A$ ,  $\tan^{-1} y = B$ ,  $\tan^{-1} z = C$   
so,  $A + B + C = \frac{\pi}{2} \Rightarrow A + B = \frac{\pi}{2} - C$   
 $\tan(A + B) = \tan\left(\frac{\pi}{2} - C\right) = \cot C$   
 $\frac{\tan A + \tan B}{1 - \tan A \cdot \tan B} = \frac{1}{\tan C} \Rightarrow \frac{x + y}{1 - xy} = \frac{1}{z}$   
 $\Rightarrow xz + yz = 1 - xy$   
 $\Rightarrow xz + yz + zx = 1$ 

# **ONE MARK QUESTIONS**

1. Principal Value of 
$$\cos^{-1}\left(\frac{\sqrt{3}}{2}\right) + \cos^{-1}\left(\frac{-1}{2}\right)is$$
  
(a)  $\pi$  (b)  $\frac{\pi}{3}$  (c)  $\frac{\pi}{6}$  (d)  $\frac{5\pi}{6}$   
2. Principal Value of  $\sin^{-1}\left(\sin\frac{3\pi}{5}\right)$  is

(a) 
$$\frac{3\pi}{5}$$
 (b)  $\frac{2\pi}{5}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{-3\pi}{5}$ 

3. Principal value of 
$$\cos^{-1}\left(\cos\frac{14\pi}{3}\right)$$
 is

(a) 
$$\frac{4\pi}{3}$$
 (b)  $\frac{2\pi}{3}$  (c)  $\pi$  (d)  $\frac{14\pi}{3}$ 

4. If the Principal value of  $\tan \tan^{-1}(\tan \frac{7\pi}{6})$  is  $\frac{a\pi}{b}$ , Where *a* & *b* are co-prime numbers, then (a + b)= (a) 13 -13 (c) 7 (d) (b) 5 5. If the Principal value of  $\cos^{-1}(\cos\frac{2\pi}{3}) + \sin^{-1}(\sin\frac{2\pi}{3})$  is  $\frac{a\pi}{b}$ , then |a-b|=2 (a) 0 (b) 1 (c) (d) 6. If  $\cos(\cos^{-1}\frac{1}{3} + \sin^{-1}x) = 0$ , then (3x + 1) =(b) 1 (c) 2 (d) (a) 0 4 7. If  $\sin(\sin^{-1}\frac{3}{5} + \cos^{-1}x) = 1$ , then (5x - 2)= (b) 1 (a) 0 (c) 2 (d) 4 Domain of the function  $\cos^{-1}(2x-1)$  is 8. (b) [-1,1] (c) [0,1] (a) R (d) [0,2] Domain of the function  $f(x) = \sin^{-1} \sqrt{x-1}$  is 9. (b) [–1,1] (c) [0,1] (d) (a) [1,2] [0,2] 10. Principal value of sec<sup>-1</sup>(2) + sin<sup>-1</sup>( $\frac{1}{2}$ ) + tan<sup>-1</sup>( $-\sqrt{3}$ ) is (a)  $\frac{\pi}{3}$  (b)  $\frac{2\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{\pi}{6}$ 

[0,1]

(d)

[0,2]

11. Domain of the function 
$$f(x) = \cos^{-1}\sqrt{x+1}$$
 is  
(a) [1,2] (b) [-1,0] (c) [0,1] (d)

[Class XII : Maths]

(b)

(a) [1,2]

12. Domain of the function  $f(x) = \sin^{-1}(-x^2)$  is [0,1] (a) [1,2] (b) [-1,0] (c) (d) [-1,1] 13. Domain of the function  $f(x) = \sin^{-1}(2x+3)$  is (a) [-2,2] (b) [-2,-1] (c) [0,1] (d) [-1,1] 14. If Domain of the function  $f(x) = \sin^{-1}(x^2 - 4)$  is  $[-b, -a] \cup [a, b]$  then the value of  $(a^2 + b^2)$ is. (a) 8 (b) 3 (c) 5 4 (d) 15. If  $\sin^{-1} x_1 + \sin^{-1} x_2 = \pi$ , then the value of  $(x_1 + x_2)$  is (a) 0 (b) 1 (c) -1 (d) 2 16. If  $\cos^{-1} a + \cos^{-1} b = 2\pi$ , then the value of  $(a - b)^2$  is (a) 0 (b) 1 (c) -1 (d) 4 17.  $\cos^{-1}[\sin(\cos^{-1}\frac{1}{2})]=$ (a)  $\frac{\pi}{6}$  (b)  $\frac{2\pi}{3}$  (c)  $\frac{\pi}{2}$  (d)  $\frac{\pi}{3}$ 18. Principal value of  $\sin^{-1}(\cos\frac{34\pi}{5})$  is (a)  $\frac{\pi}{5}$  (b)  $\frac{-\pi}{10}$  (c)  $\frac{3\pi}{10}$  (d)  $\frac{-3\pi}{10}$ 19. If cot  $(\cos^{-1}\frac{7}{25}) = x$ , then  $\sqrt{24x+2} =$ (b) 2 (c) 3 (a) 1 (d) 4 20. If  $\tan^{-1} x + \tan^{-1} y = \frac{4\pi}{5} \& \cot^{-1} x + \cot^{-1} y = \frac{k\pi}{5}$ , then k = (a) 1 (b) 2 (c) 3 (d) 4 21.  $\sum_{i=1}^{2023} \cos^{-1} x_i = 0$ , then the velue of  $\sum_{i=1}^{2023} x_i$  is (a) 0 (b) 1 (c) 2023 (d) -2023



23. If graph of f(x) is shown below, identify the function f(x) & find the value of  $f(-\frac{1}{2})$ .



#### ASSERTION-REASON BASED QUESTIONS (Q.24 & Q.25)

In the following questions, a statement of assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A.
- (b) Both A and R are true but R is not the correct explanation of A.
- (c) A is true but R is false.
- (d) A is false but R is true.
- 24. ASSERTION (A): The range of the function  $f(x) = \sin^{-1}x + \frac{3\pi}{2}$ , where

$$x \in [-1,1], \text{ is } [\frac{\pi}{2}, \frac{5\pi}{2}].$$

REASON (R): The range of the principal value branch of sin<sup>-1</sup>x is  $[0, \pi]$ .

25. ASSERTION (A): All trigonometric function have their inverses over their respective domains.

REASON (R): The inverse of  $\tan^{-1}x$  exists for some  $x \in R$ .

#### **TWO MARKS QUESTIONS**

#### 26. Match the following:

If  $\cos^{-1}a + \cos^{-1}b = 2\pi$  and  $\sin^{-1}c + \sin^{-1}d = \pi$  then

	Column 1		Column 2
А	abcd	Р	0
В	$a^2 + b^2 + c^2 + d^2$	Q	1
С	(d - a) + (c - d)	R	2
D	$a^3 + b^3 + c^3 + d^3$	S	4

27. Find the value of  $\cos\left[\cos^{-1}\left(\cos\frac{5\pi}{3}\right) + \sin^{-1}\left(\sin\frac{5\pi}{3}\right)\right]$ 

28. If  $P = \tan^2(\sec^{-1} 2) + \cot^2(\csc^{-1} 3)$ , then find the value of  $(P^2 + P + 11)$ .

29. If  $P = \sec^2(\tan^{-1} 2) + \csc^2(\cot^{-1} 3)$ , then find the value of  $(P^2 - 2P)$ .

30. Find the value of 
$$\sin\left(\frac{1}{2}\cot^{-1}\left(\frac{3}{4}\right)\right)$$
.  $\left[\text{Hint}:\sin\frac{\theta}{2}=\sqrt{\frac{1-\cos\theta}{2}}\right]$ 

31. Solve for x: 
$$\tan^{-1}\sqrt{x(x+1)} + \sin^{-1}\sqrt{x^2 + x + 1} = \frac{\pi}{2}$$

32. Find the value of x, such that  $\sin^{-1}x = \frac{\pi}{6} + \cos^{-1}x$ .

- 33. Find x, if  $\sin^{-1}x \cos^{-1}x = \frac{\pi}{2}$
- 34. If  $\tan^{-1}(\cot x) = 2x$ , find *x*.

35. Solve for 
$$x$$
:  $\cos^{-1}\left(\cos\frac{3\pi}{4}\right) + \sin^{-1}\left(\sin\frac{3\pi}{4}\right) = x$ 

#### **THREE MARKS QUESTIONS**

36. Find the value of *k*, if 100 sin(2 tan<sup>-1</sup> (0.75)) = *k* [Hint: sin2 $\theta$  = 2sin $\theta$  cos $\theta$ ]

37. Prove that:

(a) 
$$\cot^{-1}\left(\frac{\sqrt{1+\sin x} + \sqrt{1-\sin x}}{\sqrt{1+\sin x} - \sqrt{1-\sin x}}\right) = \frac{x}{2}, x \in \left(0, \frac{\pi}{4}\right)$$

(b) 
$$\tan^{-1}\left(\frac{\sqrt{1+x}-\sqrt{1-x}}{\sqrt{1+x}-\sqrt{1-x}}\right) = \frac{x}{4} - \frac{1}{2}\cos^{-1}x$$

(c) 
$$\tan^{-1}\left(\frac{1}{2}\sin^{-1}\frac{3}{4}\right) = \frac{4-\sqrt{7}}{3}$$

(d) 
$$\sin^{-1}\left(2\tan^{-1}\left(\frac{2}{3}\right)\right) = \frac{12}{13}$$

38. (a) Prove that 
$$\cos[\tan^{-1}{\sin(\cot^{-1} x)}] = \sqrt{\frac{x^2 + 1}{x^2 + 2}}$$

(b) Prove that 
$$\tan\left(\frac{\pi}{4} + \frac{1}{2}\cos^{-1}\frac{a}{b}\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\cos^{-1}\frac{a}{b}\right) = \frac{2b}{a}$$

(c) Prove that 
$$\tan\left(\frac{\pi}{4} + \frac{1}{2}\tan^{-1}\frac{a}{b}\right) + \tan\left(\frac{\pi}{4} - \frac{1}{2}\tan^{-1}\frac{a}{b}\right) = \frac{2\sqrt{a^2 + b^2}}{b}.$$

(d) Prove that : 
$$\tan^{-1}\left(\frac{\sqrt{1+x}-\sqrt{1-x}}{\sqrt{1+x}+\sqrt{1-x}}\right) = \frac{\pi}{4} - \frac{1}{2}\cos^{-1}x$$

(e) Prove that : 
$$\tan^{-1}\left(\frac{\sqrt{1+\cos x} + \sqrt{1-\cos x}}{\sqrt{1+\cos x} - \sqrt{1-\cos x}}\right) = \frac{\pi}{4} + \frac{x}{2}, x \in (0, \frac{\pi}{2})$$

(f) Prove that : 
$$\cot^{-1}\left(\frac{\sqrt{1+\sin x}+\sqrt{1-\sin x}}{\sqrt{1+\sin x}-\sqrt{1-\sin x}}\right) = \frac{x}{2}, x \in \left(0, \frac{\pi}{4}\right)$$

39. Solve for x:

(a) 
$$\sin^{-1}(6x) + \sin^{-1}(6\sqrt{3}x) = \frac{\pi}{2}$$

(b) Solve for x: 
$$\sin^{-1}(6x) + \sin^{-1}(6\sqrt{3}x) = \frac{-\pi}{2}$$

(c) 
$$(\tan^{-1} x)^2 + (\cot^{-1} x)^2 = \frac{5\pi^2}{8}$$
.

40. Solve for 
$$x : \cos(\tan^{-1} x) = \sin\left(\cot^{-1} \frac{3}{4}\right), x > 0$$

# **FIVE MARKS QUESTIONS**

Illustration: (For Solving Q.41)  
If 
$$\cos^{-1} x + \cos^{-1} y + \cos^{-1} z = \pi$$
, then prove that  $x^2 + y^2 + z^2 + 2xyz = 1$ .  
Solution: Let,  $\cos^{-1} x = A$ ,  $\cos^{-1} y = B$ ,  $\cos^{-1} z = C$   
so,  $A + B + C = \pi \implies A + B = \pi - C$   
Thus,  $\cos(A + B) = \cos(\pi - C)$   
 $\implies \cos A \cos B - \sin A \sin B = -\cos C$   
 $\implies \cos A \cos B - \sqrt{1 - \cos^2 A} \sqrt{1 - \cos^2 B} = -\cos C$   
 $\implies \cos A \cos B - \sqrt{1 - x^2} \sqrt{1 - y^2} = -z$   
 $\implies (xy + z) = \sqrt{1 - x^2} \sqrt{1 - y^2}$   
On squaring both the sides, we get  
 $(xy + z)^2 = (1 - x^2) (1 - y^2)$   
 $\implies x^2 y^2 + z^2 + 2xyz = 1 - x^2 - y^2 + x^2 y^2$   
 $\therefore x^2 + y^2 + z^2 + 2xyz = 1$ 

41. Prove the following:

(a) If 
$$\cos^{-1}\left(\frac{x}{a}\right) + \cos^{-1}\left(\frac{y}{b}\right) = \alpha$$
, then prove that  $\frac{x^2}{a^2} + \frac{y^2}{b^2} - \frac{2xy}{ab}\cos\alpha = \sin^2\alpha$ 

(b) If  $\cos^{-1}\left(\frac{x}{2}\right) + \cos^{-1}\left(\frac{y}{3}\right) = \theta$ , then prove that  $9x^2 + 4y^2 - 12xy\cos\theta = 36\sin^2\theta$ .

42. Prove the following:

(a) If  $\tan^{-1} x + \tan^{-1} y + \tan^{-1} z = \pi$ , then prove that x + y + z = xyz

(a) If  $\cot^{-1} x + \cot^{-1} y + \cot^{-1} z = \pi$ , then prove that xy + yz + zx = 1

#### **CASE STUDIES**

43. On National Mathematics Day, December 22, 2020, Mathematics Teachers of DOE organized Mathematical Rangoli Competition for the students of all DOE schools to celebrate and remembering the contribution of Srinivasa Ramanujan to the field of mathematics. The legendary Indian mathematician who was born on this date in 1887.



Team A of class XI students made a beautiful Rangoli on Trigonometric Identities as shown in the figure Above, While Team B of class XII students make the Rangoli on the graph of Trigonometric and Inverse Trigonometric Functions. As shown in the following figure.



On the basis of above information, Teacher asked few questions from Team B. Now you try to answer. Those questions which are as follows:

- (a) Write the domain & range (principal value branch) of the function  $f(x) = \tan^{-1} x$ ?
- (b) If the principal branch of  $\sec^{-1}x$  is  $[0,\pi] \{k\pi\}$ , then find the value of k.
- (c) Draw the graph of  $\sin^{-1} x$ , where  $x \in [-1, 1]$ . Also write its Principal branch Range.

#### SELF ASSESSMENT-1

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE COR-RECT ALTERNATIVE.

1. If 
$$\cos\left(\cos^{-1}\frac{2}{3} + \sin^{-1}x\right) = 0$$
, then  $(3x - 1)$ 

- (a) 0 (b) 1
- (d) 2 (c) -1

2. Domain of the function  $\cos^{-1}\left(\frac{x}{2}-1\right)$  is

- (a) [0, 2] (b) [-1, 1]
- (d) [0, 4] (c) [0, 1]

3. If  $\cos^{-1}a + \cos^{-1}b = 2\pi$  and  $\sin^{-1}c + \sin^{-1}d = \pi$ , then  $a^2 + b^2 + c^2 + d^2 = 1$ 

- (a) 0 (b) 1
- (c) 2 (d) 4
- 4. The principal value of  $\cos^{-1}\left(\cos\frac{2\pi}{3}\right) + \sin^{-1}\left(\sin\frac{2\pi}{3}\right)$  is
  - (a) 0 (b) π

(c) 
$$2\pi$$
 (d)  $\frac{4\pi}{3}$ 

5. If  $\cos^{-1}\left(\frac{1}{x}\right) = \theta$ , then  $\tan\theta =$ (a) *x* (b)  $x^2 + 1$ (c)  $\sqrt{x^2 + 1}$ (d)  $\sqrt{x^2 - 1}$ 

### **SELF ASSESSMENT-2**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE COR-RECT ALTERNATIVE.

1. If 
$$\sin^{-1}x + \sin^{-1}y + \sin^{-1}z = \frac{3\pi}{2}$$
, then  $(x^3 + y^3 + z^3 - 3xyz) =$   
(a) 0 (b) 1  
(c) -1 (d) 2

- 2. Principal Range of the function  $\sin^{-1}x$  is
  - (a) [0, π] (b) (0, π)

(c) 
$$\left[\frac{-\pi}{2},\frac{\pi}{2}\right]$$
 (d)  $\left(\frac{-\pi}{2},\frac{\pi}{2}\right)$ 

3. If 
$$\cos^{-1}\left(\cos\frac{5\pi}{3}\right) + \sin^{-1}\left(\sin\frac{5\pi}{3}\right) = x$$
, then  $x =$   
(a) 0 (b)  $\pi$ 

(c) $\frac{5\pi}{3}$	(d) $\frac{10}{3}$	$\frac{2\pi}{3}$		
4. If $\sin^{-1}\left(\frac{x}{5}\right) + \csc^{-1}\left(\frac{5}{4}\right) = \frac{1}{5}$	$\frac{\pi}{2}$ , then x	<		
(a) 0	(b) 1			
(c) 2	(d) 3			
5. Range of $f(x) = \sin^{-1}x + \tan^{-1}x$	x + sec <sup>-1</sup> )	xis		
(a) $\left[\frac{\pi}{4},\frac{3\pi}{4}\right]$	(B) $\left(\frac{1}{2}\right)$	$\left(\frac{\pi}{4},\frac{3\pi}{4}\right)$		
$(C)  \left\{\frac{\pi}{4}, \frac{3\pi}{4}\right\}$	(D) $\left[\frac{\pi}{2}\right]$	$\left(\frac{\pi}{4}, \frac{\pi}{2}\right)$		
		ANSWER		
	One M	lark Questions		
1. ( <i>d</i> ) $\frac{5\pi}{6}$	2. (b	) $\frac{2\pi}{5}$	3.	(b) $\frac{2\pi}{3}$
4. ( <i>c</i> ) 7	5. (a)	) 0	6.	(c) 2
7. (b) 1	8. (c)	) [0, 1]	9.	(a) [1,2]
π				
10. ( <i>d</i> ) $\frac{1}{6}$	11. (b	) [—1,0]	12.	(d) [-1,1]
13. ( <i>b</i> ) [–2,–1]	14. (a	a) 8	15.	( <i>d</i> ) 2
16. ( <i>a</i> ) 0	17. ( <i>a</i> )	$\frac{\pi}{6}$	18.	(d) $\frac{-3\pi}{10}$
19. (c) 3	20. (a)	) 1	21.	(c) 2023
		, О <del>л</del>		( )
22. (b) 2024	23. (d)	$\frac{2\pi}{3}$	24.	(c) A is true but R is
false.	25. (d)	) A is false but R	is true.	
	Two Ma	arks Questions		
26. $A \rightarrow Q, B \rightarrow S, C \rightarrow R, D \rightarrow Q$	P		27. 1	
28. ( <i>P</i> <sup>2</sup> + <i>P</i> + 11) = 143	29. (P	<sup>2</sup> – 2P) = 195	$30. \ \frac{1}{\sqrt{5}}$	
31. <i>x</i> = 0 or −1	32. $\frac{\sqrt{2}}{2}$	<u>3</u> 2	33. 1	

34.  $\frac{\pi}{6}$  35.  $\pi$ 

**Three Marks Questions** 

36. 96  
39. (a) 
$$x = \frac{1}{12}$$
 (b)  $x = \frac{-1}{12}$  (c)  $x = -1$   
40.  $x = \frac{3}{4}$ 

# CASE STUDIES BASED QUESTION

43. (a) Domain =  $R = (-\infty, \infty)$ , Range =  $(\frac{-\pi}{2}, \frac{\pi}{2})$  (b) k = 0.5

		SELF ASSESSMI	ENT-1	
1. (b)	2. (d)	3. (d)	4. (b)	5. (d)
		SELF ASSESSMI	ENT-2	
1. (a)	2. (c)	3. (a)	4. (d)	5. (c)
## **CHAPTER-3**

# MATRICES

Matrices find many applications is scientific field and apply to practical real life problem. Matrices can be solved physical related application and one applied in the study of electrical circuits, quantum mechanics and optics, with the help of matrices, calculation of battery power outputs, resistor conversion of electrical energy into another useful energy, these matrices play a role in calculation, with the help of matrices problem related to Kirchhoff law of voltage and current can be easily solved.



Matrices can play a vital role in the projection of three dimensional images into two dimensional screens, creating the realistic decreeing motion. Now day's matrices are used in the ranking of web pages in the Google search. It can also be used in generalization of analytical motion like experimental and derivatives to their high dimensional.

Matrices are also used in geology for seismic survey and it is also used for plotting graphs. Matrices are also used in robotics and automation in terms of base elements for the robot movements. The movements of the robots are programmed with the calculation of matrices 'row and column' controlling of matrices are done by calculation of matrices.

## TOPIC TO BE COVERED AS PER CBSE LATEST CURRICULUM (2024-25)

- Concept, notaion, order, equality, types of matrices, zero and identity matrix, transpose of a matrix, symmetric and skew symmetric matrices.
- Operation on matrices: Addition and multiplication and multiplication with a scalar. Simple
  properties of addition, multiplication and scalar multiplication. Oncommutativity of multiplication of matrices and existence of non-zero matrices whose product is the zero
  matrix (restrict to square matrices of order 2).
- Invertible matrices and proof of the uniqueness of inverse, if it exists; (Here all matrices will have real entries).

Matrices are defined as a rectangular arrangement of numbers of functions. Since it is a rectangular arrangement, it is 2-dimensional.

A two-dimensional matrix consists of the number of rows (m) and a number of columns (n). Horizontal ones are called Rows and Vertical ones are called columns.



## ORDER OF MATRIX

The order of matrix is a relationship with the number of elements present in a matrix.

The order of a matrix is denoted by  $m \times n$ , where m and n are the number of Rows and Columns Respectively and the number of elements in a matrix will be equal to the product of m and n.

## **TYPES OF MATRICES**

#### **Row Matrix**

A matrix having only one row is called a row matrix.

Thus  $A = [A_{ij}]_{mxn}$  is a row matrix if m = 1. So, a row matrix can be represented as  $A = [A_{ij}]_{1xn}$ 

It is called so because it has only one row and the order of a row matrix will hence be  $1 \times n$ .

For example,

 $A = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$  is row matrix of order 1 × 4. Another example of the row matrix is

 $B = [0 \ 9 \ 4]$  which is of the order  $1 \times 3$ .

## **Column Matrix**

A matrix having only one column is called a column matrix. Thus,  $A = [A_{ij}]_{mxn}$  is a column matrix if n = 1.

Hence, the order is  $m \times 1$ . An example of a column matrix is:

$$A = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix}, B = \begin{pmatrix} M \\ A \\ T \\ H \end{pmatrix}$$

In the above example, A and B are 3 × 1 and 4 × 1 order matrices respectively.

#### **Square Matrix**

If the number of rows and the number of columns in a matrix are equal, then it is called a square matrix.

Thus,  $A = [A_{ij}]_{mxn}$  is a square matrix if m = n; For example is a square matrix of order  $3 \times 3$ .

$$\mathbf{A} = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$

For Additional Knowledge:

F

The sum of the diagonal elements in a square matrix A is called the trace of matrix A, and which is sdenoted by tr(A);

 $tr(A) = a_{11} + a_{22} + \dots + a_{nn}$ 

#### Zero or Null Matrix

If in a matrix all the elements are zero then it is called a zero matrix and it is generally denoted by O. Thus,  $A = [A_{ij}]_{mxn}$  is a zero-matrix if  $a_{ij} = 0$  for all *i* and *j*; For example

$$\boldsymbol{A} = \begin{pmatrix} \boldsymbol{0} \\ \boldsymbol{0} \\ \boldsymbol{0} \end{pmatrix}, \, \boldsymbol{B} = \begin{pmatrix} \boldsymbol{0} & \boldsymbol{0} \\ \boldsymbol{0} & \boldsymbol{0} \end{pmatrix}$$

Here A and B are Null matrix of order 3 × 1 and 2 × 2 respectively.

#### **Diagonal Matrix**

If all the non-diagonal elements of a square matrix, are zero, then it is called a diagonal matrix.

Thus, a square matrix  $A = [a_{ij}]$  is a diagonal matrix if all i = 0, when  $i \neq j$ ;

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 3 & 0 \\ 0 & 0 & 4 \end{pmatrix}, B = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 4 \end{pmatrix}, C = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 4 \end{pmatrix}, D = \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$$

*A*, *B* and *C* are diagonal matrix of order  $3 \times 3$ , and *D* is a diagonal matrix of order  $2 \times 2$ . Diagonal matrix can also be denoted by *A* = diagonal [2 3 4], *B* = diag [2 0 4], *C* = [0 0 4]

Important things to note:

- (i) A diagonal matrix is always a square matrix.
- (ii) The diagonal elements are characterized by this general form:  $a_{ij}$ , where i = j. This means that a matrix can have only one diagonal.

#### Scalar Matrix

If all the elements in the diagonal of a diagonal matrix are equal, it is called a scalar matrix. Thus, a square matrix  $A = [a_{ij}]$  is a scalar matrix if

$$A = [a_{ij}] = \begin{cases} 0; & i \neq j \\ k; & i = j \end{cases}$$
 Where, *k* is constant.

For example A and B are scalar matrix of order 3 × 3 and 2 × 2 respectively.

$$A = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}, B = \begin{pmatrix} -7 & 0 \\ 0 & -7 \end{pmatrix}$$

#### **Unit Matrix or Identity Matrix**

If all the elements of a principal diagonal in a diagonal matrix are 1, then it is called a unit matrix.

A unit matrix of order *n* is denoted by  $I_n$ . Thus, a square matrix  $A = [a_{ij}]_{m \times m}$  is an identity matrix if

$$A = [a_{ij}] = \begin{cases} 0; & i \neq j \\ 1; & i = j \end{cases}$$

For example  $I_3$  and  $I_2$  are identity matrix of order 3 × 3 and 2 × 2 respectively.

$$I_3 = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, I_2 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$$

- All identity matrices are scalar matrices
- All scalar matrices are diagonal matrices
- · All diagonal matrices are square matrices

#### **Triangular Matrix**

A square matrix is said to be a triangular matrix if the elements above or below the principal diagonal are zero. There are two types of Triangular Matrix:

#### **Upper Triangular Matrix**

A square matrix  $[a_{ij}]$  is called an upper triangular matrix, if  $a_{ij} = 0$ , when i > j.

$$A = \begin{pmatrix} D & O & E \\ 0 & D & O \\ 0 & 0 & E \end{pmatrix}$$
 is an upper triangular matrix of order 3 × 3.

#### Lower Triangular Matrix

A square matrix is called a lower triangular matrix, if  $a_{ij} = 0$ , when i > j.

$$A = \begin{pmatrix} D & 0 & 0 \\ O & D & 0 \\ E & O & E \end{pmatrix}$$

is a lower triangular matrix of order 3 × 3.

## Transpose of a Matrix

Let A be any matrix, then on interchanging rows and columns of A. The new matrix so obtained is transpose of A donated  $A^{\tau}$  or A'.

[order of  $A = m \times n$ , then order of  $A^{T} = n \times m$ ]

Properties of transpose matrices A and B are:

(a) 
$$(A^{T})^{T} = A$$
 (b)  $(kA)^{T} = kA^{T} (k = \text{constant})$ 

(c)  $(A + B)^{T} = A^{T} + B^{T}$  (d)  $(AB)^{T} = B^{T} \cdot A^{T}$ 

Symmetric Matrix and Skew-Symmetric matrix

- A square matrix  $A = [a_{ij}]$  is symmetric if  $A^T = A$  i.e.  $a_{ij} \forall i$  and j
- A square matrix  $A = [a_{ij}]$  is skew-symmetric if  $A^T = -A$  i.e.  $a_{ij} = -a_{ij} \forall i$  and j

(All diagonal elements are zero in skew-symmetric matrix)

#### Illustration:

A is matrix of order 2022 × 2023 and B is a matrix such that  $AB^T$  and  $B^TA$  are both defined, then find the order of matrix B.

Solution: Let the order of matrix be R × C, So,

 $(A)_{2022 \times 2023} (B^{T})_{C \times R} \quad \Rightarrow \quad C = 2023 \text{ (As } AB^{T} \text{ is defined)}$ 

$$(B^{T})_{C\times R}(A)_{2022\times 2023} \Rightarrow R = 2022 \text{ (As } B^{T}A \text{ is defined)}$$

Thus order of matrix *B* is (2022 × 2023).

#### Illustration:

If A is a skew symmetric matrix, then show that  $A^2$  is symmetric.

**Solution:** As A is skew-symmetric, 
$$A^{T} = -A^{T}$$

 $(A^2)^{\mathsf{T}} = (A.A)^{\mathsf{T}} = A^{\mathsf{T}}.A^{\mathsf{T}} = (-A) (-A) = A^2$ 

As  $(A^2)^T = A^2$ 

 $\Rightarrow$  Thus,  $A^2$  is symmetric.

Illustration: If  $\begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix} + X = \begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix}$ , where  $X = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  then find the value of a + c - b - d. Solution: As,  $\begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix} + \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix}$ ,  $\Rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 3 & 4 \\ 5 & 6 \end{pmatrix} - \begin{pmatrix} 1 & -1 \\ 2 & 3 \end{pmatrix} = \begin{pmatrix} 3 - 1 & 4 + 1 \\ 5 - 2 & 6 - 3 \end{pmatrix}$   $\Rightarrow \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} 2 & 5 \\ 3 & 3 \end{pmatrix}$ On compairing the corresponding elements, we get, a = 2, b = 5, c = 3, d = 3Thus, a + c - b - d = 5 - 5 - 3 = -3

## Illustration:

If A is a diagonal matrix of order  $3 \times 3$  such that  $A^2 = A$ , then find number of possible matrices A.

Solution: As, A is a diagonal matrix of order 3 × 3

Let,  $A = \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix}$   $\Rightarrow A^{2} = \begin{pmatrix} a^{2} & 0 & 0 \\ 0 & b^{2} & 0 \\ 0 & 0 & c^{2} \end{pmatrix}$ As  $A^{2} = A \Rightarrow \begin{pmatrix} a & 0 & 0 \\ 0 & b & 0 \\ 0 & 0 & c \end{pmatrix} = \begin{pmatrix} a^{2} & 0 & 0 \\ 0 & b^{2} & 0 \\ 0 & 0 & c^{2} \end{pmatrix}$ So, a = 0 or -1, similarly b and c can take 2 values (0 and -1). Thus, total number of possible matrices are  $2 \times 2 \times 2 = 8$ .

## **ONE MARK QUESTIONS**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. If 
$$A = \begin{bmatrix} a_{ij} \end{bmatrix}_{2\times 2} = \begin{cases} 0, when i = j \\ 1, when i \neq j \end{cases}$$
, then  $A^2 =$   
(a)  $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  (b)  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  (c)  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  (d)  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$   
2. If  $A = \begin{bmatrix} a_{ij} \end{bmatrix}_{2\times 2} = \begin{cases} 0, when i = j \\ 1, when i \neq j \end{cases}$ , then  $A^{3025} =$   
(a)  $\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$  (b)  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  (c)  $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$  (d)  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$   
3. If  $A = \begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix}, B = \begin{pmatrix} x & 0 \\ 1 & 1 \end{pmatrix}$  and  $A = B^2$ , then x equals  
(a)  $\pm 1$  (b) 1 (c)  $-1$  (d) 2  
4. If  $A = \begin{pmatrix} 1 & x^2 - 2 & 3 \\ 7 & 5 & 7 \\ 3 & 7 & -5 \end{pmatrix}$  be a symmetric matrix, then x equals  
(a)  $\pm 3$  (b)  $\pm 2$  (c)  $\pm \sqrt{2}$  (d) 0  
5. If  $A = \begin{pmatrix} 0 & x^2 + 6 & 1 \\ -5x & x^2 - 9 & 7 \\ -1 & -7 & 0 \end{pmatrix}$  be a skew -symmetric matrix, then x equals  
(a)  $\pm 3$  (b) 3 (c)  $-3$  (d) 0  
6. If  $A = \begin{pmatrix} 2y - 7 & 0 & 0 \\ 0 & x - 3 & 0 \\ 0 & 0 & 7 \end{pmatrix}$  be a scalar matrix, then (x+y) equals  
(a) 7 (b) 14 (c) 16 (d) 17

7.	If A is matrix of orde defined, then the orde	r 2023 er of m	3 × 2024 and B is natrix B is	sam	atrix such that A	AB' and	B'A both are
	(a) 2023 × 2024	(b)	2023 × 2023	(c)	2024 × 2024	(d) 2	2024 × 2023
8.	If A is matrix of order defined, then the order	er 2023 er of m	3 × 2024 and B i natrix B is	is a r	natrix such that	AB and	d BA both are
	(a) 2023 × 2024	(b)	2023 × 2023	(c)	2024 × 2024	(d)	2024 × 2023
	( 2	0 y	(-x)				
9.	If $A =  x + y - 2 $	3	0 be a diago	nal m	natrix then (xy) e	quals	
	( O	0	4 )				
	(a 1	(b)	2	(c)	3	(d)	4
10.	If all entries of a squa matrices are possible	ire ma ?	trix of order 2 are	eithe	er 3, –3 <i>or</i> 0, then	how m	any Non-zero
	(a) 80	(b)	81	(c)	27	(d)	64
11.	If all entries of a squ matrices are possible	are m ?	atrix of order 3 a	ire ei	ther 1 <i>or</i> 0, then	how n	any Diagonal
	(a) 512	(b)	8	(b)	6	(d)	2
12.	If all entries of a squar are possible?	re mat	rix of order 3 are e	either	3 <i>or</i> 0, then how I	many S	calar matrices
	(a) 1	(b)	8	(c)	6	(d)	2
13.	If all entries of a squ matrices are possible	uare m ?	natrix of order 3	are e	ither 5 <i>or</i> 0, the	n how	many Identity
	(a) 1	(b)	8	(c)	2	(d)	0
14.	If there are five one' symmetric matrices of	s i.e. 1 of orde	l, 1, 1, 1, 1 & four er 3 ×3 possible?	zero	es i.e. 0, 0, 0, 0, 0,	then to	otal number of
	(a) 10	(b)	12	(c)	3	(d)	9
15.	If $x \begin{pmatrix} 1 \\ 2 \end{pmatrix} + y \begin{pmatrix} 2 \\ 5 \end{pmatrix} = \begin{pmatrix} 2 \\ 5 \end{pmatrix}$	(4), tł	nen				
	(a) x= 1, y = 2	(b)	x= 2, y=1	(c)	x= 1, y=–1	(d)	x= 3, y=2
16.	The product $\begin{pmatrix} a & b \\ -b & a \end{pmatrix}$	$\binom{a}{b}$	$\begin{pmatrix} -b \\ a \end{pmatrix}$ , is equal to	0			
	(a) $\begin{pmatrix} a^2 + b^2 \\ 0 & a^2 \end{pmatrix}$	$\left( \begin{array}{c} 0 \\ + b^2 \end{array} \right)$	(b)		$\begin{pmatrix} a^2 + b^2 & 0 \\ a^2 + b^2 & 0 \end{pmatrix}$		

(c) 
$$\begin{pmatrix} a^2 + b^2 & 0 \\ 0 & 0 \end{pmatrix}$$
 (d)  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$   
17. If A is a square matrix such that  $A^2 = I$ , then  $(A - I)^3 + (A + I)^3 - 7A$  is equal to  
(a) 1 (b) A (c)  $3A - I$  (d)  $A - I$   
18. If A and B are two non-zero matrices such that AB = A, BA = B and  $(A + B)^3 = k$   $(A + B)$ , then k is equal to  
(a) 1 (b) 2 (c) 4 (d) 8  
19. If A is a square matrix such that  $A^2 = A$ , then $(A + I)^2 - 3A$  is equal to  
(a) 1 (b) A (c)  $2A$  (d)  $3I$   
20. If a matrix  $A = (123)$ , then the matrix A.A' (where A' is the transpose of A) is  
(a)  $(123)_{1\times 3}$  (b)  $(14)_{1\times 1}$  (c)  $(6)_{1\times 1}$  (d)  $\begin{pmatrix} 3 \\ 2 \\ 1 \\ 3_{3\times 1} \end{pmatrix}$   
21. If  $A = \begin{pmatrix} 3 & 4 \\ 5 & 2 \end{pmatrix}$  and  $2A + B$  is a null matrix, then B is equal to]  
(a)  $\begin{pmatrix} 3 & 4 \\ 5 & 2 \end{pmatrix}$  (b)  $\begin{pmatrix} -3 & -4 \\ -5 & -2 \end{pmatrix}$  (c)  $\begin{pmatrix} -6 & -8 \\ -10 & -2 \end{pmatrix}$  (d)  
(e)  $\begin{pmatrix} -6 & -8 \\ -10 & -4 \end{pmatrix}$   
22. If  $A = \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  and  $(3I + 4A) (3I - 4A) = x^2 I$ , then value of x is/are  
(a)  $\pm 3$  (b)  $\pm \sqrt{7}$  (c)  $\pm 5$  (d) 0  
23. If  $A = \begin{pmatrix} 2 & 8 \\ 6 & 4 \end{pmatrix} = p + Q$ , where P is a symmetric and Q is a skew-symmetric matrix, then Q is equal to  
(a)  $\begin{pmatrix} 2 & 6 \\ 8 & 4 \end{pmatrix}$  (b)  $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$  (c)  $\begin{pmatrix} 0 & 2 \\ -2 & 0 \end{pmatrix}$  (d)  
(c)  $\begin{pmatrix} 0 - 2 \\ 2 & 0 \end{pmatrix}$ 

ASSERTION-REASON BASED QUESTIONS (Q.24 & Q.25)

In the following questions, a statement of assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices

- (a) Both A and R are true and R is the correct explanation of A.
- (b) Both A and R are true but R is not the correct explanation of A.
- (c) A is true but R is false.
- (d) A is false but R is true.

24. ASSERTION: Matrix A =  $\begin{pmatrix} 0 & 1 & -2 \\ -1 & 1 & 3 \\ 2 & -3 & 0 \end{pmatrix}$  is a skew-symmetric matrix.

**REASONING:** A matrix A is skew-symmetric if  $A^{I} = -A$ .

25. ASSERTION : For matrices 
$$A = \begin{pmatrix} 1 & 4 \\ 2 & 3 \end{pmatrix}$$
 and  $B = \begin{pmatrix} 4 & 2 \\ 9 & 1 \end{pmatrix}$ ,

 $(A + B) (A - B) = A^2 - AB + BA - B^2$ 

REASONING : Matrix multiplication is not commutative.

#### **TWO MARKS QUESTIONS**

- 26. If A is a square matrix, then show that
  - (a)  $(A + A^{T})$  is symmetric matrix.
  - (b)  $(A A^{T})$  is symmetric matrix.
  - (c)  $(AA^{T})$  is symmetric matrix.
- 27. Show that every square matrix can be expressed as the sum of a symmetric and a skewsymmetric matrix.
- 28. If A and B are two symmetric matrices of same order, then show that
  - (i) (AB BA) is skew-symmetric Matrix.
  - (ii) (AB + BA) is symmetric Matrix.

29. (a) If 
$$A = \begin{pmatrix} 0 & 6 & 7 \\ -6 & 0 & 8 \\ 7 & -8 & 0 \end{pmatrix}$$
,  $B = \begin{pmatrix} 0 & 1 & 1 \\ 1 & 0 & 2 \\ 1 & 2 & 0 \end{pmatrix}$ ,  $C = \begin{pmatrix} 2 \\ -2 \\ 3 \end{pmatrix}$ . Verify that  $(A + B)C = AC + BC$ .

(b) If 
$$A + B = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix}$$
 and  $A - 2B = \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}$  then show that  $A = \begin{pmatrix} \frac{1}{3} & \frac{1}{3} \\ \frac{2}{3} & \frac{1}{3} \end{pmatrix}$ 

[Class XII : Maths]

/ .

30. If 
$$A = \begin{pmatrix} i & 0 \\ 0 & -1 \end{pmatrix}$$
 and  $B = \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}$ , show that  $AB \neq BA$ 

31. Find a matrix X, for which 
$$\begin{pmatrix} 5 & 4 \\ 1 & 1 \end{pmatrix} \times = \begin{pmatrix} 1 & -2 \\ 1 & 3 \end{pmatrix}$$

- 32. If A and B are symmetric matrices, show that AB is symmetric, if AB = BA.
- 33. Match the following:

Possible Number of Matrices  $(A_n)$  of order  $3 \times 3$  with entry 0 or 1 which are

	Condition		No. of matrices
(1)	A <sub>n</sub> is diagonal Matrix	Р	<b>2</b> <sup>0</sup>
(2)	$A_n$ is upper triangular Matrix	Q	2 <sup>1</sup>
(3)	$A_n$ is identity Matrix	R	2 <sup>3</sup>
(4)	$A_n$ is scalar Matrix	S	2 <sup>6</sup>

34. If 
$$A = \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$$
 then prove that  $A^3 = \begin{pmatrix} \cos 3x & -\sin 3x \\ \sin 3x & \cos 3x \end{pmatrix}$ .

35. Express the following Matrices as a sum of a symmetric and skey-symmetric matrix. (Note: Part (b) and (c) can be asked for one marker, <u>SO THINK ABOUT THIS!</u>)

(a) 
$$A = \begin{pmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ -2 & -4 & -5 \end{pmatrix}$$
 (b)  $A = \begin{pmatrix} 1 & 2 & 5 \\ 2 & 5 & 7 \\ 5 & 7 & -5 \end{pmatrix}$  (c)  $A = \begin{pmatrix} 0 & 2 & -3 \\ -2 & 0 & 4 \\ 3 & -4 & 0 \end{pmatrix}$ 

36. Show that the Matrix  $A = \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}$  satisfies the equation  $A^2 - 4A + I = 0$ .

- 37. Find the values of x and y, if  $A = \begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$  satisfies the equation  $A^2 + xA + yI = 0$ .
- 38. Find f(A), if  $A = \begin{pmatrix} 2 & 3 \\ -1 & 1 \end{pmatrix}$  such that  $f(x) = x^2 3x + 5$
- 39. Find  $A^2$  if  $A = \begin{pmatrix} 5 & 4 \\ 1 & 1 \end{pmatrix}$ .
- 40. Find  $2A^2$  when  $x = \frac{\pi}{3}$  where  $A = \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$ .

#### **THREE MARKS QUESTIONS**

41. Let  $P = \begin{pmatrix} 1 & 0 & 0 \\ 3 & 1 & 0 \\ 9 & 3 & 1 \end{pmatrix}$  and  $Q = [q_{ij}]$  be two 3 × 3 matrices such that  $Q = P^5 + I_3$ , then Prove that  $\left(\frac{q_{21} + q_{31}}{q_{32}}\right) = 10.$ 

42. Construct a 3 × 3 matrix  $A = [a_{ij}]$  such that

(a) 
$$\mathbf{a}_{ij} = \begin{cases} i+j; & i>j\\ \frac{i}{j}; & i=j\\ i-j; & i  
(b)  $\mathbf{a}_{ij} = \begin{cases} 2^{i}; & i>j\\ i.j; & i=j\\ 3^{j}; & i  
(c)  $\mathbf{a}_{ij} = \begin{cases} i^{2}+j^{2}; & i\neq j\\ 0; & i=j \end{cases}$   
(d)  $\mathbf{a}_{ij} = \frac{|2i-3j|}{5}$$$$

(e)  $a_{ij} = \left[\frac{i}{j}\right]$ , where [.] represents Greatest Integer Function.

43. If 
$$A = \begin{pmatrix} \cos \theta & i \sin \theta \\ i \sin \theta & \cos \theta \end{pmatrix}$$
, then prove that  $A^2 = \begin{pmatrix} \cos 2\theta & i \sin 2\theta \\ i \sin 2\theta & \cos 2\theta \end{pmatrix}$ , where  $i = \sqrt{-1}$ 

44. If 
$$A = \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix}$$
, evaluate  $A^3 - 4A^2 + A$ .

45. If 
$$f(x) = \begin{pmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{pmatrix}$$
, then prove that  $f(x).f(y) = f(x + y)$ 

46. If 
$$f(x) = \frac{1}{\sqrt{1-x^2}} \begin{pmatrix} 1 & -x \\ -x & 1 \end{pmatrix}$$
, Prove that  $f(x).f(y) = f\left(\frac{x+y}{1+xy}\right)$ . Hence show that  $f(x).f(-x) = 1$ , where  $|x| < 1$ .

#### **FIVE MARKS QUESTIONS**

47. Find x, y and z if 
$$A^{T} = A^{-1}$$
 and  $A = \begin{pmatrix} 0 & 2y & z \\ x & y & -z \\ x & -y & z \end{pmatrix}$ . Also find how many triplets of (x, y, z) are possible. (NOTE:  $A \cdot A^{-1} = A^{-1}A = I$ )

48. If *A* is a symmetric Matrix and *B* is skew-symmetric Matrix such that  $A + B = \begin{pmatrix} 2 & 3 \\ 5 & -1 \end{pmatrix}$ 

then show that 
$$AB = \begin{pmatrix} 4 & -2 \\ -1 & -4 \end{pmatrix}$$
.  
49. If  $A = \begin{pmatrix} 4 & 1 \\ -9 & -2 \end{pmatrix}$  and  $A^{50} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$  then show that  $(a + b + c + d + 398) = 0$ .

## **CASE STUDIES**

50. (A) Two farmers Ramkishan and Gurcharan Singh cultivates only three varieties of rice namely Basmati, Permal and Naura. The Quantity of sale (in Kg) of these varieties of rice by both the farmers in the month of September and October are given by the following matrices *A* and *B*.



A(September sales) =	BASMATI = (1000 5000	PERMAI 2000 3000	NAURA 3000 1000	RAMAKRISHAN GURCHARAN SINGH
$B(\text{October sales}) = \begin{pmatrix} 1 \\ 2 \end{pmatrix}$	basmati f 5000 1 20000 1	PERMAL 0000 0000	NAURA 6000 10000	RAMAKRISHAN GURCHARAN SINGH

If Ramakrishan sell the variety of rice (per kg)i.e. Basmati, Permal and Naura at Rs.30, Rs. 20 & Rs.10 respectively, While Gurcharan Singh sell the variety of rice (per kg) i.e. Basmati, Permal and Naura at Rs. 40, Rs. 30, & Rs.20 respectively.

Based on the above information answer the following:

- (a) Find the Total selling Price received by Ramakrishan in the month of september.
- (b) Find the Total Selling Price received by Gurcharan Singh in the month of september.
- (c) Find the Total selling Price received by Ramakrishan in the month of september & october.

(B) A manufacture produces three stationery products Pencil, Eraser and Sharpener which he sells in two markets. Annual sales are indicated below



Market	Products (in numbers)						
	Pencil	Eraser	Sharpener				
А	10,000	2000	18,000				
В	600	20,000	8,00				

If the unit Sale price of Pencil, Eraser and Sharpener are Rs. 2.50, Rs. 1.50 and Rs. 1.00 respectively, and unit cost of the above three commodities are Rs. 2.00, Rs. 1.00 and Rs. 0.50 respectively, then, Based on the above information answer the following:

- (a) Find the total Revenue of both the markets.
- (b) Find the total Profit for both the markets.
- (C) Three schools ABC, PQR and MNO decided to organize a fair for collecting money for helping the flood victims. They sold handmade fans, mats and plates from recycled material at a cost of Rs. 25, Rs. 100 and Rs. 50 each respectively. The numbers of articles sold are given as



School/Article	ABC	PQR	MNO
Hand made fans	40	25	35
Marks	50	40	50
Plates	20	30	40

Based on the information given above, answer the following questions.

- (a) What is the total amount of money (in Rs.) collected by all the three schools ABC, PQR & MNO?
- (b) If the number of handmade fans and plates are interchanged for all the schools, then what is the total money collected by all schools?

#### **SELF ASSESSMENT-1**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. If A is a symmetric matrix then which of the following is not Symmetric matrix,
  - (a)  $A + A^{T}$  (b)  $A.A^{T}$
  - (c)  $A A^{T}$  (d)  $A^{T}$
- 2. Suppose *P*, *Q* and *R* are different matrices of order 3 × 5, *a* × *b* and *c* × *d* respectively, then value of *ac* + *bd* is, if matrix *P* + *Q R* is defined
  - (a) 9 (b) 14
  - (c) 24 (d) 34
- If A and B are two square matrices of same order such that, AB = A and BA = B, then (A + B) (A B) =
- (a) O (b) A(c)  $A^2 - B^2$  (d) B4. If  $\begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \begin{bmatrix} 6 \\ 3 \\ 2 \end{bmatrix}$ , then 2x + y - z =(a) 1 (b) 3(c) 5 (d) 7
- 5. If a matrix has 2022 elements, how many orders it can have?
  - (a) 6 (b) 2
    - (c) 4 (d) 8

#### SELF ASSESSMENT-2

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. If matrix  $A = [a_{ij}]_{2\times 2}$  where

	$\boldsymbol{a}_{ij} = \begin{cases} 1, & \text{if} \\ 0, & \text{if} \end{cases}$	$i \neq j$ , then $A^{2021}$ = $i = j$ ,	
	(a) O	(b	) A
	(c) –A	(d	) /
2.	$If A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}$	1 1 1 , then A⁴ = 1 ]	
	(a) <i>A</i>	(b	) 3A
	(c) 9 <i>A</i>	(d	) 27A

3. If 
$$A = \begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$$
 and  $A^2 + pA + qI = 0$ , then  $pq =$   
(a) 0 (b) 1  
(c) -1 (d) 2  
4. If  $\begin{bmatrix} 2a+b & a-2b \\ 5c-d & 4c+3d \end{bmatrix} = \begin{bmatrix} 4 & -3 \\ 11 & 24 \end{bmatrix}$ , then  $a + b + c + d =$   
(a) 0 (b) 4  
(c) 6 (d) 10  
5. If A is a square Matrix such that  $A^2 = A$ , then  $(I + A)^3 - 7A$  is equal to  
(a)  $2A + I$  (b)  $A + 2I$   
(c)  $I$  (d)  $A + I$ 

## ANSWER

## **One Mark Questions**

1. (b) $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$	2. (c) $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$
3. (b) + 1	4. (a) ±3
5. (b) 3	6. (d) 17
7. (a) 2023 × 2024	8. (d) 2024 × 2023
9. (a) 1	10. (a) 80
11. (b) 8	12. (d) 2
13. (d) 0	14. (b) 12
15. (b) x =2, y = 1	16. (a) $\begin{pmatrix} a^2 + b^2 & 0 \\ 0 & a^2 + b^2 \end{pmatrix}$
17. (b)A	18. (c) 4
19. (a) I	20. (b)(14)
21. (d) $\begin{pmatrix} -6 & -8 \\ -10 & -4 \end{pmatrix}$	22. (c)±5
23. (b) $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$	24. (d) A si false but R is true.
25. (a) Both A and R are true and	R is the correct explanation of A

## **Two Marks Questions**

31.	$X = \begin{pmatrix} -3 & -14 \\ 4 & 17 \end{pmatrix}$	33.	$(1) \to R  (2) \to S  (3)$	$B) \to P  (4) \to Q$
35. (a)	$ \begin{pmatrix} 1 & 2 & \frac{1}{2} \\ 2 & 5 & \frac{3}{2} \\ \frac{1}{2} & \frac{3}{2} & -5 \end{pmatrix} + \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ \frac{-5}{2} & \frac{-11}{2} \end{pmatrix} $			
35. (b)	$ \begin{pmatrix} 1 & 2 & 5 \\ 2 & 5 & 7 \\ 5 & 7 & -5 \end{pmatrix} + \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} $	35.	$(c) \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \begin{pmatrix} 0 \\ -2 \\ 3 \end{pmatrix}$	$ \begin{array}{ccc} 2 & -3 \\ 0 & 4 \\ -4 & 0 \end{array} \right) $
37. <i>x</i> =	2, <i>y</i> = 0	38.	$\begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$	$39. \begin{pmatrix} 29 & 24 \\ 6 & 5 \end{pmatrix}$
40. (	$\begin{pmatrix} 1 & -\sqrt{3} \\ \overline{3} & -1 \end{pmatrix}$			
	-	Thre	e Marks Questions	
42. (a)	$ \begin{pmatrix} 1 & -1 & -2 \\ 3 & 1 & -1 \\ 4 & 5 & 1 \end{pmatrix} $	42.	(b) $ \begin{pmatrix} 1 & 9 & 27 \\ 4 & 4 & 27 \\ 8 & 8 & 9 \end{pmatrix} $	42. (c) $\begin{pmatrix} 0 & 5 & 10 \\ 5 & 0 & 13 \\ 10 & 13 & 0 \end{pmatrix}$
40. (d)	$ \begin{pmatrix} \frac{1}{5} & \frac{4}{5} & \frac{7}{5} \\ \frac{1}{5} & \frac{2}{5} & 1 \\ \frac{3}{5} & 0 & \frac{3}{5} \end{pmatrix} $	42.	$(e) \begin{pmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 1 & 1 \end{pmatrix}$	$44. \begin{pmatrix} 0 & 0 \\ 0 & 0 \end{pmatrix}$
47. <i>x</i> =	$\pm \frac{1}{\sqrt{2}}, y = \pm \frac{1}{\sqrt{6}}, z = \pm \frac{1}{\sqrt{3}};$			
50. Cas	se Studv: A			
(a)	Rs. 1,00,000	(b)	Rs. 3,10,000	(c) Rs. 5,10,000

50.	Cas	e Study: B				
	(b)	Rs. 46,000 (F	or Market A)	(b) Rs. 15,000 (For	Market A)	
		Rs. 43,000 (R	or Market B)	Rs. 17,000 (For	<sup>-</sup> Market A)	
50.	Cas	e Study C:	50.	(iv) Option (d)	50. (v) Option (c	:)
	(a)	Rs. 21,000				
		Rs. 21,250				
			SEL	F ASSESSMENT-1		
1.	(c)	2.	(d)	3. (a)	4. (c)	5. (d)
			SEL	F ASSESSMENT-2		
1.	(b)	2.	(d)	3. (a)	4. (d)	5. (c)

# **CHAPTER-4**

# DETERMINANTS



One of the important aplications of inverse of a non-singular square matrix is in cryptography.

Cryptography is an art of communication between two people by keeping the information not known to others. It is based upon two factors, namely encryption and decryption.

<u>Encryption</u> means the process of transformation of an information (plain form) into an unreadable form (coded form). On the other hand, <u>Decryption</u> means the transformation of the coded message back into original form. Encryption and decryption require a secret technique which is known only to the sender and the receiver.

This secret is called a <u>key</u>. One way of generating a key is by using a non-singular matrix to encrypt a message by the sender. The receiver decodes (decrypts) the message to retrieve the original mesage by using the inverse of the matrix. The matrix used for encryption is called encryption matrix (encoding matrix) and that used for decoding is called decryption matrix (decoding matrix).

## TOPIC TO BE COVERED AS PER CBSE LATEST CURRICULUM (2024-25)

- Determinant of a square matrix (up to 3 × 3 matrice), minors, co-factors and applications of determinants in finding the area of a triangle.
- Adjoint and inverse of a square matrix.
- Consistency, inconsistency and number of solutions of system of linear equations by examples, solving system of linear equations in two or three variables (having unique solution) using inverse of a matrix.

A determinant of order 2 is written as  $|A| = \begin{vmatrix} a & b \\ c & d \end{vmatrix}$  where a, b, c, d are complex numbers (As

Complex Number Include Real Number). It denotes the complex number ad - bc.

Even though the value of determinnants Represented by Modulus symbol but the value of a determinant may be positive, negative or zero.

In other words,

. .

.

$$|A| = \begin{vmatrix} a & b \\ c & d \end{vmatrix} = ad - bc$$
 (Product of diagonal elements – Product of non-diagonal elements)

- Determinant of order 1 is the number itself.
- We can expand the determinants along any Row or Column, but for easier calculations we shall expand the determinant along that row or column which contains maximum number of zeroes.

## **MINORS AND COFATORS**

## **Minor of an Element**

If we take an element of the determinant and delete/remove the row and column containing that element, the determinant of the elements left is called the minor of that element. It is denoted by  $M_{i}$ . For example,

Let us consider a Determinant |A|

$$|A| = \begin{vmatrix} a & b & c \\ d & e & f \\ p & q & r \end{vmatrix} \Rightarrow$$

$$\begin{vmatrix} \widehat{a} & -b & -c \\ d & e & f \\ p & q & r \end{vmatrix} \Rightarrow M_{11} = \begin{vmatrix} e & f \\ q & r \end{vmatrix} (Minor of a_{11} = M_{11})$$

$$\begin{vmatrix} \widehat{a} & -b & -c \\ d & e & f \\ p & q & r \end{vmatrix} \Rightarrow M_{11} = \begin{vmatrix} e & f \\ q & r \end{vmatrix} (Minor of a_{11} = M_{11})$$

$$\begin{vmatrix} \widehat{a} & -b & -c \\ d & e & f \\ p & q & r \end{vmatrix} \Rightarrow M_{11} = \begin{vmatrix} e & f \\ q & r \end{vmatrix} (Minor of a_{11} = M_{11})$$

Hence a determinant of order two will have "4 minors" and a determinant of order three will have "9 minors".

## Minor of an Element:

Cofactor of the element  $a_{ij}$  is  $c_{ij} = (-1)^{i+j} M_{ij}$ ; where *i* and *j* denotes the row and column in which the particular element lies. (Means Magnitude of Minor and Cofactor of aij are equal).

• **Property:** If we multiply the elements of any row/column with their respective Cofactors of the same row/column, then we get the value of the determinant.

For example,

$$|A| = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$
$$|A| = a_{31}C_{31} + a_{32}C_{32} + a_{33}C_{33}$$

• **Property:** If we multiply the elements of any row/column with their respective Cofactors of the other row/column, then we get zero as a result.

For example,

 $a_{11}C_{21} + a_{12}C_{22} + a_{13}C_{23} = a_{11}C_{31} + a_{12}C_{32} + a_{13}C_{33}$ 

Note that the value of a determinant of order three in term sof 'Minor' and 'Cofactor' can be written as:

$$|A| = a_{11}M_{11} - a_{12}M_{12} + a_{13}M_{13} \text{ OR } |A| = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$
  

$$|A| = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13}$$
  
Clearly, we see that, if we apply the appropriate sign to the minor of an element, we have its  
Cofactor. The signs form a check-board pattern.  

$$\begin{vmatrix} + & - & + \\ - & + & - \\ + & - & + \end{vmatrix}$$

## **PROPERTIES OF DETERMINANTS**

• The value of a determinant remains unaltered, if the row and solumns are inter changed.

 $|A| = |A^{T}|$   $a \quad p \quad x \\ b \quad q \quad y \\ c \quad r \quad z \end{vmatrix} = \begin{vmatrix} a & b & c \\ p & q & r \\ x & y & z \end{vmatrix}$ 

• If any two rows (or columns) of a determinant be interchanged, the value of determinant is changed in sign only . e.g.

а	р	Х		а	Х	р		b	У	q
b	q	У	=	b	У	q	=	а	x	р
С	r	Ζ		с	Ζ	r		c	Ζ	r

• If all the elements of a row (or column) are zero, then the determinant is zero.

а	0	x		0	0	0	
b	0	y	=	p	q	r	= 0
с	0	z		x	у	z	

• If the all elements of a row (or column) are proportional (identical) to the elements sof some other row (or column), then the determinant is zero.

$$\begin{vmatrix} a & ka & x \\ b & kb & y \\ c & kc & z \end{vmatrix} = \begin{vmatrix} mp & mq & mr \\ p & q & r \\ x & y & z \end{vmatrix} = 0$$

• If all the elements of a determinant above or below the main diagonal consist of zeros (Triangular Matrix), then the determinant is equal to the product of diagonal elements.

а	0	0		а	Х	У		а	0	0	
Х	b	0	=	0	b	Ζ	=	0	b	0	= abc
y	Ζ	С		0	0	С		0	0	С	

• If all the elements of one row/column of a determinant are multiplied by "k" (A scalar), the value of the new determinant is k times the original determinant.

$$\begin{vmatrix} ka & p & x \\ kb & q & y \\ kc & r & z \end{vmatrix} = \begin{vmatrix} a & p & x \\ b & q & y \\ c & r & z \end{vmatrix}$$
$$\begin{vmatrix} ka & kp & x \\ kb & kq & y \\ kc & kr & z \end{vmatrix} = k^2 \begin{vmatrix} a & p & x \\ b & q & y \\ c & r & z \end{vmatrix}$$
$$\begin{vmatrix} ka & kp & kx \\ kb & kq & ky \\ kc & kr & kz \end{vmatrix} = k^3 \begin{vmatrix} a & p & x \\ b & q & y \\ c & r & z \end{vmatrix}$$

 $|kA| = k^n |A|$ , where *n* is the order of determinant.

## **AREA OF A TRIANGLE**

Area of a triangle whose vertices are  $(x_1, y_1), (x_2, y_2)$  and  $(x_3, y_3)$  is given by

$$A = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$
(sq. units)

## **ADJOINT OF A MATRIX**

Let  $A = [a_{ij}]_{m \times n}$  be a square matrix and  $C_{ij}$  be cofactor of  $a_{ij}$  in |A|.

Then, (adj A) = 
$$[C_{ij}] \Rightarrow adj A = \begin{pmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{pmatrix}$$

- *A*. (adj *A*) = (adj *A*).*A* = |*A*|
- (adj *AB*) = (adj *B*).(adj *A*)
- $|\text{adj } A| = |A|^{n-1}$ , where *n* is the order of a Matrix A

## SINGULAR MATRIX

A Matrix A is singular if |A| = 0 and it is non-singular if  $|A| \neq 0$ 

$$|A| = \begin{vmatrix} 2 & 3 \\ 1 & 4 \end{vmatrix} = 5 \neq 0. \text{ So } A \text{ is Non-singular Matrix.}$$
$$|A| = \begin{vmatrix} 2 & 8 \\ 1 & 4 \end{vmatrix} = 8 - 8 = 0. \text{ So } A \text{ is singular Matrix.}$$

## **INVERSE OF A MATRIX**

A square matrix *A* is said to be invertible if there exists a square matrix *B* of the same order such that AB = BA = I then we write  $A^{-1} = B$ ,  $(A^{-1}$  exists only if  $|A| \neq 0$ )

$$A^{-1} = \frac{1}{|A|} (adj A) = \frac{1}{|A|} \begin{pmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{pmatrix}$$

- $(AB)^{-1} = B^{-1}.A^{-1}$
- $(A^{-1})^{-1} = A$
- $(A^{T})^{-1} = (A^{-1})^{T}$
- $AA^{-1} = A^{-1}A = I$
- $|A^{-1}| = \frac{1}{|A|}$
- $|A.adj A| = |A|^n$  (Where *n* is the order of Matrix A)

Illustration:
For what value of <i>k</i> , the matrix $A = \begin{pmatrix} 2 & 10 \\ 5k - 2 & 15 \end{pmatrix}$ is singular matrix.
Solution: As, Matrix is singular, so its determinant will be zero.
A  = 2(15) - 10(5k - 2) = 30 - 50k + 20
A  = 50 - 50k = 0
$\Rightarrow$ 50k = 50
$\therefore$ $k = 1$

## Illustration:

Without expanding the determinants prove that  $\begin{vmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{vmatrix} = 0$ Solution: Let  $A = \begin{vmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{vmatrix}$ 

We observe here  $a_{ij} = -a_{ij}$  (*A* is skew-symmetric matrix)  $\Rightarrow A^{T} = -A$   $\Rightarrow |A^{T}| = |-A|$   $\Rightarrow |A| = (-1)^{3} |A|$ Property USED:  $|A^{T}| = |A|$ ,  $|kA| = k^{n}|A|$ Where *n* is the order of the determinant  $\Rightarrow |A| = -|A|$   $\Rightarrow 2|A| = 0$   $\Rightarrow |A| = \begin{vmatrix} 0 & a & -b \\ -a & 0 & -c \\ b & c & 0 \end{vmatrix} = 0$ 

#### Illustration:

If A is an invertible matrix of order 2 and |A| = 4, then write the value of  $|A^{-1}|$ .

Solution: As we know that,

$$|A^{-1}| = \frac{1}{|A|} = \frac{1}{4}$$
$$\Rightarrow |A^{-1}| = \frac{1}{4}$$

Illustration:Find the inverse of the matrix $\begin{pmatrix} 3 & 4 & 5 \\ 2 & -1 & 8 \\ 5 & -2 & 7 \end{pmatrix}$ and hence solve the system of equations:3x + 4y + 5z = 185x - 2y + 7z = 202x - y + 8z = 13

Solution: Let,  $A = \begin{pmatrix} 3 & 4 & 5 \\ 2 & -1 & 8 \\ 5 & -2 & 7 \end{pmatrix}$ Cofactors are.  $C_{11} = \begin{vmatrix} -1 & 8 \\ -2 & 7 \end{vmatrix} = -7 + 16 = 9$   $C_{21} = -\begin{vmatrix} 4 & 5 \\ -2 & 7 \end{vmatrix} = -38$   $C_{31} = \begin{vmatrix} 4 & 5 \\ -1 & 8 \end{vmatrix} = 37$  $C_{12} = -\begin{vmatrix} 2 & 8 \\ 5 & 7 \end{vmatrix} = -(14 - 40) = 26$   $C_{22} = \begin{vmatrix} 3 & 5 \\ 5 & 7 \end{vmatrix} = -4$   $C_{32} = -\begin{vmatrix} 3 & 5 \\ 2 & 8 \end{vmatrix} = -14$  $C_{13} = \begin{vmatrix} 2 & -1 \\ 5 & -2 \end{vmatrix} = -4 + 5 = 1 \qquad C_{23} = -\begin{vmatrix} 3 & 4 \\ 5 & -2 \end{vmatrix} = 26 \qquad C_{33} = \begin{vmatrix} 3 & 4 \\ 2 & -1 \end{vmatrix} = -11$  $\begin{vmatrix} Adj \ A = \begin{pmatrix} C_{11} & C_{21} & C_{31} \\ C_{12} & C_{22} & C_{32} \\ C_{13} & C_{23} & C_{33} \end{pmatrix} = \begin{pmatrix} 9 & -38 & 37 \\ 26 & -4 & -14 \\ 1 & 26 & -11 \end{pmatrix}$  $|A| = a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} = 3(9) + 4(26) + 5(1) = 27 + 104 + 5 = 136$ So,  $A^{-1} = \frac{1}{|A|} (Adj A) = \frac{1}{136} \begin{pmatrix} 9 & -38 & 37 \\ 26 & -4 & -14 \\ 1 & 26 & -11 \end{pmatrix}$ Given system of equation can be written as  $\begin{pmatrix} 3 & 4 & 5 \\ 2 & -1 & 8 \\ 5 & -2 & 7 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 18 \\ 20 \\ 13 \end{pmatrix} \implies \begin{pmatrix} 3 & 4 & 5 \\ 5 & -2 & 7 \\ 2 & -1 & 8 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 18 \\ 13 \\ 20 \end{pmatrix}$  $\Rightarrow A.X = B \Rightarrow A^{-1} AX = A^{-1}.B$  $IX = A^{-1}.B \Rightarrow X = A^{-1}.B$  $X = \frac{1}{136} \begin{pmatrix} 9 & -38 & 37 \\ 26 & -4 & -14 \\ 1 & 26 & -11 \end{pmatrix} \begin{pmatrix} 18 \\ 13 \\ 20 \end{pmatrix} = \frac{1}{136} \begin{pmatrix} 9 \times 18 - 38 \times 13 + 37 \times 20 \\ 26 \times 18 - 4 \times 13 - 14 \times 20 \\ 1 \times 18 + 26 \times 13 - 11 \times 20 \end{pmatrix}$  $X = \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{136} \begin{pmatrix} 408 \\ 136 \\ 126 \end{pmatrix} = \begin{pmatrix} 3 \\ 1 \\ 1 \\ 1 \end{pmatrix}$  $\Rightarrow$  So, x = 3, y = 1, z = 1

[Class XII : Maths]

77

## ONE MARK QUESTIONS

1. If 
$$f(x) = \begin{pmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{pmatrix}$$
, then determinant of  $\left(f\left(\frac{\pi}{6}\right), f\left(\frac{\pi}{3}\right)\right) =$   
(a) 0 (b) 1 (c) -1 (d)  $\frac{\pi}{2}$   
2. If for a square matrix  $A, A^2 - A + I = 0$ , then  $A^{-1}$  equal  
(a) A (b)  $I + A$  (c)  $A - I$  (d)  $I - A$   
3. If  $\begin{vmatrix} x & 3 & 4 \\ 1 & 2 & 1 \\ 1 & 4 & 1 \end{vmatrix} = 0$ , then value of x is  
(a) 0 (b) 1 (c) 4 (d) 2  
4. The value of  $\begin{vmatrix} x + y & y + z & z + x \\ z & x & y \\ 1 & 1 & 1 \end{vmatrix}$  is  
(a) xyz (b)  $(x + y + z)$  (c)  $2(x + y + z)$  (d) 0  
5. If  $A = \begin{pmatrix} 2 & 2023 & 2024 \\ 0 & 1 & 2022 \\ 0 & 0 & 5 \end{pmatrix}$ , then A. (adj A) equals  
(a) 2! (b) 1 (c) 5! (d) 10!  
6. If  $A = \begin{pmatrix} 3 & 1 \\ 19 & 7 \end{pmatrix}$ , then A.(adj A) equals  
(a)  $\begin{pmatrix} 3 & 1 \\ 19 & 7 \end{pmatrix}$  (b)  $\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}$  (c)  $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$  (d)  $\begin{pmatrix} 7 & -1 \\ -19 & 3 \end{pmatrix}$   
7. If the area of a triangle with vertices (-3, 0), (3, 0) and (0, k) is 9 sq. units, then |k| =

(b) 6

(c) 3

[Class XII : Maths]

(d) 9

-1 3

(a) 0

- 8. If the area of a triangle with vertices (2, –6), (5, 4) and (k, 4) is 35sq. units, then the sum of all possible values of k is
- (a) 2 (b) 10 (c) 12 (d) 14 9. If  $A = \begin{pmatrix} 2023 & 1 \\ 2024 & 1 \end{pmatrix}$ , then  $A^{-1} =$ (a)  $\begin{pmatrix} -2023 & 1 \\ 2024 & -1 \end{pmatrix}$ (b)  $\begin{pmatrix} -1 & 1 \\ 2024 & -2023 \end{pmatrix}$ (c)  $\begin{pmatrix} 1 & 1 \\ 1 & 1 \end{pmatrix}$ (d)  $\begin{pmatrix} -1 & -1 \\ -2024 & -2023 \end{pmatrix}$ 10. If  $A = \begin{pmatrix} k & 16 \\ -9 & -k \end{pmatrix}$  is singular matrix, then sum of all possible values of k is (a) 0 (b) 12 (c) 10 (d) 24 11. If  $A = \begin{pmatrix} k & 12 \\ 3 & 6 \end{pmatrix}$  is non-invertible matrix, then value of k is (a) 0 (b) 3 (c) 6 (d) 12 12. If  $A.(adjA) = \begin{pmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{pmatrix}$ , then |A| + |adjA| =(a) 5 (b) 10 (c) 25 30 (d) 13. If  $A.(adjA) = \begin{pmatrix} 2 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 2 \end{pmatrix}$ , then  $\frac{|A| + |A^{T}|}{|A^{-1}|} =$ (a) 2 (b) 8 (c) 4 (d) 16

## ASSERTION-REASON BASED QUSTIONS (Q. 14 & Q.15)

In the following questions, a statement of assertion (A) is following by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A are R are true and R is the correct explanation of A.
- (b) Both A and R are true but R is not the correct explnantion of A.
- (c) A is true but R is false
- (d) A is false but R is true

14. Assertion: For Matrix 
$$A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$$
, value of  $4C_{31} + 5C_{32} + 6C_{33}$  is 0.

Reasoning : The sum of the products of elements of any row of a matrix A with the cofactors of elements of other row is always equal to Zero.

15. Assertion : If  $A = \begin{pmatrix} 1 & 2 & 3 \\ 4 & 5 & 6 \\ 7 & 8 & 9 \end{pmatrix}$ , then determinant of matrix A is zero.

Reasoning : The determinant of a skew-symmetric matrix of order 3 × 3 is always zero.

## **TWO MARKS QUESTIONS**

17. Let *A* be a 3 × 3 matrix such that |A| = -2, then find the value of  $|-2A^{-1}| + 2|A|$ .

18. If 
$$A = \begin{pmatrix} a & b & c \\ x & y & z \\ p & q & r \end{pmatrix}$$
,  $B = \begin{pmatrix} yr - zq & cq - br & bz - cy \\ zp - xr & ar - cp & cx - az \\ xq - yp & bp - aq & ay - bx \end{pmatrix}$ . Find  $|B|$  if  $|A| = 4$ 

19. If 
$$A = \begin{pmatrix} a & b & c \\ x & y & z \\ p & q & r \end{pmatrix}$$
,  $B = \begin{pmatrix} yr - zq & cq - br & bz - cy \\ zp - xr & ar - cp & cx - az \\ xq - yp & bp - aq & ay - bx \end{pmatrix}$ . Find  $|A|$  if  $|B| = 25$ 

20. Find the Adjoint of Matrix A,

$$A = \begin{pmatrix} 2\cos\frac{\pi}{3} & -2\sin\frac{\pi}{3} \\ 2\sin\frac{\pi}{3} & -2\cos\frac{\pi}{3} \end{pmatrix}$$

#### **THREE MARKS QUESTIONS**

- 21. If A is a square matrix of order 3, such that |Adj A| = 25, then find the value of
  - (a) |A| (b)  $|-2A^{\tau}|$  (c)  $|4A^{-1}|$ (d) |5A| (e) A.Adj A (f) |A.Adj A|
  - (g)  $|A^3|$

22. If A is a square matrix of order 3, such that |A| = 5, then find the value of

	(a)  3 <i>A</i>	(b)  −2 <i>A</i> <sup>7</sup>	(c)  4 <i>A</i> <sup>-1</sup>
	(d)  Adj <i>A</i>	(e) A.Adj A	(f)  A.Adj A
	(f)  A <sup>3</sup>		
23.	If $A = \begin{pmatrix} 1 & 2020 & 2021 \\ 0 & 1 & 2022 \\ 0 & 0 & 3 \end{pmatrix}$ , $B = \begin{pmatrix} 1 & 2022 \\ 0 & 0 & 3 \end{pmatrix}$	$\begin{pmatrix} 2 & 0 & 0 \\ 2021 & 1 & 0 \\ 2020 & 2022 & 1 \end{pmatrix}$ then find	ind the value of
	(a)   <i>AB</i>	(b)  ( <i>AB</i> ) <sup>-1</sup>	(c)   <i>A</i> <sup>2</sup> . <i>B</i> <sup>3</sup>
	(d) $ 3(AB)^{T} $	(e)  Adj ( <i>AB</i> )	
24.	Find matrix 'X' such that $\begin{pmatrix} 2\\ 1 \end{pmatrix}$		
25.	Find matrix ' $X$ ' such that		

(a)	$\mathbf{X} \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} = \begin{pmatrix} 7 & -2 \\ 1 & 1 \end{pmatrix}$	(b) $\begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} X = \begin{pmatrix} 7 \\ 1 \end{pmatrix}$	-2 1
(c)	$ \begin{pmatrix} 2 & 3 \\ 1 & 2 \end{pmatrix} X \begin{pmatrix} 7 & -2 \\ 1 & 1 \end{pmatrix} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} $	0 1	

#### FOUR/FIVE MARKS QUESTIONS

- 26. (a) A school wants to award its students for regularity and hardwork with a total cash award of ₹ 6,000. If three times the award money for hardwork added to that given for regularity amounts of ₹ 11,000 represent the above situation algebraically and find the award money for each value, using matrix method.
  - (b) A shopkeeper has 3 varieties of pen A, B and C. Rohan purchased 1 pen of each variety for total of ₹ 21. Ayush purchased 4 pens of A variety, 3 pens of B variety and 2 pen of C variety for ₹ 60. While Kamal purchased 6 pens of A variety, 2 pens of B variety and 3 pen of C variety for ₹ 70. Find cost of each variety of pen by Matrix Method.
- 27. Find  $A^{-1}$ , where  $A = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 3 & 2 \\ 3 & -3 & -4 \end{pmatrix}$ . Hence use the result to solve the following system of

linear equations:

$$x + 2y - 3z = -4$$
  
2x + 3y + 2z = 2  
3x - 3y - 4z = 11

28. Find  $A^{-1}$ , where  $A = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 3 & 2 \\ 3 & -3 & -4 \end{pmatrix}$ . Hence, solve the system of linear equations: x + 2y + 3z = 8 2x + 3y - 3z = -3 -3x + 2y - 4z = -629. If  $A = \begin{pmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{pmatrix}$  and  $B = \begin{pmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{pmatrix}$  find *AB*. Hence using the product solve the system of eq. x - y + z = 4 x - 2y - 2z = 9 2x + y + 3z = 130. Find the product of matrices *AB*, where  $A = \begin{pmatrix} 1 & 3 & -2 \\ -3 & 0 & -1 \\ 2 & 1 & 0 \end{pmatrix}$ ,  $B = \begin{pmatrix} 1 & -2 & -3 \\ -2 & 4 & 7 \\ -3 & 5 & 9 \end{pmatrix}$  and use the result to solve following system of equations: x - 2y - 3z = 1

$$-2x + 4y + 5z = -1$$
  
$$-3x + 7y + 9z = -4$$

#### CASE STUDY BASED QUESTIONS

A. A family wanted to buy a home, but they wanted it to be close both to both the children's school and the parents' workplace. By looking at a map, they cold find a point that is equidistant from both the workplace and the school by finding the *circumcenter* of the triangular region.



If the coordinates are A(12, 5), B(20, 5) and C(16, 7), on the basis of this answer the following: (Figure is for reference only, Not as per scale)

- (a) Using the concept of Determinants. Find the equation of AC.
- (b) If any point P(2, k) is collinear with point A(12, 5) and O(16, 2), then find the value of (2k 15).
- (c) If any point P(2, k) is collinear with point A (12, 5) and 0 (16, 2), then find the value of (2k 15).
- B. For keeping Fit, X people believes in morning walk, Y people believes in yoga and Z people join Gym. Total no of people are 70. Further 20%, 30% and 40% people are suffering from any disease who believe in morning walk, yoga and GYM respectively. Total no. of such people is 21. If morning walk cost ₹ 0 Yoga cost ₹ 500/month and GYM cost ₹ 400/ month and total expenditure is ₹ 23000.



On the basis of above information, answer the following:

(a) If matrix 
$$A = \begin{pmatrix} 1 & 1 & 1 \\ 2 & 3 & 4 \\ 0 & 5 & 4 \end{pmatrix}$$
, then find  $A^{-1}$ .

- (b) On solving the given situational problem using matrix method, find the total number of person who prefer GYM.
- C. An amount of ₹ 600 crores is spent by the government in three schemes. Scheme A is for saving girl child from the cruel parents who don't want girl child and get the abortion before her birth.

Scheme *B* is for saving of newlywed girls from death due to dowry. Scheme *C* is planning for good health for senior citizen. Now twice the amount spent on Scheme *C* together with amount spent on Scheme *A* is ₹ 700 crores. And three times the amount spent on Scheme *A* together with amount spent on Scheme *B* and Scheme *C* is ₹ 1200 crores.

If we assume government invest (In crores)  $\notin X$ ,  $\notin Y$  and  $\notin Z$  in scheme *A*, *B* and *C* respectively. Solve the above problem using Matrices and answer the following:

C. Gautam buys 5 pens, 3 pens, 3 bags & 1 instrumental box and pays a sum of Rs. 160. From the same shop, Vikram buys 2 pens, 1 bag & 3 instrumental boxes and pays a sum of Rs. 190. Also Ankur buys 1 pen, 2 bags & 4 instrumental boxes and pays a sum of Rs. 250.

Based on above informatin answer the following questions:



- (a) Convert the given situation into a matrix equation of the form AX = B.
- (b) Find |A|.
- (c) Find  $A^{-1}$ .

OR

Determine  $P = A^2 - 5A$ 

#### SELF ASSESSMENT-1

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. If  $A = \begin{bmatrix} 2 & 3 & 6 \\ 0 & 1 & 8 \\ 0 & 0 & 5 \end{bmatrix}$ , then |A| =(a) 2 (b) 5 (c) 8 (d) 10 2. If  $A = \begin{bmatrix} 1 & 0 & 0 \\ 3 & 2 & 0 \\ 1 & 0 & 5 \end{bmatrix}$ , then  $|A^{T}| =$ (a) 2 (b) 5 (c) 8 (d) 10 3. If  $A = \begin{bmatrix} \cos x & -\sin x \\ \sin x & \cos x \end{bmatrix}$ , then  $|A^{-1}| =$ (a) 0 (b) 1 (c) cosx.sinx (d) –1 4. If  $A = \begin{bmatrix} 6x & 8 \\ 3 & 2 \end{bmatrix}$  is singular matrix, then the value of x is (a) 2 (b) 3 (c) 5 (d) 7 5. The area of a triangle with vertices (-3, 0), (3, 0) and (0, k) is 9 sq. units. The value of k will be (a) 6 (b) 9 (c) 3 (d) 0 **SELF ASSESSMENT-2** EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE COR-**RECT ALTERNATIVE** 1. If the value of a third order determinant is 12, then the value of the determinant formed by replacing each element by its co-factor will be (a) 0 (b) 1 (c) 12 (d) 144 2. If the points (3, -2), (x, 2), (8, 8) are collinear, then x =
  - (a) 2 (b) 5
  - (c) 4 (d) 3

3.	co   sii	s15⁰ n15⁰	sin75° cos75°	=						
	(a)	0			(b)	1				
	(c)	-1			(d)	2				
						1	2	3		
4.	The	mino	r of 6 in th	e determir	nant	4	5	6	is	5
						7	8	9		
	(a)	9			(b)	-6	6			
	(c)	6			(d)	10	)			
						Í	1	2	3	
5.	The	cofac	tor of 4 in	the deterr	ninaı	nt	4	5	6	is
							7	8	9	
	(a)	9			(b)	-6	6			
	(c)	6			(d)	10	)			

## ANSWER **One Mark Questions** 1. (b) 1 2. (d) I – A 3. (c) 4 6. (c) $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$ 4. (d) 0 5. (d) 10/ 9. (b) $\begin{pmatrix} -1 & 1 \\ 2024 & -2023 \end{pmatrix}$ 7. (c) 3 (b) 10 8. 10. (a) 0 11. (c)6 12. (d) 30 13. (b) 8 14. (a) Both A and R are true and R is the correct explanation of A. (b) Both A and R are true and R is not the correct explanation of A. **Two Marks Questions** 17.0 18. 16 19. ±5

 $20. \begin{pmatrix} -1 & \sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix}$ 

## **Three Marks Questions**

21. (a)  $\pm 5$  (b)  $\pm 40$  (c)  $\frac{\pm 64}{5}$  (d)  $\pm 625$  (e)  $\pm 5/$ (f)  $\pm 125$  (g)  $\pm 125$ 22. (a) 135 (b) -40 (c)  $\frac{64}{5}$  (d) 25 (e) 5/ (f) 125 (e) 125

23. (a) 6 (b)  $\frac{1}{6}$  (c) 72 (d) 162 (e) 36 24.  $X = \frac{1}{9} \begin{pmatrix} 2 & 31 \\ -1 & -11 \end{pmatrix}$ 

25. (a) 
$$X = \begin{pmatrix} 16 & -25 \\ 1 & -1 \end{pmatrix}$$
 (b)  $X = \begin{pmatrix} 11 & -7 \\ -5 & 4 \end{pmatrix}$  (c)  $X = \frac{1}{9} \begin{pmatrix} 5 & -17 \\ -3 & 12 \end{pmatrix}$ 

**Five Marks Questions** 

26. (a) Award money given for Honesty = ₹ 500

Regularity = ₹ 2000 and

Hard work = ₹ 3500  
(b) Cost of pen of  
Variety 
$$A = ₹ 5$$
  
Variety  $B = ₹ 8$  and  
Variety  $C = ₹ 8$   
27.  $x = 3, y = -2, z = 1$   
30.  $x = -4, y = -1, z = -1$   
28.  $x = 0, y = 1, z = 2$   
29.  $x = 3, y = -2, z = -1$ 

## CASE STUDIES QUESTIONS

A. (a) x – 2y = 2		(b)	10 sq. units	(c) 10
B. (a) $A^{-1} = \frac{-1}{6} \begin{pmatrix} -8 \\ -8 \\ 10 \end{pmatrix}$	1 4 -5	1 -2 1	(b)20	

C. (a) 
$$\begin{pmatrix} 5 & 3 & 1 \\ 2 & 1 & 3 \\ 1 & 2 & 4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 160 \\ 190 \\ 250 \end{pmatrix}$$
 (b)-22  
A X = B

$$(c) \frac{1}{-22} \begin{pmatrix} -2 & -10 & 8 \\ -5 & 19 & -13 \\ 3 & -7 & -1 \end{pmatrix}$$

OR	PA	PART				
(7	5	13)				
5	8	2				
8	3	3)				

		SELF ASSESSM	ENT-1	
1. (d)	2. (d)	3. (b)	4. (a)	5. (c)
		SELF ASSESSM	ENT-2	
1. (d)	2. (b)	3. (a)	4. (b)	5. (c)
# **CHAPTER 5**

# CONTINUITY AND DIFFERENTIABILITY



Many real life events, such as trajectory traced by Football where you see player hit the soccer ball, angle and the distance covered animation on the screen is shown to the viewers using technology can be described with the help of mathematical functions. The knowledge of Continuity and differentiation is popularly used in finding speed, directions and other parameters from a given function.

# CONTINUITY AND DIFFERENTIABILITY

Topics to be covered as per C.B.S.E. revised syllabus (2024-25)

- Continuity and differetiability
- Chain rule
- Derivative of inverse trigonometric functions, like  $\sin^{-1}x$ ,  $\cos^{-1}x$  and  $\tan^{-1}x$
- Derivative of implicit functions.
- Concept of exponential and logarithmic function
- Derivatives of logarithmic and exponential functions.
- Logarithmic differentiation, derivative of functions expressed in parametric forms.
- Second order derivatives.

## POINTS TO REMEMBER

• A function f(x) is said to be continuous at x = c iff  $\lim_{x \to c} f(x) = f(c)$ 

*i.e.*,  $\lim_{x \to c^-} f(x) = \lim_{x \to c^+} f(x) = f(c)$ 

- f(x) is continuous in (a, b) iff it is continuous at  $x = c \forall c \in (a, b)$ .
- *f*(*x*) is continuous in [*a*, *b*] iff
  - (i) f(x) is continuous in (a, b)
  - (ii)  $\lim_{x \to a^+} f(x) = f(a)$

(iii) 
$$\lim_{x \to b^-} f(x) = f(b)$$

- Modulus functions is Continuous on R
- Trigonometric functions are continuous in their respective domains.
- Exponential function is continuous on R
- Every polynomial function is continuous on R.
- Greatest integer function is continuous on all non-integral real numbers
- If f(x) and g(x) are two continuous functions at x = a and if  $c \in R$  then
  - (i)  $f(x) \pm g(x)$  are also continuous functions at x = a.
  - (ii) g(x) f(x), f(x) + c, cf(x), |f(x)| are also continuous at x = a.

(iii) 
$$\frac{f(x)}{g(x)}$$
 is continuous at  $x = a$ , provided  $g(a) \neq 0$ .

• A function f(x) is derivable or differentiable at x = c in its domain iff

$$\lim_{x \to c} \frac{f(x) - f(c)}{x - c} = \lim_{x \to c^+} \frac{f(x) - f(c)}{x - c}$$
, and is finite

The value of above limit is denoted by f'(c) and is called the derivative of f(x) at x = c.

$$\frac{d}{dx}(u\pm v) = \frac{du}{dx} + \frac{dv}{dx}$$

• 
$$\frac{d}{dx}(u,v) = u.\frac{dv}{dx} + v.\frac{du}{dx}$$
 (Product Rule)

• 
$$\frac{d}{dx}\left(\frac{u}{v}\right) = \frac{v \cdot \frac{du}{dx} - u \cdot \frac{dv}{dx}}{v^2}$$
 (Quotient Rule)

• If y= f(u) and u=g(t) then 
$$\frac{dy}{dt} = \frac{dy}{du} \times \frac{du}{dt} = f'(u)g'(t)$$
 (Chain Rule)

• If 
$$y = f(u)$$
,  $x = g(u)$  then,

$$\frac{dy}{dx} = \frac{dy}{du} \times \frac{du}{dx} = \frac{f'(u)}{g'(u)}$$

## Illustration:

Discuss the continuity of the function f(x) given by

$$f(x) = \begin{cases} 4 - x, & x < 4 \\ 4 + x, & x \ge 4 \end{cases} \text{ at } x = 4$$
  
Solution: We have  $f(x) = \begin{cases} 4 - x, & x < 4 \\ 4 + x, & x \ge 4 \end{cases}$   
LHL =  $\lim_{x \to 2^{i}} f(x) = \lim_{x \to 4^{-}} (4 - x) = \lim_{h \to 0^{-}} 4 - (4 - h) = 0$   
RHL =  $\lim_{x \to 4^{+}} f(x) = \lim_{x \to 4^{+}} (4 + x) = \lim_{h \to 0^{+}} 4 + (h + 4) = 8 + 0 = 8$   
Here LHL  $\neq$  RHL  
Hence  $f(x)$  is not continuous at  $x = 4$ 

Illustration:

Show that the function 
$$f(x)$$
 given by  

$$f(x) = \begin{cases} \frac{\tan x}{x} + \cos x, & x \neq 0 \\ 2, & x = 0 \end{cases}$$
is continuous at  $x = 0$ 
Solution: We have  $f(x) = \begin{cases} \frac{\tan x}{x} + \cos x, & x \neq 4 \\ 2, & x = 4 \end{cases}$ 
Now  $f(0) = 2$  ... (i)  
LHL =  $\lim_{x \to 0^-} f(x) = \lim_{x \to 0^-} \left( \frac{\tan x}{x} + \cos x \right) = \lim_{h \to 0^-} \frac{\tan(0-h)}{(o-h)} + \cos(0-h) = \lim_{h \to 0} \left[ \frac{-\tan h}{-h} + \cos h \right]$ 

$$= \lim_{h \to 0} \frac{\tan h}{h} + \lim_{h \to 0} \cos h = 1 + \cos(0) = 1 + 1 = 2 \qquad ...(ii)$$

[Class XII : Maths]

$$RHL = \lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{+}} \left( \frac{\tan x}{x} + \cos x \right) = \lim_{h \to 0^{+}} \frac{\tan(0+h)}{(0+h)} + \cos(0+h)$$
$$= \lim_{h \to 0} \frac{\tan h}{h} + \lim_{h \to 0} \cos h = 1 + \cos(0) = 1 + 1 = 2 \qquad \dots (iii)$$
$$LHL = RHL = f(0)$$
Hence  $f(x)$  is continuous at  $x = 0$ 

# ONE MARK QUESTIONS

# **Continuity and Differentiability**

This section comprises Multiple Choice Questions (MCQ) of one mark each

1. The value of k for which the function f given by

$$f(x) = \begin{pmatrix} \frac{k \cos x}{\pi - 2x}, & \text{if } x \neq \frac{\pi}{2} \\ 5, & \text{if } x = \frac{\pi}{2} \end{pmatrix} \text{ is continuous at } x = \frac{\pi}{2} \text{ is}$$
(a) 6
(b) 5
(c)  $\frac{5}{2}$ 
(d) 10
2. The value of k for which

$$f(x) = \begin{cases} 3x + 5, x \ge 2\\ k x^2, x < 2 \end{cases}$$
 is a continuous function is a

(a) 
$$\frac{-11}{4}$$
 (b)  $\frac{4}{11}$ 

(c) 11 (d) 
$$\frac{11}{4}$$

3. For what value of k, may the function  $f(x) = \begin{cases} k(3x^2 - 5x), x \le 0\\ \cos x, x > 0 \end{cases}$  becomes continuous ?

- (a) 0 (b) 1
- (c)  $\frac{-1}{2}$  (d) No value

4. If  $f(x) = \begin{cases} \frac{\sin \pi x}{5x}, x \neq 0\\ k, x = 0 \end{cases}$  is continuous at x = 0, then k is equal to : (a)  $\frac{5}{\pi}$ (b)  $\frac{\pi}{5}$ (c) 1 (d) 0 5. if  $f(x) = \begin{cases} \frac{\sqrt{x^2 + 5 - 3}}{x + 2}, & x \neq -2 \\ k & x = -2 \end{cases}$  is continuous at x = -2, then the value of k is equal to : (a)  $\frac{-2}{3}$ (b) 0 (c)  $\frac{2}{2}$ (d) none of these 6. If  $f(x) \begin{cases} \frac{\tan\left(\frac{\pi}{4} - x\right)}{\cot 2x}, x \neq \frac{\pi}{4} \text{ is continuous at } x = \frac{\pi}{4}, \text{ then the value of k is :} \\ k , x = \frac{\pi}{4} \end{cases}$ (a) 1 (b) 2 (c)  $\frac{1}{2}$ (d) none of these 7. The number of points of discontinuity of the rational function  $f(x) = \frac{x^2 - 3x + 2}{4x - x^3}$  is : (a) 1 (b) 2 (c) 3 (d) none of these 8. The function f(x) = [x], where [x] denotes the greatest integer function, is continuous at x = (a) -2 (b) 1 (c) 4 (d) 1.5 9. The function f(x) = |x| at x = 0 is : (a) continuous but not differentiable (b) differentiable but not continuous (c) continuous and differentiable (d) neither continuous nor differentiable

[Class XII : Maths]

- 10. The function f(x) = |x| + |x-1| is :
  - (a) differentiable at x = 0 but not at x = I
  - (b) differentiable at x = I but not at x = 0
  - (c) neither differentiable at x = 0 nor at x = 1
  - (d) differentiable at x = 0 as well as at x = 1

11. The set of numbers where the function f given by  $f(x) = |2x - 1| \cos x$  is differentiable is:

- (a) R (b)  $R \left(\frac{1}{2}\right)$
- (c)  $(0,\infty)$  (d) none of these

12. If 
$$y = \log\left(\frac{1-x^2}{1+x^2}\right)$$
,  $|x| < 1$  then  $\frac{dy}{dx} =$ 

(a) 
$$\frac{4x^3}{1-x^4}$$
 (b)  $\frac{-4x}{1-x^4}$ 

(c) 
$$\frac{1}{4-x^4}$$
 (d)  $\frac{-4x^3}{1-x^4}$ 

13. The derivative of sec(tan<sup>-1</sup>x) w.r.t. x is

(a) 
$$\frac{x}{1+x^2}$$
 (b)  $\frac{1}{\sqrt{1+x^2}}$   
(c)  $\frac{x}{\sqrt{1+x^2}}$  (d)  $x\sqrt{1+x^2}$ 

14. If 
$$y = \cos^{-1}\left(\frac{\sqrt{x}-1}{\sqrt{x}+1}\right) + \csc^{-1}\left(\frac{\sqrt{x}+1}{\sqrt{x}-1}\right)$$
 then  $\frac{dy}{dx}$  is equal to :

(a)  $\frac{\pi}{2}$  (b) 0

15. Differential of log [log (log x<sup>5</sup>)] w.r.t. x is :

(a) 
$$\frac{5}{x \log(x^5) \log(\log x^5)}$$
 (b) 
$$\frac{5}{x \log(\log x^5)}$$
  
(c) 
$$\frac{5x^4}{\log(x^5) \log(\log x^5)}$$
 (d) 
$$\frac{5x^4}{\log(\log x^5)}$$

16. If  $y = \sin(m \sin^{-1} x)$  then which of the following equations is true?

- (a)  $(1-x^2)\frac{d^2y}{dx^2} + x\frac{dy}{dx} + m^2y = 0$
- (b)  $(1-x^2)\frac{d^2y}{dx^2} x\frac{dy}{dx} + m^2y = 0$
- (c)  $(1+x^2)\frac{d^2y}{dx^2} x\frac{dy}{dx} m^2y = 0$

(d) 
$$(1+x^2)\frac{d^2y}{dx^2} + x\frac{dy}{dx} - m^2y = 0$$

17. If 
$$y = \sqrt{\sin x + y}$$
, then  $\frac{dy}{dx}$  is equal to :

(a) 
$$\frac{\cos x}{2y-1}$$
 (b)  $\frac{\cos x}{1-2y}$ 

(c) 
$$\frac{\sin x}{1-2y}$$
 (d)  $\frac{\sin x}{2y-1}$ 

- Q no (18-22) are Assertion Reason Based questions carrying one mark each. These type of questions consists of two statements, one labelled Assertion (A) and other labelled Reason (R). Select the correct answer from the codes (a), (b),(c), and (d) as given below.
  - (a) Bolh Assertion (A) and Reason (R)arc true and Reason (R) is the correct explanation of the Assertion (A).
  - (b) Both Assertion (A) and Reason (R)arc true and Reason (R) is **not** the correct explanation of the Assertion (A).
  - (c) Assertion (A) is true and Rcason(R) is false.
  - (d) Assertion (A) is false and Reason(R) is true

18. Let  $f(x) = \frac{1}{1-x} - \frac{3}{1-x^3}, x \neq 1$ 

Statement -I: The value of f(1) so that f is continuous function is 1

**Statement-II** :  $g(x) = \frac{x+2}{x^2+x+1}$  is continuous function

Answer (d) Assertion (A) is false and Reason(R) is true

19. Consider the function  $f(x) = |x - 2| + |x - 5|, x \in R$ Statement - I : f'(4)=0

[Class XII : Maths]

**Statement -II :** f is continuous on [2, 5] differentiable on (2, 5) and f(2) = f(5)

Solution (b) Both Assertion (A) and Reason (R) are true and Reason (R) is **not** the correct explanation of the Assertion (A).

20. Statement -I:  $f(x) = \begin{cases} x^2 \sin \frac{1}{x}, & x \neq 0 \\ 0, & x = 0 \end{cases}$ 

Statement-II : Both  $h(x) = x^2$  and  $g(x) = \begin{cases} \sin \frac{1}{x}, x \neq 0\\ 0, x = 0 \end{cases}$  continuous at x = 0

21. F(x) is defined as the product of two real functions  $f_x(x) = x \quad \forall x \in R$  and

$$f_2(x) = \begin{cases} x \sin \frac{1}{x}, x \neq 0 \text{ as follows} \\ 0, x = 0 \end{cases}$$

$$F(x) = \begin{cases} f_1(x), f_2(x) \text{ if } x \neq 0\\ 0, \text{ if } x = 0 \end{cases}$$

Statement -I: F(x) is continuous on R

Statement-II :  $f_1(x)$  and  $f_2(x)$  are continuous on R

22. Let f(x) be a differentiable function such that f(2)=4 and f'(2)=4

Statement -I: 
$$\lim_{x \to 2} \frac{xf(2) - 2f(x)}{x - 2} = -4$$

Statement -II :  $f'(a) = \lim_{x \to a} = \frac{f(x) - f(a)}{x - a}$ 

#### CASE BASED

23. A plotter made a mud vessel , where the shape of pot is based on f(x) = |x-3| + |x-2|, where f(x) represents the height of the pot.



[Class XII : Maths]

#### Based on the information given above answer the following questions

- (1) When x > 4 what will be the height in terms of x?
- (2) When the value of x lies between (2, 3) then find the value of f(x).
- (3) If the potter is trying to make pot using the function f(x)=[x], will he get a pot or not? why?
- **Q24.** Let x = f(t) and y = g(t) be the parametric forms with t as parameter, then dy = dy dt = g'(t)

$$\frac{dy}{dx} = \frac{dy}{dt} \cdot \frac{dt}{dx} = \frac{g(t)}{f'(t)}$$
 where  $f'(t) \neq 0$ 

On the basis of the above information answer the following questions :

- (1) What will be the derivative of f(tanx) w.r.t g(secx) at  $x = \frac{\pi}{4}$  where f'(1) and  $g'(\sqrt{2}) = 4$ ?
- (2) Find the derivative of  $\cos^{-1}(2x^2 1)$  w.r.t  $\cos^{-1}x$ .

(3) If 
$$y = \frac{1}{4}u^4$$
 and  $u = \frac{2}{3}x^3$  then find  $\frac{dy}{dx}$ 

25. A function f(x) is said to be differentiable at x=c if

(i) Left hand derivative (L.H.D) = 
$$f'(c) = \lim_{h \to 0^-} \frac{f(c+h) - f(c)}{h}$$
 exists finitely.

- (ii) Right hand derivative (R.H.D) =  $f'(c) = \lim_{h \to 0^+} \frac{f(c+h) f(c)}{h}$  exists finitely.
- (iii) R.H.D = L.H.D, i.e. if the function f(x) is differentiable at x = c, then  $f'(c) = \lim_{h \to c} \frac{f(x) f(c)}{x c}$

#### Based on the above information answer the following :

(1) If f(x) is differentiable at x = 3. then find the value of  $\lim_{h \to 3} \frac{x^2 f(3) - 9f(x)}{x - 3}$ 

(2) Find 
$$\lim_{h\to 0} \frac{f(x+h)-f(x-h)}{h}$$
 if it exists.

# **TWO MARKS QUESTIONS**

1. Differentiate sin (x<sup>2</sup>) w. r. t. 
$$e^{\sin x}$$
  
2.  $y = x^{y}$  then find  $\frac{dy}{dx}$   
3. If  $y = x^{x} + x^{3} + 3^{x} + 3^{3}$ , find  $\frac{dy}{dx}$   
4. If  $y = 2\sin^{-1}(\cos x) + 5 \csc^{-1}(\sec x)$ . Find  $\frac{dy}{dx}$   
5. If  $y = e^{[\log (x+1) - \log x]}$  find  $\frac{dy}{dx}$   
6. Differentiate Sin<sup>-1</sup> [x  $\sqrt{x}$ ] w. r. t. x.  
7 Find the derivative of |x<sup>2</sup>+2| w.r.t. x  
8. Find the domain of the continuity of f(x) = sin<sup>-1</sup>x - [x]  
9. Find the derivative of cos (sin x<sup>2</sup>) w.r.t. x at  $x = \sqrt{\frac{p}{2}}$   
10. If  $y=e^{3\log x+2x}$ , Prove that  $\frac{dy}{dx} = x^{2}(2x+3)e^{2x}$ .  
11. Differentiate sin<sup>2</sup>( $\theta^{2}+1$ ) w.r.t.  $\theta^{2}$   
12. Find  $\frac{dy}{dx}$  if  $y = \sin^{1/4}\left(\frac{\sqrt{x}-1}{\sqrt{x}+1}\right) + \sec^{-1}\left(\frac{\sqrt{x}+1}{\sqrt{x}-1}\right)$ 

13. If 
$$x^2 + y^2 = 1$$
 verify that  $\frac{dy}{dx} \cdot \frac{dx}{dy} = 1$ 

14. Find 
$$\frac{dy}{dx}$$
 when  $y = 10^{x^{10^x}}$ 

15. If 
$$y = x^x$$
 find  $\frac{d^2y}{dx^2}$ 

16. Find 
$$\frac{dy}{dx}$$
 if y = cos<sup>-1</sup> (sinx)

17. If 
$$f(x) = x + 7$$
, and  $g(x) = x - 7$ ,  $x \in R$ , them find  $\frac{d}{dx}$  (fog) (x).

18. Differntiate log (7 logx) w.r.t x

19. If 
$$y = f(x^2)$$
 and  $f'(x) = sinx^2$ . Find  $\frac{dy}{dx}$ 

20. Find 
$$\frac{dy}{dx}$$
 if  $y = \sqrt{\sin^{-1}\sqrt{x}}$ 

# **THREE MARKS QUESTIONS**

1. Examine the continuity of the following functions at the indicated points.

(I) 
$$f(x) = \begin{cases} x^2 \cos(\frac{1}{x}), & x \neq 0 \\ 0, & x = 0 \end{cases}$$
 at  $x = 0$ 

(II) 
$$f(x) = \begin{cases} x - [x], x \neq 1 \\ 0, x = 1 \end{cases}$$
 at  $x = 1$ 

(III) 
$$f(x) = \begin{cases} \frac{e^{\frac{1}{x}} - 1}{e^{\frac{1}{x}} + 1} & , x \neq 0 \\ 0 & , x = 0 \end{cases}$$
 at  $x = 0$ 

(IV) 
$$f(x) = \begin{cases} \frac{x - \cos(\sin^{-1}x)}{1 - \tan(\sin^{-1}x)} & x \neq \frac{1}{\sqrt{2}} \\ \frac{-1}{\sqrt{2}} & x = \frac{1}{\sqrt{2}} \end{cases}$$

2. For what values of constant K, the following functions are continuous at the indicated points.

(i) 
$$f(x) = \begin{cases} \frac{\sqrt{1+Kx} - \sqrt{1-Kx}}{x} & x < 0\\ \frac{2x+1}{x-1} & x > 0 \end{cases} \text{ at } x = 0$$

(ii) 
$$f(x) = \begin{cases} \frac{e^x - 1}{\log(1 + 2x)} & x \neq 0 \\ K & x = 0 \end{cases}$$
 at  $x = 0$ 

(iii) 
$$f(x) = \begin{cases} \frac{1 - \cos 4x}{x^2} & x < 0\\ K & x = 0\\ \frac{\sqrt{x}}{\sqrt{16 + \sqrt{x} - 4}} & x > 0 \end{cases} \text{ at } x = 0$$

[Class XII : Maths]

3. For what values a and b

$$f(x) = \begin{cases} \frac{x+2}{|x+2|} + a & \text{if } x < -2\\ a+b & \text{if } x = -2\\ \frac{x+2}{|x+2|} + 2b & \text{if } x > -2 \end{cases}$$

Is continuous at x = -2

4. Find the values of a, b and c for which the function

$$f(x) = \begin{cases} \frac{\sin[(a+1)x] + \sin x}{x} & x < 0\\ C & x = 0\\ \frac{\sqrt{x+bx^2} - \sqrt{x}}{bx^{3/2}} & x > 0 \end{cases}$$

Is continuous at x = 0

5. 
$$f(x) = \begin{cases} [x] + [-x] \ x \neq 0 \\ \lambda & x = 0 \end{cases}$$

Find the value of  $\lambda$ , *f* is continuous at = 0 ?

6. Let 
$$f(x) = \begin{cases} \frac{1-\sin^3 x}{3\cos^2 x}; & x < \frac{\pi}{2} \\ a; & x = \frac{\pi}{2} \\ \frac{b(1-\sin x)}{(\pi-2x)^2}; & x > \frac{\pi}{2} \end{cases}$$
  
If  $f(x)$  is continuous at  $x = \frac{\pi}{2}$ , find *a* and *b*.

7. If 
$$f(x) = \begin{cases} x^3 + 3x + a \, x \le 1 \\ bx + 2 & x > 1 \end{cases}$$

Is everywhere differentiable, find the value of *a* and *b*.

8. Find the relationship between a and b so that the function defined by

$$f(x) = \begin{cases} ax+1 , x \le 3 \\ bx+3 , x > 3 \end{cases}$$
 is continuous at  $x = 3$ .

9. Differentiate 
$$\tan^{-1}\left(\frac{\sqrt{1-x^2}}{x}\right)$$
 w.r.t  $\cos^{-1}\left(2x\sqrt{1-x^2}\right)$  where  $x \neq 0$ .

10. If 
$$y = x^{x^x}$$
, then find  $\frac{dy}{dx}$ .

11. Differentiate 
$$(x \cos x)^x + (x \sin x)^{\frac{1}{x}}$$
 w.r.t. x.

12. If 
$$(x + y)^{m+n} = x^m \cdot y^n$$
 then prove that  $\frac{dy}{dx} = \frac{y}{x}$ 

13. If 
$$(x - y) \cdot e^{\frac{x}{x - y}} = a$$
, prove that  $y\left(\frac{dy}{dx}\right) + x = 2y$ 

14. If 
$$x = \tan\left(\frac{1}{a}\log y\right)$$
 then show that

$$(1+x^2)\frac{d^2y}{dx^2} + (2x-a)\frac{dy}{dx} = 0$$

15. If 
$$y = x \log\left(\frac{x}{a+bx}\right)$$
 prove that  $x^3 \frac{d^2y}{dx^2} = \left(x \frac{dy}{dx} - y\right)^2$ .

16. Differentiate 
$$\sin^{-1}\left[\frac{2^{x+1}.3^x}{1+(36)^x}\right]$$
 w.r.t *x*.

17. If 
$$\sqrt{1 - x^6} + \sqrt{1 - y^6} = a(x^3 - y^3)$$
, prove that  
 $\frac{dy}{dx} = \frac{x^2}{y^2} \sqrt{\frac{1 - y^6}{1 - x^6}}$ , Where  $-1 < x < 1$  and  $-1 < y < 1$  [HINT: put  $x^3 = \sin A$  and  
 $y^3 = \sin B$ ]  
18. If  $f(x) = \sqrt{x^2 + 1}$ ,  $g(x) = \frac{x + 1}{x^2 + 1}$  and  $h(x) = 2x - 3$  find  $f'[h'(g'(x))]$ .  
19. If  $x = \sec \theta - \cos \theta$  and  $y = \sec^n \theta - \cos^n \theta$ , then prove that  $\frac{dy}{dx} = n \sqrt{\frac{y^2 + 4}{x^2 + 4}}$ 

20. If 
$$x^{y} + y^{x} + x^{x} = m^{n}$$
, then find the value of  $\frac{dy}{dx}$ .

21. If 
$$x = a \cos^3 \theta$$
,  $y = a \sin^3 \theta$  then find  $\frac{d^2 y}{dx^2}$  at  $x = \frac{\pi}{6}$ 

[Class XII : Maths]

22. If 
$$y = \tan^{-1} \left[ \frac{\sqrt{1 + \sin x} - \sqrt{1 - \sin x}}{\sqrt{1 + \sin x} + \sqrt{1 - \sin x}} \right]$$
 where  $0 < x < \frac{\pi}{2}$  find  $\frac{dy}{dx}$ 

23. If 
$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$
, then show that  $\frac{d^2y}{dx^2} = -\frac{b^4}{a^2y^3}$ .

24. If 
$$= [x + \sqrt{x^2 + 1}]^m$$
, show that  $(x^2 + 1)y_2 + xy_1 - m^2y = 0$ .

25. If 
$$x^y = e^{x-y}$$
, prove that  $\frac{dy}{dx} = \frac{\log x}{(1+\log x)^2}$ 

26. If 
$$y^{1/m} + y^{-1/m} = 2x$$
 then prove that  $(x^2 - 1)y_2 + xy_1 = m^2 y$ .

## **SELF ASSESSMENT-1**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ONE

- 1. If  $y = \sin^2 x \cos^2 x$ , then  $\frac{dy}{dx} =$ (a)  $2 \sin x$  (b)  $2 \cos x$ (c)  $2 \sin 2x$  (d)  $-2 \sin 2x$
- 2. The value of 4k' for which the function f(x) is continuous at x = 3.

$f(x) = \begin{cases} \frac{(x+3)^2 - 36}{x-3}, \end{cases}$	when $x \neq 3$			
$\lfloor 2k+1, \rfloor$	when $x = 3$			
(a) 4		(b)	6	
(c) 11		(d)	22	
Derivative of $\sin x$ with respect to $\cos x$ is				

(c)  $\cot x$  (d)  $-\cot x$ 

3.

4. If 
$$y = (x + \sqrt{1 + x^2})^n$$
, then  $(1 + x^2)\frac{d^2y}{dx} + x\frac{dy}{dx} =$   
(a)  $n^2y$  (b)  $ny$   
(c)  $y$  (d)  $-ny$   
5. If  $x = a(\cos\theta + \theta\sin\theta), y = a(\sin\theta - \theta\cos\theta)$  then  $\frac{d^2y}{dx^2} =$   
(a)  $\frac{\sec^3\theta}{a}$  (b)  $\frac{\sec^3\theta}{a\theta}$   
(c)  $\sec^3\theta$  (d)  $\theta\sec^3\theta$ 

#### **SELF ASSESSMENT-2**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ONE

1. A Function defined as

 $f(x) = \begin{cases} |x| - 3, & \text{when } x < 0 \\ 5 - |x|, & \text{when } x \ge 0 \end{cases}$ is continuous on (b)  $R - \{0\}$ (a) R (c) [0,∞) (d) (−∞, 0] 2. The function  $g(x) = (\sin x + \cos x)$  is continuous at (a) R (b)  $R - \{0\}$ (c)  $R - \left\{\frac{\rho}{2}\right\}$ (d)  $R - \{\pi\}$ 3. The value of the derivative of |x-2| + |x-3| at x = 2 is (b) 3 (a) 1 (c) 2 (d) 0 4. If  $\sin y = x \cdot \cos(a + y)$  then  $\frac{dy}{dx}$ ? (b)  $\frac{\cos^2(a+y)}{\sin a}$ (a)  $\frac{\cos^2(a+y)}{\cos a}$ (c)  $\frac{\sin^2(a-y)}{\cos a}$ (d)  $\frac{\sin^2(a+y)}{\sin a}$ 5. If  $y = \left(\frac{x^a}{x^b}\right)^{a-b} \cdot \left(\frac{x^b}{x^c}\right)^{b+c} \cdot \left(\frac{x^c}{x^a}\right)^{c+a}$ , then  $\frac{dy}{dx} =$ (b) abc (a) 1 (c) a + b + c (d) 0

[Class XII : Maths]

# **ANSWERS**

# **ONE MARK QUESTIONS**

1.	(d) 10		
2.	(d) $\frac{11}{4}$		
3.	(d) No value		
4.	(b) $\frac{\pi}{5}$		
5.	(a) $\frac{-2}{3}$		
6.	(c) $\frac{1}{2}$		
7.	(c) 3		
8. 9. 10.	<ul> <li>(d) 1.5</li> <li>(c) continuous and differentiable at</li> </ul>	itiable x = 0 r	nor at x = 1
11.	(b) $R - \left\{ \frac{1}{2} \right\}$		
12.	(b) $\frac{-4x}{1-x^4}$	13.	(c) $\frac{x}{\sqrt{1+x^2}}$
14.	(b) 0	15.	(a) $\frac{5}{x \log(x^5) \log(\log x^5)}$
16.	(b) $(1-x^2)\frac{d^2y}{dx^2} - x\frac{dy}{dx} + x^2$	$y^{2} = 0$	
17.	(a) $\frac{\cos x}{2y-1}$	18.	(a) $\frac{x}{\sqrt{1+x^2}}$

## **ASSERTION REASONING**

- 18. Answer (d) Assertion (A) is false and Reason (R) is true
- 19. (b) Both Assertion (A) and Reason (R) are true and Reason (R) is not the correct explanation of the Assertion (A).
- 20. Solution (c) Assertion (A) and Reason (R) is false
- 21. Ans (a) Both Assertion (A) is true and Reason (R) are true.
- 22. (a) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A)

## **CASE BASED QUESTIONS**

23. (a) f(x) = 2x - 5(b) f(x) = 1

(c) Since the function is not continuous he will not get a pot.

24. (a) 
$$\frac{1}{\sqrt{2}}$$
 (b) 2 (c)  $\frac{16}{27}x^{11}$ 

25. (a) 9f' (3) + 6f(3) (b) 
$$2f'(x)$$

# **TWO MARKS QUESTIONS**

1.	$\frac{2x\cos(x^2)}{\cos(x^2)}$	11.	$\sin(2\theta^2+2), \theta\neq 0$
	$\cos x e^{-xxx}$	12.	0
2.	$x[1-y\log x]$	14.	10 <sup>x<sup>10<sup>x</sup></sup></sup> 10 <sup>x</sup> log10(1 – x log10)
3.	$x^{x} [1 + \log x] + 3x^{2} + 3^{x} \log_{e} 3$	15.	$x^{x}[1-\log x]$
4.	-7	16	-1
5.	$-\frac{1}{x^2}$	17.	1
6.	$\frac{3}{2}\sqrt{\frac{x}{1-x^3}}$	18.	$\frac{1}{x \log x}$
7.	$\frac{2x(x^2+2)}{ x^2+2 }$	19.	2x sinx⁴
8.	$(-1,0) \cup (0,1)$		1
9.	0	20.	$\frac{1}{4\sqrt{x}\sqrt{1-x}\sqrt{\sin^{-1}\sqrt{x}}}$ , where $0 < x < 1$

# THREE MARKS QUESTIONS

1.	(I)	Continuous		(II)	Discontinuous
	(III)	Not Continuous at $x =$	: 0	(IV)	Continuous
2.	(I)	K = -1		(II)	$K = \frac{1}{2}$
	(111)	K = 8			-
3.	<i>a</i> = 0,	b = -1			
4.	$a = \frac{-3}{2}$	, $b = R - \{0\}, \ c = \frac{1}{2}$			
5.	$\lambda = -i$	1			
6.	$a=\frac{1}{2},$	b = 4			
7.	a = 3,	b = 5			
8.	3a — 3	b = 2			
9.	$-\frac{1}{2}$				
10.	$x^x x^{x^x}$	$\left\{ (1 + \log x) \log x + \frac{1}{x} \right\}$			
11.	$(x\cos x)$	$(x)^x [1 - x \tan x + (\log x)]$	$(\cos x)] +$	$(x \sin x)$	$)^{1/x} \left[ \frac{1 + x \cot x - \log(x \sin)^x}{x^2} \right]$
16.	$\left[\frac{2^{x+1}3}{1+(36)}\right]$	$\left[\frac{x}{x}\right]\log 6$			
18.	$\frac{2}{\sqrt{5}}$				
20.	$\frac{dy}{dx} = \frac{y}{dx}$	$\frac{x^{x}(1+\log x)+yx^{y-1}-y^{x}\log x}{x^{y}\log x+xy^{x-1}}$	<u>g y</u>		
21.	$\frac{32}{27a}$				
22.	$-\frac{1}{2}$				
		SELF ASS	SESS	IENT	TEST-1
1. (C)		2. (C)	3. (D)		4. (A) 5. (B)

# SELF ASSESSMENT TEST-2

1. (D) 2. (A) 3. (C) 4. (A) 3. (D)	1. (B)	2. (A)	3. (C)	4. (A)	5. (D)
------------------------------------	--------	--------	--------	--------	--------

# **CHAPTER 6**

# **APPLICATION OF DERIVATIVES**



The sight of soap bubble produced using a bubble wand is very exciting! One application of derivative is finding the rate of increase of size of the bubble (dv/dt) due to increasing radius, where *V* is the volume of spherical bubble and *r* is the radius. This can be calculated by knowing the rate of increase of radius with time (dr/dt).

# **APPLICATION OF DERIVATIVES**

Topics to be covered as per C.B.S.E. revised syllabus (2024-25)

- Applications of derivatives:
- rate of change of quantities,
- increasing/decreasing functions,
- maxima and minima (first derivative test motivated geometrically and second derivative test given as a provable tool).
- Simple problems (that illustrate basic principles and understanding of the subject as well as real life situations).

## POINTS TO REMEMBER

• Rate of change: Let y = f(x) be a function then the rate of change of y with respect to x is given by  $\frac{dy}{dx} = f'(x)$  where a quantity y varies with another quantity x.

 $\left\{\frac{dy}{dx}\right\}_{x=x_1}$  or  $f'(x_1)$  represents the rate of change of y w.r.t. x at  $x = x_1$ .

#### Increasing and Decreasing Function

Let f be a real-valued function and let I be any interval in the domain of f. Then f is said to be

a) Strictly increasing on I, if for all  $x_1, x_2 \in I$ , we have

$$x_1 < x_2 \Rightarrow f(x_1) < f(x_2)$$

b) Increasing on I, if for all  $x_{1}$ ,  $x_2 \in I$ , we have

$$x_1 < x_2 \Rightarrow f(x_1) \le f(x_2)$$

c) Strictly decreasing in I, if for all  $x_{1, x_2} \in I$ , we have

$$x_1 < x_2 \Rightarrow f(x_1) > f(x_2)$$

d) Decreasing on I, if for all  $x_1, x_2 \in I$ , we have

$$x_1 < x_2 \Rightarrow f(x_1) \ge f(x_2)$$

- **Derivative Test:** Let f be a continuous function on [a, b] and differentiable on (a, b). Then
  - a) *f* is strictly increasing on [a, b] if f'(x) > 0 for each  $x \in (a, b)$ .
  - b) f is increasing on [a, b] if  $f'(x) \ge 0$  for each  $x \in (a, b)$ .
  - c) *f* is strictly decreasing on [a, b] if f'(x) < 0 for each  $x \in (a, b)$ .

- d) f is decreasing on [a, b] if  $f'(x) \le 0$  for each  $x \in (a, b)$ .
- e) *f* is constant function on [a, b] if f'(x) = 0 for each  $x \in (a, b)$ .

#### Maxima and Minima

- a) Let f be a function and c be a point in the domain of f such that either f '(x)=0 or f '(x) does not exist are called critical points.
- b) First Derivative Test: Let *f* be a function defined on an open intervalI. Let *f* be continuous at a critical point *c* in interval I.
  - i. f'(x) changes sign from positive to negative as x increases through *c*, then *c* is called the point of the local maxima.
  - ii. f'(x) changes sign from negative to positive as x increases through c, then c is a point of *local minima*.
  - iii. f '(x) does not change sign as x increases through c, then c is neither a point of *local maxima* nor a point of *local minima*.
     Such a point is called a point of *inflexion*.
- c) Second Derivative Test : Let f be a function defined on an interval I and let  $c \in I$ . Let f be twice differentiable at c. Then
  - i. x = c is a point of local maxima if f'(c) = 0 and f''(c) < 0. The value f(c) is local maximum value of f.
  - ii. x = c is a point of local minima if f'(c) = 0 and f''(c) > 0. The value f(c) is local minimum value of f.
  - iii. The test fails if f'(c) = 0 and f''(c) = 0.

### **EXTREME VALUE OF A FUNCTION**

Let y = f(x) be a real function defined on an interval I and C be any point in I. Then f is said to have an extreme value in / if f(c) is either maximum or minimum value of f in /.

Here, f(c) is called the extreme value and C is called one of the extreme points.

#### Illustration:

Let  $f(x) = (2x-1)^2 + 3$ . Then,  $f(x) \ge 3$ , as  $(2x-1)^2 \ge 0$ For any real number 'x'  $\Rightarrow (2x-1)^2 + 3 \ge 0 + 3$ Thus, minimum value of f(x) is 3, which occurs at  $x = \frac{1}{2}$ Also f(x) has no maximum value as  $f(x) \to \infty$  as  $|x| \to \infty$ 

#### Illustration:

Let  $g(x) = -(x-1)^2 + 10$ . Then,  $g(x) = 10 - (x-1)^2 \le 10 \quad \forall x \in R \text{ as } (x-1)^2$  is Always greater them or equal to zero. Thus maximum value of g(x) is 10, which occurs at x = 1Also g(x) has no minimum value of  $f(x) \rightarrow -\infty$  as  $|x| \rightarrow \infty$ .

#### Illustration:

Neither maximum nor minimum value of a function.

Let as consider a function  $f(x) = x^3$ ,  $x \in (-1, 1)$ 

Since this function is an increasing function in (-1, 1), it should have minimum value at a point nearest to -1 and maximum value at a point nearest to 1.

But we can not locate such points (see figure)

So,  $f(x) = x^3$ , has neither maximum nor-minimum value in (-1, 1).



But, if we extend the domain of *f* to [-1, 1], then the function  $f(x) = x^3$  has maximum value 1 at x = 1 and minimum value -1 at x = -1

Note: Every continuous function on an closed interval has a maximum and minimum

# **ONE MARK QUESTIONS**

# Multiple Choice Questions(MCQ)

- 1. If a function  $f: R \rightarrow R$  is defined by  $f(x) = 2x + \cos x$ , then
  - (a) f has a minimum at  $x = \pi$
  - (b) has a maximum at x=0
  - (c) f is a decreasing function
  - (d) f is an increasing function
- 2. If the radius of circle is increasing at the rate of 2cm/sec, then the area of circle when its radius is 20 cm is increasing at the rate of
  - (a)  $80 \pi m^2 / sec$  (b)  $80 m^2 / sec$
  - (c)  $80 \pi cm^2 / sec$  (d)  $80 cm^2 / sec$

3.	The	maximum value of	log x is:			
	(a)	е	x	(b)	2e	
	(c)	$\frac{1}{e}$		(d)	$\frac{2}{e}$	

- 4. The interval on which the function  $f(x) = 2x^3 + 9x^2 + 12x 1$ is decreasing is :
  - (a)  $[-l, \infty)$  (b)  $(-\infty, -2]$ (c) [-2, -l) (d) [-l, 1)
- 5. The sides of an equilateral triangle are increasing at the rate of 2cm/sec. The rate at which its area increases, when its side is 10 cm is :
  - (a)  $10 \ cm^2 \ / \ sec$  (b)  $10 \ \sqrt{3} \ cm^2 \ / \ sec$
  - (c)  $\frac{10}{3}$  cm<sup>2</sup> / sec (d)  $\sqrt{3}$  cm<sup>2</sup> / sec

- 6. The function  $f(x) = x^x$ , x > o is increasing on the interval
  - (a) (0, e] (b) (0, 1/e)
  - (c)  $[1/e, \infty)$  (d) None of these
- 7. The function  $f(x) = 2x^3 15x^2 + 36x + 6$  is increasing in the interval:
  - (a)  $(-\infty, 2)U[3, \infty)$  (b)  $(-\infty, 2)$
  - (c)  $(-\infty, 2]U[3, \infty)$  (d)  $[3, \infty)$
- 8. A point on the curve  $y^2 = 18 \text{ x}$  at which ordinate increases twice the rate of abscissa is :
  - (a) (2, 4) (b) (2, -4)
  - (c)  $\left(\frac{-9}{8},\frac{9}{2}\right)$  (d)  $\left(\frac{9}{8},\frac{9}{2}\right)$

9. The least value of function  $f(x) = ax + \frac{b}{x}(x > 0, a > 0, b > 0)$  is:

- (a)  $\sqrt{ab}$  (b)  $2\sqrt{ab}$
- (c) ab (d) 2ab

10. At  $X = \frac{5\pi}{6}$ , the function f(x) = 2 sin 3x + 3cos 3x is

- (a) Maximum (b) Minimum
- (c) zero (d) Neither maximum nor minimum
- 11. The function tanx-x :
  - (a) always increases (b) always deccreases
  - (c) Remains contain (d) Sometime increases sometime decreases

12. The minimum value of  $x^2 + \frac{250}{x}$  is: (a) 75 (b) 55

- (c) 50 (d) 20
- 13. In a sphere of radius r, a right circular cone of height having maximum curved surface area is inscribed. The expression for the square of curved surface of the cone is:
  - (a)  $2\pi^2 rh(2rh+h^2)$  (b)  $\pi^2 hr(2rh+h^2)$
  - (c)  $2\pi^2 r (2rh^2 h^3)$  (d)  $2\pi^2 r^2 (2rh h^2)$

## **ASSERTION REASON TYPE QUESTIONS 1 Marks**

Statement I is called Assertion (A) Statements II is called Reason R. Read the given statiments carefully and chose the correct answer from the four options given below.

- (a) Both the statement are true and statement II is correct explantion of statement I
- (b) Both the statmetns are treu and statement II is not the correct explanation of statement I.
- (c) Statement I is true statement II is false
- (d) Statement I is false and statement II is true

14. Statement I. The function 
$$f(x) = x^x$$
,  $x > 0$ , is strictly increasing in  $\left(\frac{1}{R}, \infty\right)$ 

Statement II :  $\log_a x > b \Rightarrow x > a^b$  if a > 1

15. Let  $a, b \in R$  be such that the function f given by  $f(x) = \log |x| + bx^2 + an, x \neq 0$ has extreme values at x = -1, and x = 2Statement I : f has local maximum at x = -1 and x = 2

Statement II :  $a = \frac{1}{2}$  and  $b = \frac{-1}{4}$ 

16. Let  $f(x) = 2x^3 - 15x^3 + 36x + 1$ 

Statement I :: f is strictly decreasing in [2, 3] Statement I :: f is strictly increasing in  $(-\infty, 2] \cup [3, \infty)$ 

### **TWO MARKS QUESTIONS**

- 1. The sum of the two numbers is 8, what will be the maximum value of the sum of their reciprocals.
- 2. Find the maximum value of  $f(x) = 2x^3 24x + 107$  in the interval [1, 3]
- 3. If the rate of change of Area of a circle is equal to the rate of change its diameter. Find the radius of the circle.
- The sides of on equilateral triangle are increasing at the rate of 2 cm/s.
   Find the rate at which the area increases, when side is 10 cm.
- 5. If there is an error of a% in measuring the edge of cube, then what is the percentge error in its surface?
- 6. If an error of k% is made in measuring the radius of a sphere, then what is the percentage error in its volume?
- 7. If the curves  $y = 2e^x$  and  $y = ae^{-x}$  intersect orthogonally, then find a.
- 8. Find the point on the curve  $y^2 = 8x$  for which the abscissa and ordinate change at the same rate.
- 9. Prove that the function  $f(x) = \tan x 4x$  is strictly decreasing on  $\left[\frac{-\pi}{3}, \frac{\pi}{3}\right]$ .
- 10. Find the point on the curve  $y = x^2$ , where the slope of the tangent is equal to the x coordinate of the point.
- 11. Use differentials to approximate the cube root of 66.
- 12. Find the maximum and minimum values of the function  $f(x) = \sin(\sin x)$
- 13. Find the local maxima and minima of the function  $f(x) = 2x^3 21x^2 + 36x 20$ .
- 14. If  $y = a \log x + bx^2 + x$  has its exteme values at x = -1 and x = 2, then find a and b.
- 15. If the radius of the circle increases from 5 into 5.1 cm, then find the increase in area.

## THREE MARKS QUESTIONS

- 1. In a competition, a brave child tries to inflate a huge spherical balloon bearing slogans against child labour at the rate of 900  $cm^3$  of gas per second. Find the rate at which the radius of the balloon is increasing, when its radius is 15 cm.
- 2. An inverted cone has a depth of 10 cm and a base of radius 5 cm. Water is poured into it at the rate of  $\frac{3}{2}$  c.c. per minute. Find the rate at which the level of water in the cone is rising when the depth is 4 cm.
- 3. The volume of a cube is increasing at a constant rate. Prove that the increase in its surface area varies inversely as the length of an edge of the cube.
- 4. A kite is moving horizontally at a height of 151.5 meters. If the speed of the kite is 10m/sec, how fast is the string being let out when the kite is 250 m away from the boy who is flying the kite ? The height of the boy is 1.5 m.
- 5. A swimming pool is to be drained for cleaning. If L represents the number of litres of water in the pool t seconds after the pool has been plugged off to drain and  $L = 200(10 t)^2$ . How fast is the water running out at the end of 5 sec. and what is the average rate at which the water flows out during the first 5 seconds?
- 6. A man 2m tall, walk at a uniform speed of 6km/h away from a lamp post 6m high. Find the rate at which the length of his shadow increases.
- 7. A water tank has the shape of an inverted right circular cone with its axis vertical and vertex lower most. Its semi- vertical angle is  $\tan^{-1}(0.5)$ . water is poured into it at a constant rate of  $5m^3/h$ . Find the rate at which the level of the water is rising at the instant, when the depth of Water in the tank is 4m.

- 8. A spherical ball of salt is dissolving in water in such a manner that the rate of decrease of the volume at any instant is proportional to the surface area. Prove that the radius is decreasing at a constant rate.
- 9. A conical vessel whose height is 10 meters and the radius of whose base is half that of the height is being filled with a liquid at a uniform rate of  $1.5m^3/min$ . find the rate at which the level of the water in the vessel is rising when it is 3m below the top of the vessel.
- 10. Let *x* and *y* be the sides of two squares such that  $y = x x^2$ . Find the rate of change of area of the second square w.r.t. the area of the first square.
- 11. The length of a rectangle is increasing at the rate of 3.5 cm/sec. and its breadth is decreasing at the rate of 3 cm/sec. Find the rate of change of the area of the rectangle when length is 12 cm and breadth is 8 cm.
- 12. If the areas of a circle increases at a uniform rate, then prove that the perimeter various inversely as the radius.
- 13. Show that  $f(x) = x^3 6x^2 + 18x + 5$  is an increasing function for all  $x \in R$ . Find its value when the rate of increase of f(x) is least.

[Hint: Rate of increase is least when f'(x) is least.]

- 14. Determine whether the following function is increasing or decreasing in the given interval:  $f(x) = \cos\left(2x + \frac{\pi}{4}\right), \frac{3\pi}{8} \le x \le \frac{5\pi}{8}$ .
- 15. Determine for which values of x, the function  $y=x^4 \frac{4x^3}{3}$  is increasing and for which it is decreasing.
- 16. Find the interval of increasing and decreasing of the function  $f(x) = \frac{\log x}{x}$
- 17. Find the interval of increasing and decreasing of the function  $f(x) = \sin x \cos x$ ,  $0 < x < 2\pi$ .
- 18. Show that  $f(x) = x^2 e^{-x}$ ,  $0 \le x \le 2$  is increasing in the indicated interval.

- 19. Prove that the function  $y = \frac{4 \sin \theta}{2 + \cos \theta} \theta$  is an increasing function of  $\theta$  in  $\left[0, \frac{\pi}{2}\right]$ .
- 20. Find the intervals in which the following functionis decreasing.

$$f(x) = x^4 - 8x^3 + 22x^2 - 24x + 21$$

- 21. Find the interval in which the function  $f(x) = 5x^{\frac{3}{2}} 3x^{\frac{5}{2}}$ , x > 0 is strictly decreasing.
- 22. Show that the function  $f(x) = \tan^{-1}(\sin x + \cos x)$ , is strictly increasing the interval  $\left(0, \frac{\pi}{4}\right)$ .
- 23. Find the interval in which the function  $f(x) = \cos^{-1}\left(\frac{1-x^2}{1+x^2}\right)$  is increasing or decreasing.
- 24. Find the interval in which the function given by

$$f(x) = \frac{3x^4}{10} - \frac{4x^3}{5} - 3x^2 + \frac{36x}{5} + 11$$

- (i) strictly increasing
- (ii) strictly decreasing
- 25. Show that the curves  $xy = a^2$  and  $x^2 + y^2 = 2a^2$  touch each other.
- 26. For the curve  $y = 5x 2x^{3}$ , if x increases at the rate of 2 Units/sec. then how fast is the slope of the curve changing when x=3?
- 27. If the radius of a circle increases from 5 cm to 5.1 cm, find the increase in area.
- 28. If the side of a cube be increased by 0.1%, find the corresponding increase in the volume of the cube.

- 29. Find the maximum and minimum values of  $f(x) = \sin x + \frac{1}{2}\cos 2x$  in  $\left[0, \frac{\pi}{2}\right]$ .
- 30. Find the absolute maximum value and absolute minimum value of the following question  $f(x) = \left(\frac{1}{2} x\right)^2 + x^3$  in [-2, 2.5]
- 31. Find the maximum and minimum values of  $f(x) = x^{50} x^{20}$  in the interval [0, 1]
- 32. Find the absolute maximum and absolute minimum value of  $f(x) = (x 2)\sqrt{x 1}$  in [1, 9]
- 33. Find the difference between the greatest and least values of the function  $f(x) = \sin 2x x$  on  $\left[-\frac{\pi}{2}, \frac{\pi}{2}\right]$ .

#### **FIVE MARKS QUESTIONS**

- 1. Prove that the least perimeter of an isosceles triangle in which a circle of radius r can be inscribed is  $6\sqrt{3} r$ .
- 2. If the sum of length of hypotenuse and a side of a right angled triangle is given, show that area of triangle is maximum, when the angle between them is  $\frac{\pi}{3}$ .
- 3. Show that semi-vertical angle of a cone of maximum volume and given slant height is  $\cos^{-1}\left(\frac{1}{\sqrt{3}}\right)$ .
- 4. The sum of the surface areas of cuboids with sides x, 2x and  $\frac{x}{3}$  and a sphere is given to be constant. Prove that the sum of their volumes is minimum if x = 3 radius of the sphere. Also find the minimum value of the sum of their volumes.
- 5. Show that the volume of the largest cone that can be inscribed in a sphere of radius R is  $\frac{8}{27}$  of the volume of the sphere.
- 6. Show that the cone of the greatest volume which can be inscribed in a given sphere has an altitude equal to  $\frac{2}{3}$  of the diameter of the sphere.

- 7. Prove that the radius of the right circular cylinder of greatest curved surface area which can be inscribed in a given cone is half of that of the cone.
- 8. Show that the volume of the greatest cylinder which can be inscribed in a cone of height h and semi-vertical angle  $\alpha$  is  $\frac{4}{27}\pi h^3 tan^2 \alpha$ . Also show that height of the cylinder is  $\frac{h}{2}$
- 9. Find the point on the curve  $y^2 = 4x$  which is nearest to the point (2,1).
- 10. Find the shortest distance between the line y x = 1 and the curve $x = y^2$ .
- 11. A wire of length 36 m is to be cut into two pieces. One of the pieces is to be made into a square and the other into a circle. What should be the length of the two pieces, so that the combined area of the square and the circle is minimum?
- 12. Show that the height of the cylinder of maximum volume that can be inscribed in a sphere of radius r is  $\frac{2r}{\sqrt{3}}$ .
- 13. Find the area of greatest rectangle that can be inscribed in an ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1.$

## SELF ASSESSMENT-1

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. For the curve  $y = 5x 2x^3$ , if x increases at the rate of 2 units/sec, then how fast is the slope of curve changing when x = 3
  - (a) 72 units/sec (b) -72 units/sec
  - (c) 54 units/sec (d) -54 units/sec
- 2. The function  $f(x) = \tan x 4x$ , on  $\left(-\frac{\pi}{3}, \frac{\pi}{3}\right)$  is
  - (a) strictly decreasing (b) strictly increasing
  - (c) neither increasing nor deceasing
  - (d) None of these

3. The curve  $y = xe^x$  has minimum value equal to

(a) 1 (b) 0  
(c) 
$$-e$$
 (d)  $/\frac{1}{e}$ 

- 4. The sides of an equilateral triangle are increasing at the rate of 2 cm/sec. The rate (in cm<sup>2</sup>/sec) at which the area increases, when side is 10 cm is
  - (a) 10 (b) 5
  - (c)  $10\sqrt{3}$  (d)  $5\sqrt{3}$
- 5. If ab = 2a + 3b, a > 0, b > 0 then the minimum value of ab is
  - (a) 6 (b) 12 (c) 24 (d) 48

#### **SELF ASSESSMENT-2**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. If the function  $f(x) = 2x^3 9ax^2 + 12a^2x + 1$  where a > 0, attains its maximum and minimum at *p* and *q* respectively such that  $p^2 = q$ , then a =
  - (a) 0 (b) 1
  - (c) 2 (d) 3
- 2. The interval in which  $y = -x^3 + 3x^2 + 2022$  is increasing is
  - (a)  $(-\infty, 0) \cup (2, \infty)$  (b)  $(2, \infty)$
  - (c) (0,2) (d)  $(-\infty,0)$
- 3. The maximum value of the function  $f(x) = 4\sin x \cdot \cos x$  is
  - (a) 1 (b) 2
  - (c) 3 (d) 4

4. Which of the following function is decreasing on  $\left(0, \frac{\pi}{2}\right)$ 

- (a)  $\cos x$  (b)  $\sin x$
- (c)  $\tan x$  (d)  $\sin 2x$
- 5. A man of height 2 metres walks at a uniform speed of 5 km/h away from a lamp post which is 6 metres high. The rate at which the length of his shadow increases is
  - (a) 5 km/hr (b) 2 km/hr
  - (c) 3 km/hr (d) 2.5 km/hr

# <u>Answers</u>

ONE MARK QUESTIONS	IWO		
Answer	1. $\frac{1}{2}$		
1. (d) f is an increasing	2. 89		
2. (c) $80\pi$ cm <sup>2</sup> / sec	3. $\frac{1}{\pi}$		
3. (c) <u>1</u>	4.		
e 4. (c) [-21]	5.		
5. (b) $10\sqrt{3}$ cm <sup>2</sup> / sec	6. 7.		
(-)	8.		
6. (C) [1/e, ∞)	10.		
7. (c) (–∞, 2] U [3, ∞)	11.		
$\left(\frac{9}{2},\frac{7}{2}\right)$	12.		
o. (u) (8'2)	13.		
9. (b) 2√ab	1.1		
10. (d) Neither maximum	14.		
nor minimum			
11 (a) always increases	INKEE		
12. (a) 75	1.		
13.(c) $2\pi^2 r (2rh^2 - h^3)$	2.		
14 (a)	4.		
14.(d)	5.		
15. (a)	б.		
16. (b)	7.		
× /	0		

TW	O MARKS QUESTIONS
1.	$\frac{1}{2}$
2.	89
3.	$\frac{1}{\pi}$ units
4.	$10\sqrt{3} \ cm^2$ / s
5.	2a%
6.	3k %
7.	1/2
8.	(2, 4)
10.	(0, 0)
11.	4.042
12.	sin 1, – sin 1
13.	Local maxima at x = 1 Local minima at x = 6
14.	a = 2, b = -1/ <sub>2</sub>
15.	$\pi$ cm <sup>2</sup>
THRE	E MARKS QUESTIONS
1	$\frac{1}{\pi}$ cm / s
2	$\frac{3}{8\pi}$ cm / min
4	. 8 m/sec.

5. 3000 L/s 6. 3 km/h 7.  $\frac{35}{88}$  m/h 9.  $\frac{6}{49\pi}$  m/min.

- 10.  $1 3x + 2x^2$ 11.  $8 \frac{\text{cm}^2}{\text{sec}}$ 13. 25
- 14. Increasing
- 15. Increasing for all  $x \ge 1$ Decreasing for all  $x \le 1$
- 16. Increasing on (o, e)

Decreasing on [e,  $\infty$ )

17. Increasing on

$$\left(\mathbf{O},\frac{3\pi}{4}\right)\cup\left(\frac{7\pi}{4},2\pi\right)$$

Decreasing on  $\left[\frac{3\pi}{4}, \frac{7\pi}{4}\right]$ 

- 20.  $(-\infty, 1]$ U[2, 3]
- 21. [1,∞]
- 23. increasing on  $[0, \infty)$ Decreasing  $(-\infty, 0]$
- 24. (i) Strictly increasing [-2,1] U [3, ∞)
  - (ii) Strictly decreasing
     (-∞, -2] U [1, 3]

- 26. decrease 72 units/sec.
- 27.  $\pi cm^2$
- 28. 0.3%
- 29. max. value  $=\frac{3}{4}$ , mim value  $=\frac{1}{2}$
- 30. ab. Max.= $\frac{157}{8}$ , ab. Min. = $\frac{-7}{4}$
- 31. max.value=0,

min.value =  $\frac{-3}{5} \left[\frac{2}{5}\right]^{2/3}$ 

- 32. ab. Max = 14 at x = 9 ab. Min.= $\frac{-3}{4^{4/3}}$  at  $x = \frac{5}{4}$
- 33. *π*

### **FIVE MARKS QUESTIONS**

4.  $18r^{3} + \frac{4}{3}\pi r^{3}$ 9. (1, 2) 10.  $\frac{3\sqrt{2}}{8}$ 11.  $\frac{144}{\pi+4}m, \frac{36\pi}{\pi+4}m$ 13. 2ab sq. Units.

## **SELF ASSESSMENT TEST-1**

1.	(b)	2. (a)	3. (d)	4. (c)	5. (c)
		SELF AS	SESSMENT	TEST-2	
1.	(c)	2. (c)	3. (b)	4. (a)	5. (d)

# CHAPTER 7

# INTEGRALS



There are many applications of integration in the field such as Physics, Engineering, Business, Economics etc. One of the important application of integration is finding the profit function of producing a certain number of cars if the marginal cost and revenue function are known. Companies can thus determine the maximum profit that can be earned and in this way plan their production, labour and other infrastructure accordingly.

# INTEGRALS

Topics to be covered as per C.B.S.E. revised syllabus (2024-25)

- Integration as inverse process of differentiation
- Integration of a variety of functions by substitution, by partial fractions and by parts
- Evaluation of simple integrals of the following types and problems based on them.

$$\int \frac{dx}{x^2 \pm a^2}, \int \frac{dx}{\sqrt{x^2 \pm a^2}}, \int \frac{dx}{\sqrt{a^2 - x^2}}, \int \frac{dx}{ax^2 + bx + c}, \int \frac{dx}{\sqrt{ax^2 + bx + c}}$$
$$\int \frac{px + q}{ax^2 \pm bx + c} dx, \int \frac{px + q}{\sqrt{ax^2 + bx + c}} dx, \int \sqrt{a^2 \pm x^2} dx, \int \sqrt{x^2 - a^2} dx$$
$$\int \sqrt{ax^2 + bx + c} dx$$

Fundamental Theorem of Calculus (without proof).

## POINTS TO REMEMBER

- Integration or anti derivative is the reverse process of Differentiation.
- Let  $\frac{d}{dx}F(x) = f(x)$  then we write  $\int f(x) dx = F(x) + c$ .
- These integrals are called indefinite integrals and c is called constant of integration.
- From geometrical point of view, an indefinite integral is the collection of family of curves each of which is obtained by translating one of the curves parallel to itself upwards or downwards along y-axis.

## STANDARD FORMULAE

1. 
$$\int x^{n} dx = \begin{cases} \frac{x^{n+1}}{n+1} + c, & n \neq -1 \\ \log_{e} |x| + c, & n = -1 \end{cases}$$

2. 
$$\int (ax+b)^n dx = \begin{cases} \frac{(ax+b)^{n+1}}{(n+1)a} + c, & n \neq -1 \\ \frac{1}{a} \log|ax+b| + c, & n = -1 \end{cases}$$

- 3.  $\int \sin x \, dx = -\cos x + c.$
- 4.  $\int \cos x \, dx = \sin x + c$
- 5.  $\int \tan x \, dx = -\log|\cos x| + c = \log|\sec x| + c.$
- 6.  $\int \cot x \, dx = \log |\sin x| + c.$
- 7.  $\int \sec^2 x \, dx = \tan x + c$
- 8.  $\int \csc^2 x \, dx = -\cot x + c$
- 9.  $\int \sec x \, \tan x \, dx = \sec x + c$
- 10.  $\int \csc x \cot x \, dx = -\csc x + c$
11.  $\int \sec x \, dx = \log |\sec x + \tan x| + c$ 

$$= \log \left| \tan \left( \frac{x}{2} + \frac{\pi}{4} \right) \right| + c$$

12. 
$$\int \csc x \, dx = \log |\csc x - \cot x| + c$$

$$= \log \left| \tan \frac{x}{2} \right| + c$$

$$13. \qquad \int e^x \, dx = e^x + c$$

14. 
$$\int a^x \, dx = \frac{a^x}{\log a} + c$$

15. 
$$\int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + c$$
,  $|x| < 1$ 

$$= -\cos^{-1}x + c$$

16. 
$$\int \frac{1}{1+x^2} dx = \tan^{-1} x + c$$
$$= -\cot^{-1} x + c$$

17. 
$$\int \frac{1}{|x|\sqrt{x^2-1}} \, dx = \sec^{-1} x + c, |x| > 1$$
$$= -\csc^{-1} x + c$$

- 18.  $\int \frac{1}{a^2 x^2} dx = \frac{1}{2a} \log \left| \frac{a + x}{a x} \right| + c$
- 19.  $\int \frac{1}{x^2 a^2} dx = \frac{1}{2a} \log \left| \frac{x a}{x + a} \right| + c$
- 20.  $\int \frac{1}{a^2 + x^2} dx = \frac{1}{a} \tan^{-1} \frac{x}{a} + c$
- 21.  $\int \frac{1}{\sqrt{a^2 x^2}} dx = \sin^{-1} \frac{x}{a} + c = -\cos^{-1} \frac{x}{a} + c$

22. 
$$\int \frac{1}{\sqrt{a^2 + x^2}} dx = \log |x + \sqrt{a^2 + x^2}| + c$$

23. 
$$\int \frac{1}{\sqrt{x^2 - a^2}} dx = \log |x + \sqrt{x^2 - a^2}| + c$$

24. 
$$\int \sqrt{a^2 - x^2} \, dx = \frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} + c$$

25. 
$$\int \sqrt{a^2 + x^2} \, dx = \frac{x}{2} \sqrt{a^2 + x^2} + \frac{a^2}{2} \log \left| x + \sqrt{a^2 + x^2} \right| + c$$

26. 
$$\int \sqrt{x^2 - a^2} \, dx = \frac{x}{2} \sqrt{x^2 - a^2} - \frac{a^2}{2} \log \left| x + \sqrt{x^2 - a^2} \right| + c$$

# **RULES OF INTEGRATION**

- 1.  $\int [(f_1(x) \pm f_2(x) \pm \dots \pm f_x(x)] dx = \int f_1(x) dx \pm \int f_2(x) dx \pm \dots + \int f_x(x) dx = \int f_1(x) dx = \int f_2(x) dx + \dots + \int f_x(x) dx$
- 2.  $\int k. f(x) dx = k \int f(x) dx$ .

3. 
$$\int e^{x} \{f(x) + f'(x)\} dx = e^{x} f(x) + c$$

# INTEGRATION BY SUBSTITUTION

1. 
$$\int \frac{f'(x)}{f(x)} dx = \log|f(x)| + c$$

2. 
$$\int [f(x)]^n f'(x) dx = \frac{[f(x)]^{n+1}}{n+1} + c$$

3. 
$$\int \frac{f'(x)}{[f(x)]^n} dx = \frac{(f(x))^{-n+1}}{-n+1} + c$$

# **INTEGRATION BY PARTS**

$$\int f(x) g(x) dx = f(x) \int g(x) dx - \int \left[ f(x) \int g(x) dx \right]$$

### **DEFINITE INTEGRALS**

$$\int_{a}^{b} f(x)dx = F(b) - F(a), \text{ where } F(x) = \int f(x) dx$$

## DEFINITE INTEGRAL AS A LIMIT OF SUMS.

$$\int_{a}^{b} f(x)dx = \lim_{h \to o} h \left[ f(a) + f(a+h) + f(a+2h) + \dots + f(a+n-1h) \right]$$
  
Where  $h = \frac{b-a}{h}$  or  $\int_{a}^{b} f(x)dx = \lim_{h \to o} \left[ h \sum_{r=1}^{n} f(a+rh) \right]$ 

# **PROPERTIES OF DEFINITE INTEGRAL**

1. 
$$\int_{a}^{b} f(x) = -\int_{b}^{a} f(x) dx$$
  
2. 
$$\int_{a}^{b} f(x) dx = \int_{a}^{b} f(t) dt.$$
  
3. 
$$\int_{a}^{b} f(x) dx = \int_{a}^{c} f(x) dx + \int_{c}^{b} f(x) dx.$$
  
4. 
$$(i) \int_{a}^{b} f(x) dx = \int_{a}^{b} f(a + b - x) dx.$$
  
(ii) 
$$\int_{0}^{a} f(x) dx = \int_{0}^{a} f(a - x) dx$$
  
5. 
$$\int_{-a}^{a} f(x) dx = 2 \int_{0}^{a} f(x) dx, \quad if f(x) is even function$$
  
6. 
$$\int_{-a}^{a} f(x) dx = 0 if f(x) is an odd function$$
  
7. 
$$\int_{0}^{2a} f(x) dx = \begin{cases} 2 \int_{0}^{a} f(x) dx, & \text{if } f(2a - x) = f(x) \\ 0 & \text{if } f(2a - x) = -f(x) \end{cases}$$

Illustration:  
Evaluate 
$$\int e^x \left(\frac{x-2}{x+4}\right)^2 dx$$
  
Solution:  $I = \int e^x \left(\frac{x-2}{x+4}\right)^2 dx = \int e^x \left(1 - \frac{2}{x+4}\right)^2 dx$   
 $= \int e^x \left[\left(1 - \frac{4}{x+4}\right) + \frac{4}{(x+4)^2}\right] dx$  It is of the form  $e^x \left[f(x) + f'(x)\right] dx$   
 $= \int e^x [f(x) + f'(x)] dx$ , where  $f(x) = 1 - \frac{4}{x+4}$   
 $= e^x f(x) + C = e^x \left(1 - \frac{4}{x+4}\right) + C = \frac{xe^x}{x+4} + C$ 

Illustration:  
Find 
$$\int \frac{x^2 + 1}{(x+1)^2} dx$$
  
Solution:  $\int \frac{x^2 + 1}{(x+1)^2} dx = \int \frac{(x+1)^2 - 2x}{(x+1)^2} dx$   
 $= \int \frac{(x+1)^2 - 2(x+1) + 2}{(x+1)^2} dx$   
 $= \int \left[ 1 - \frac{2}{x+1} + \frac{2}{(x+1)^2} \right] dx$   
 $= x - 2 \log |x+1| - \frac{2}{x+1} + C$ 

Illustration:  
Evaluate 
$$\int_{0}^{\pi/4} \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx$$
Solution: 
$$\int_{0}^{\pi/4} \frac{\sin^2 x \cos^2 x}{(\sin^3 x + \cos^3 x)^2} dx = \int_{0}^{\pi/4} \frac{\tan^2 x \sec^2 x}{(\tan^3 x + 1)^2} dx$$
[dividing Num and Den by  $\cos^6 x$ ]  
Put  $z = \tan^3 x + 1$ ,  
then  $dz = 3\tan^2 x \sec^2 x \, dx$ 

Also when 
$$x = 0, z = 0$$
 and when  $x = \frac{\pi}{4}, z = 2$   
Now  $I = \frac{1}{3}\int_{2}^{1}\frac{dz}{z^2} = -\frac{1}{3}\left[\frac{1}{z}\right]_{1}^{2} = -\frac{1}{3}\left[\frac{1}{2}-1\right] = \frac{1}{6}$ 

Illustration:  $\operatorname{Find}_{-\pi/4}^{\pi/4} \frac{x + \pi/4}{2 - \cos 2x} dx$ Solution:  $\int_{-\pi/4}^{\pi/4} \frac{x + \frac{\pi}{4}}{2 - \cos 2x} dx = \int_{-\pi/4}^{\pi/4} \frac{x}{2 - \cos 2x} dx + \frac{\pi}{4} \int_{-\pi/4}^{\pi/4} \frac{1}{2 - \cos 2x} dx$  $= 0 + \frac{\pi}{4} \cdot 2 \int_{0}^{\pi/4} \frac{dx}{2 - \cos x}$  [Since first function is an even function and second function is an odd function]  $=\frac{\pi}{2}\int_{0}^{\pi/4}\frac{dx}{2(1-2\sin^{2}x)}$  $=\frac{\pi}{2}\int_{0}^{\pi/4}\frac{dx}{2\sin^2 x+1}$  $=\frac{\pi}{2}\int_{0}^{\pi/4}\frac{\sec^2 x}{3\tan^2 x+1}dx$  [dividing num and den by  $\cos^2 x$ ] Put  $z = \sqrt{3} \tan x$ , then  $dz = \sqrt{3} \sec^2 x \, dx$ Also when x = 0, z = 0, and when  $x = \frac{\pi}{4}$ ,  $z = \sqrt{3}$ :. From (i),  $I = \frac{\pi}{2} \frac{1}{\sqrt{3}} \int_{2}^{\sqrt{3}} \frac{dz}{z^2 + 1} = \frac{\pi}{2\sqrt{3}} \left[ \tan^{1/2} z \right]_{0}^{\sqrt{3}}$  $=\frac{\pi}{2\sqrt{3}}\left[\tan^{-1}\sqrt{3}-\tan^{-1}0\right]$  $=\frac{\pi}{2\sqrt{3}}\tan^{-1}\sqrt{3}$  $=\frac{\pi}{2\sqrt{3}}\cdot\frac{\pi}{3}=\frac{\pi^2}{6\sqrt{3}}$ 

#### **ONE MARK QUESTIONS**

Evaluate the following integrals:

1. Integrate  $\int_{0}^{2} (x^{2} + x + 1) dx$ (a)  $\frac{15}{2}$ (b) 20/5 (c) 20/3 (d) 3/20 2.  $\int_{0}^{\pi} \sin^{2} x \, dx =$ (a)  $\frac{\pi}{2}$ (b)  $\frac{\pi}{4}$ (c) 2π (d) 4π 3.  $\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx$  equal to: (a)  $-\frac{1}{\sin x + \cos x} + c$  (b)  $\log |\sin x + \cos x| + c$ (c)  $\frac{1}{(\sin x + \cos x)^2}$  (d)  $\log |\sin x - \cos x| + c$ 4.  $\int \frac{(1+\log x)^2}{1+x^2} dx$  is : (a)  $\frac{1}{3}(1+\log x)^3 + c$  (b)  $\frac{1}{2}(1+\log x)^2 + c$ (c)  $\log(\log 1 + x) + c^2$  (d) None of these 5.  $\int \frac{\sin^2 x - \cos^2 x}{\sin^2 x \cos^2 x} dx$  is equal to (a)  $\tan x + \cos x + c$ (b)  $\tan x + \operatorname{cosec} x + c$ (c)  $\tan x + \cot x + c$ (d)  $\tan x + \sec x + c$ 

6. The value of 
$$\int_{-6}^{\frac{\pi}{3}} \frac{dx}{\sin 2x}$$
 is :  
(a)  $\frac{1}{2} \log(-1)$  (b)  $\log(-1)$   
(c)  $\log 3$  (d)  $\log \sqrt{3}$   
7. The value of  $\int_{0}^{1} \tan^{-1} \left(\frac{2x-1}{1+x-x^{2}}\right) dx$  is:  
(a) 1 (b) 0  
(c) -1 (d)  $\frac{\pi}{4}$   
8.  $\int \frac{x^{9}}{(4x^{2}+1)^{6}} dx$  is equal to  
(a)  $\frac{1}{5x} \left(4 + \frac{1}{x^{2}}\right)^{-5} + c$  (b)  $\frac{1}{5} \left(4 + \frac{1}{x^{2}}\right)^{-5} + c$   
(c)  $\frac{1}{10x} \left(\frac{1+4}{x^{2}}\right)^{-5} + c$  (d)  $\frac{1}{10} \left(\frac{1}{x^{2}} + 4\right)^{-5} + c$   
9. If  $\int \frac{x^{3}}{\sqrt{1+x^{2}}} dx = 9(1+x^{2})^{3/2} + b\sqrt{1+x^{c}} + c$   
(a)  $a = \frac{1}{3}, b = 1$  (b)  $a = -\frac{1}{3}, b = -1$   
(c)  $a = -\frac{1}{3}, b = -1$  (d)  $a = \frac{1}{3}, b = -1$ 

С

#### ASSERTION-REASON BASED QUESTIONS

In the following questions, a statement of ssertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both A and R are true and R is the correct explanation of A
- (b) Both A and R are true but R is not the correct explanation of A
- (c) A is true but R is false
- (d) A is false but R is true.

1. Assertion (A): 
$$\int \frac{dx}{x^2 + 2x + 3} = \frac{1}{\sqrt{2}} \tan^{-1} \left( \frac{x+1}{\sqrt{2}} \right) + c$$

Reason (R): 
$$\int \frac{dx}{x^2 + a^2} = \frac{1}{a} \tan^{-1} \left( \frac{x}{a} \right) + c$$

2. Assertion (A) :  $\int e^x [\sin x - \cos x] dx = e^x \sin x + c$ 

Reason (R): 
$$\int e^{x} [f(x) + f'(x)] dx = e^{x} (f(x) + c)$$

3. Assertion (A) : 
$$\int_{-2}^{2} \log\left(\frac{1+x}{1-x}\right) dx = 0$$

Reason (R) : 
$$\int_{0}^{2a} f(x) dx = 0$$
 if  $f(2a - x)$ 

4. Assertion (A) :  $\int_{\pi/6}^{\pi/3} \frac{1}{1 + (\tan x)^{11/5}} dx = \frac{\pi}{12}$ 

Reason (R) :  $\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$ 

# **TWO MARKS QUESTIONS**

#### Evaluate :

1.	$\int e^{[\log(x+1) - \log x]} dx$
2.	$\int \frac{1}{\sqrt{x+1} + \sqrt{x+2}} dx$
3.	$\int \sin x \sin 2x dx$
4.	$\int \left[\frac{x}{a} + \frac{a}{x} + x^a + a^x\right] dx$
5.	$\int_0^{\pi/2} \log\left(\frac{5+3\cos x}{5+3\sin x}\right) dx$
6.	$\int \frac{a^x + b^x}{c^x} dx$
7.	$\int \left(\sqrt{ax} - \frac{1}{\sqrt{ax}}\right)^2 dx$
8.	$\int e^x 2^x dx$
9.	$\int 2^{2^{2^{x}}} 2^{2^{x}} 2^{x} dx$

$$10. \qquad \int \frac{\sin\left(2\tan^{-1}x\right)}{1+x^2} dx$$

11.	∫xlog2x dx
12.	$\int_0^{\pi/4} \sqrt{1 + \sin 2x}  \mathrm{d}x$
13.	$\int_0^{\pi/2} e^x (\sin x - \cos x) dx$
14.	$\int_{4}^{9} \frac{\sqrt{x}}{(30 - x^{3/2})} dx$
15.	$\int_0^1 \frac{dx}{e^x + e^{-x}} dx$
16.	∫ <sup>log sinx </sup> tanx
17.	$\int \frac{\sin^4 x + \cos^4 x}{\sin^3 x + \cos^3 x}  dx$
18.	$\int \sqrt{\tan x} (1 + \tan^2 x) dx$
19.	$\int \frac{\sin 2x}{(a+b\cos x)^2} dx$
20.	$\int \frac{x^2 - x + 2}{x^2 + 1} dx$

## **THREE MARKS QUESTIONS**

#### Evaluate :

1. (i) 
$$\int \frac{x \operatorname{cosec} (\tan^{-1} x^2)}{1+x^4} dx$$
  
(ii) 
$$\int \frac{\sqrt{x+1} - \sqrt{x-1}}{\sqrt{x+1} + \sqrt{x-1}} dx$$
  
(iii) 
$$\int \frac{1}{\sin(x-a)\sin(x-b)} dx$$
  
(iv) 
$$\int \frac{\cos(x+a)}{\cos(x-a)} dx$$
  
(v) 
$$\int \cos 2x \cos 4x \cos 6x dx$$
  
(vi) 
$$\int \tan 2x \tan 3x \tan 5x dx$$
  
(vii) 
$$\int \sin^2 x \cos^4 x dx$$
  
(viii) 
$$\int \cot^3 x \csc^4 x dx$$
  
(ix) 
$$\int \frac{\sin x \cos x}{\sqrt{a^2 \sin^2 x + b^2 \cos^2 x}} dx \quad [\text{Hint: Put } a^2 \sin^2 x + b^2 \cos^2 x = t \text{ or } t^2$$
  
(x) 
$$\int \frac{1}{\sqrt{\cos^3 x \cos(x+a)}} dx$$
  
(xi) 
$$\int \frac{\sin^6 x + \cos^6 x}{\sin^2 x \cos^2 x} dx$$
  
(xii) 
$$\int \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$$

Evaluate :

2.

(i) 
$$\int \frac{x}{x^4 + x^2 + 1} dx$$
  
(ii) 
$$\int \frac{1}{x[6(\log x)^2 + 7\log x + 2]} dx$$
  
(iii) 
$$\int \frac{1}{\sqrt{\sin^3 x \cos^5 x}} dx$$
  
(iv) 
$$\int \frac{x^2 + 1}{x^4 + 1} dx$$
  
(iv) 
$$\int \frac{1}{\sqrt{(x-a)(x-b)}} dx$$
  
(v) 
$$\int \frac{5x-2}{3x^2 + 2x + 1} dx$$
  
(vi) 
$$\int \frac{x^2}{x^2 + 6x + 1} dx$$
  
(vii) 
$$\int \frac{x+2}{\sqrt{4x-x^2}} dx$$
  
(ix) 
$$\int x \sqrt{1 + x - x^2} dx$$
  
(x) 
$$\int \frac{\sin^4 x}{\cos^8 x} dx$$

(xi)  $\int \sqrt{\sec x - 1} \, dx$  [Hint: Multiply and divided by  $\sqrt{\sec x + 1}$ ]

Evaluate :

3. (i) 
$$\int \frac{\mathrm{d}x}{x(x^7+1)}$$

(ii) 
$$\int \frac{3x+5}{x^3-x^2-x+1} \, dx$$

(iii) 
$$\int \frac{\sin\theta\cos\theta}{\cos^2\theta - \cos\theta - 2} \, \mathrm{d}\theta$$

(iv) 
$$\int \frac{\mathrm{d}x}{(2-x)(x^2+3)}$$

(v) 
$$\int \frac{x^2 + x + 2}{(x-2)(x-1)} dx$$

(vi) 
$$\int \frac{(x^2+1)(x^2+2)}{(x^2+3)(x^2+4)} dx$$

(vii) 
$$\int \frac{\mathrm{dx}}{(2x+1)(x^2+4)}$$

(viii) 
$$\int \frac{x^2 - 1}{x^4 + x^2 + 1} dx$$

(ix)  $\int \sqrt{\tan x} \, dx$ 

(x) 
$$\int \frac{dx}{\sin x - \sin 2x}$$

### 4. Evaluate:

- (i)  $\int x^5 \sin x^3 dx$
- (ii)  $\int \sec^3 x \, dx$
- (iii)  $\int e^{ax} \cos(bx + c) dx$
- (iv)  $\int \sin^{-1}\left(\frac{6x}{1+9x^2}\right) dx$  [Hint: Put  $3x = \tan \theta$ ]
- (v)  $\int \cos \sqrt{x} \, dx$
- (vi)  $\int x^3 \tan^{-1} x \, dx$

$$\begin{array}{ll} \text{(vii)} & \int e^{2x} \left(\frac{1+\sin 2x}{1+\cos 2x}\right) \, dx \\ \text{(viii)} & \int \left[\frac{1}{\log x} - \frac{1}{(\log x)^2}\right] \, dx \\ \text{(ix)} & \int \sqrt{2ax - x^2} \, dx \\ \text{(ix)} & \int \sqrt{2ax - x^2} \, dx \\ \text{(x)} & \int e^x \frac{(x^2+1)}{(x+1)^2} \, dx \\ \text{(xi)} & \int x^3 \sin^{-1} \left(\frac{1}{x}\right) \, dx \\ \text{(xii)} & \int \left\{\log(\log x) + \frac{1}{(\log x)^2}\right\} \, dx \\ \text{(xiii)} & \int \left\{\log(\log x) + \frac{1}{(\log x)^2}\right\} \, dx \\ \text{(xiii)} & \int (6x + 5)\sqrt{6 + x - x^2} \, dx \\ \text{(xiv)} & \int \frac{1}{x^3 + 1} \, dx \\ \text{(xv)} & \int \tan^{-1} \left(\frac{x - 5}{1 + 5x}\right) \, dx \\ \text{(xvi)} & \int \frac{dx}{5 + 4 \cos x} \end{array}$$

- 5. Evaluate the following definite integrals:
  - (i)  $\int_0^{\pi/4} \frac{\sin x + \cos x}{9 + 16 \sin 2x} dx$
  - (ii)  $\int_0^{\pi/2} \cos 2x \, \log \sin x \, dx$

(iii) 
$$\int_0^1 x \sqrt{\frac{1-x^2}{1+x^2}} \, dx$$

(iv) 
$$\int_0^{1/\sqrt{2}} \frac{\sin^{-1}x}{(1-x^2)^{3/2}} dx$$

(v) 
$$\int_0^{\pi/2} \frac{\sin 2x}{\sin^4 x + \cos^4 x} \, dx$$

(vi) 
$$\int_0^1 \sin\left(2\tan^{-1}\sqrt{\frac{1+x}{1-x}}\right) dx$$

(vii) 
$$\int_0^{\pi/2} \frac{x + \sin x}{1 + \cos x} dx$$

(viii) 
$$\int_0^1 x \log\left(1 + \frac{x}{2}\right) dx$$

(ix) 
$$\int_{-1}^{1/2} |x \cos \pi x| \, dx$$

(x) 
$$\int_{-\pi}^{\pi} (\cos a x - \sin b x)^2 dx$$

### 6. Evaluate:

- (i)  $\int_{2}^{5} [|x-2| + |x-3| + |x-4|] dx$
- (ii)  $\int_0^{\pi} \frac{x}{1+\sin x} dx$
- (iii)  $\int_{-1}^{1} e^{\tan^{-1} x} \left[ \frac{1 + x + x^2}{1 + x^2} \right] dx$
- (iv)  $\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} \, dx$
- (v)  $\int_0^2 [x^2] dx$
- (vi)  $\int_0^{\pi/2} \frac{x \sin x \cos x}{\sin^4 x + \cos^4 x} \, dx$
- (vii)  $\int_0^{\pi} \frac{x}{a^2 \cos^2 x + b^2 \sin^2 x} dx [Hint: use \int_0^a f(x) dx = \int_0^a f(a x) dx$

7. Evaluate the following integrals:

(i) 
$$\int_{\pi/6}^{\pi/3} \frac{dx}{1+\sqrt{\tan x}}$$
  
(ii) 
$$\int_{-\pi/2}^{\pi/2} \left( \sin |x| + \cos |x| \right) dx$$
  
(iii) 
$$\int_{0}^{\pi} \frac{e^{\cos x}}{e^{\cos x} + e^{-\cos x}} dx$$
  
(iv) 
$$\int_{0}^{\pi} \frac{x \tan x}{\sec x + \csc x} dx$$

(v) 
$$\int_{-a}^{a} \sqrt{\frac{a-x}{a+x}} dx$$

### 8. Evaluate

(i) 
$$\int \frac{\sin^{-1} \sqrt{x} - \cos^{-1} \sqrt{x}}{\sin^{-1} \sqrt{x} + \cos^{-1} \sqrt{x}} \, \mathrm{dx} \ x \in [0, 1]$$

(*ii*) 
$$\int \sqrt{\frac{1-\sqrt{x}}{1+\sqrt{x}}} \, \mathrm{dx}$$

(iii) 
$$\int \frac{x^2 e^x}{(x+2)^2} dx$$

(iv) 
$$\int \frac{x^2}{(x \sin x + \cos x)^2} dx$$

(v) 
$$\int \sin^{-1} \sqrt{\frac{x}{a+x}} \, dx$$

$$(vi) \int_{\pi/6}^{\pi/3} \frac{\sin x + \cos x}{\sqrt{\sin 2x}} dx$$
$$(vii) \int \frac{\sin x}{\sin 4x} dx$$
$$(viii) \int_{-1}^{3/2} |x \sin \pi x| dx$$
$$(ix) \int \frac{\sin(x-a)}{\sin(x+a)} dx$$
$$(x) \int \frac{x^2}{(x^2+4)(x^2+9)} dx$$
$$(xi) \int \frac{\cos 5x + \cos 4x}{1 - 2\cos 3x} dx$$

### **FIVE MARKS QUESTIONS**

# 9. Evaluate the following integrals:

$$(i) \int \frac{x^{5} + 4}{x^{5} - x} dx$$
  
(ii)  $\int \frac{2e^{t}}{e^{3t} - 6e^{2t} + 11e^{t} - 6} dt$   
(iii)  $\int \frac{2x^{3}}{(x + 1)(x - 3)^{2}} dx$   
(iv)  $\int \frac{1 + \sin x}{\sin x (1 + \cos x)} dx$   
(v)  $\int_{0}^{\pi/2} (\sqrt{\tan x} + \sqrt{\cot x}) dx$ 

$$(\text{vi}) \int_{0}^{1} x \sqrt{\frac{1-x^2}{1+x^2}} dx$$
$$(\text{vii}) \int_{0}^{\pi/2} \frac{\cos x}{1+\cos x + \sin x} dx$$

10. Evaluate the following integrals as limit of sums:

(i) 
$$\int_{2}^{4} (2x + 1) dx$$
  
(ii)  $\int_{0}^{2} (x^{2} + 3) dx$   
(iii)  $\int_{1}^{3} (3x^{2} - 2x + 4) dx$   
(iv)  $\int_{0}^{4} (3x^{2} + e^{2x}) dx$   
(v)  $\int_{0}^{1} e^{2-3x} dx$   
(vi)  $\int_{0}^{1} (3x^{2} + 2x + 1) dx$ 

11. Evaluate:

(i) 
$$\int \frac{dx}{(\sin x - 2\cos x)(2\sin x + \cos x)}$$
  
(ii) 
$$\int_{0}^{1} \frac{\log(1+x)}{1+x^{2}} dx$$
  
(iii) 
$$\int_{0}^{\pi/2} (2\log\sin x - \log\sin 2x) dx$$

12. 
$$\int_{0}^{1} x(\tan^{-1} x)^{2} dx$$
  
13. 
$$\int_{0}^{\pi/2} \log \sin x dx$$

14. Prove that  $\int_{0}^{1} \tan^{-1} \left( \frac{1}{1 - x + x^2} \right) dx = 2 \int_{0}^{1} \tan^{-1} x \, dx$ 

Hence or otherwise evaluate the integral  $\int tan^{-1}(1-x+x^2) dx$ .

15. Evaluate 
$$\int_0^{\pi/2} \frac{\sin^2 x}{\sin x + \cos x} dx$$
.

#### **SELF ASSESSMENT-1**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. 
$$I = \int (x^8 + 1)(x^4 + 1)(x^2 + 1)(x + 1)(x - 1)dx =$$
  
(a)  $x^{16} - 1 + c$  (b)  $x^{17} - x + c$   
(c)  $\frac{x^{17}}{17} - x + c$  (d)  $\frac{x^{16}}{16} - x + c$   
2.  $\int \sin(x^2 + 2022) \cdot d(x^2) =$   
(a)  $2x \cdot \sin(x^2 + 2022) + c$  (b)  $-2x \cdot \cos(x^2 + 2022) + c$   
(c)  $\sin(x^2 + 2022) + c$  (d)  $-\cos(x^2 + 2022) + c$   
3.  $\int \cos^3 x \cdot \sqrt{\sin x} \, dx = \frac{2\sin^8 x}{3} - \frac{2\sin^8 x}{7} + c$ , then  $(a + b) =$   
(a)  $2$  (b)  $4$   
(c)  $5$  (d)  $6$   
4.  $\int \frac{\sin^2 x - \cos^2 x}{\sin^2 x \cdot \cos^2 x} \, dx =$   
(a)  $\tan x + \cot x + c$  (b)  $-\tan x + \cot x + c$   
(c)  $\tan x + \csc x + c$  (d)  $\tan x + \sec x + c$ 

5. 
$$\int_{0}^{\pi/2} \sin^2 x \, dx = \frac{\pi}{k}, \text{ then } k =$$
(a) 0.25
(b) 0.5
(c) 1
(

#### **SELF ASSESSMENT-2**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. 
$$\int_{0}^{\pi/2} \log \tan x \, dx =$$
(a) 0 (b) 1  
(c)  $\pi$  (d)  $\frac{\pi}{2}$   
2. 
$$\int_{0}^{\pi} \frac{x}{1+\sin x} \, dx =$$
(a)  $4\pi$  (b)  $\frac{\pi}{2}$   
(c)  $\pi$  (d)  $2\pi$   
3. 
$$\int \log(x^{2}+1) \, dx =$$
(a)  $x \log(x^{2}+1) - 2x + 2 \tan^{-1} x + c$  (b)  $x \log(x^{2}+1) - 2x - 2 \tan^{-1} x + c$   
(c)  $x \log(x^{2}+1) + 2x + 2 \tan^{-1} x + c$  (d) None of these  
4. 
$$\int e^{x} . \sin x \, dx =$$
(a)  $\frac{e^{x} (\sin x - \cos x)}{2} + c$  (b)  $\frac{e^{x} (\sin x - \cos x)}{2} - c$   
(c)  $\frac{e^{x} (-\sin x + \cos x)}{2} + c$  (d)  $\frac{-e^{x} (\sin x - \cos x)}{2} + c$   
5. 
$$\int \cos^{2} x \, dx = ax + b \sin 2x + c$$
, then  $(2a + 4b + 1) =$ 
(a) 0 (b) 1  
(c) 3 (d) -7

## Answers ONE MARKS QUESTIONS

1. (c) 
$$\frac{20}{3}$$
  
2. (a)  $\frac{\pi}{2}$   
3. (b)  $\log |\sin x + \cos x| + c$   
4. (a)  $\frac{1}{3}(1 + \log x)^3 + c$   
5. (c)  $\tan x + \cot x + c$   
6. (c)  $\log 3$   
7. (b) 0  
8.  $\frac{1}{10}(\frac{1}{x^2} + 4)^{-5} + c$ 

9. 
$$a = \frac{1}{3}, b = -1$$

#### INTEGRAL ASSERTION REASONS

- 1. A is true and R is correct explanation of A
- 2. Option (d) is correct
- 3. Option (b) is correct
- 4. (a) A is true and R is correct explanation of A

# **TWO MARKS QUESTIONS**

11.

12.

1.	x + log x + c
2.	$\frac{2}{3}\left[\left(x+2\right)^{3/2}-\left(x+1\right)^{3/2}\right]+c$
3.	$\frac{-1}{2}\left[\frac{\sin 3x}{3} - \sin x\right] + c$
4.	$\frac{1}{a}\frac{x^2}{2} + a\log x  + \frac{x^{a+1}}{a+1} + \frac{a^x}{\log a} + c$
5.	0
6.	$\frac{\left(\frac{a}{c}\right)^{x}}{\log\left \frac{a}{c}\right } + \frac{\left(\frac{b}{c}\right)^{x}}{\log\left \frac{b}{c}\right } + c$
7.	$\frac{ax^2}{2} + \frac{\log x }{a} - 2x + c$
8.	$\frac{2^{x}e^{x}}{\log(2e)}+c$
9.	$\frac{2^{2^{2^{x}}}}{(\log 2)^{3}} + C$
10.	$\frac{-\left[\cos\left(2\tan^{-1}x\right)\right]}{2}+C$

$$\frac{x^2}{2}\log 2x - \frac{x^2}{4} + C$$

13. 1  
14. 
$$\frac{19}{99}$$

1

15. 
$$\tan^{-1} e -\frac{\pi}{4}$$

$$\frac{\log|\sin x|^2}{2} - C$$

18. 
$$\frac{2}{3}(\tan x)^{3/2} + C$$

19. 
$$-\frac{2}{b^2} \left[ \log |a + b \cos x| + \frac{a}{a + b \cos x} \right] + C$$
  
20.  $x - \frac{1}{2} \log |x^2 + 1| + \tan^{-1} x + C$ 

## **THREE MARKS QUESTIONS**

1. (i) 
$$\frac{1}{2} \log \left[ \csc(\tan^{-1} x^2) - \frac{1}{x^2} \right] + c$$
  
(ii)  $\frac{1}{2} (x^2 - x\sqrt{x^2 - 1}) + \frac{1}{2} \log |x + \sqrt{x^2 - 1}| + c$   
(iii)  $\frac{1}{\sin(a-b)} \log \left| \frac{\sin(x-a)}{\sin(x-b)} \right| + c$   
(iv)  $x \cos 2a - \sin 2a \log |\sec(x-a)| + c$   
(v)  $\frac{3}{8}x - \frac{1}{4} \sin 2x + \frac{1}{32} \sin 4x + c$   
(vi)  $\frac{1}{5} \log |\sec 5x| - \frac{1}{2} \log |\sec 2x| - \frac{1}{3} \log |\sec 3x| + c$   
(vii)  $\frac{1}{32} \left[ 2x + \frac{1}{2} \sin 2x - \frac{1}{2} \sin 4x - \frac{1}{6} \sin 6x \right] + c$   
(viii)  $- \left( \frac{\cot^6 x}{6} + \frac{\cot^4 x}{4} \right) + c$   
(ix)  $\frac{1}{a^2 - b^2} \sqrt{a^2 \sin^2 x + b^2 \cos^2 x} + c$   
(x)  $-2 \csc a \sqrt{\cos a - \tan x \sin a} + c$   
(xi)  $\tan x - \cot x - 3x + c$   
(vi)  $\sin^{-1} [\sin x - \cos x] + c$   
2. (i)  $\frac{1}{\sqrt{3}} \tan^{-1} \left( \frac{2x^2 + 1}{\sqrt{3}} \right) + c$   
(ii)  $\log \left| \frac{2 \log x}{3 \log x} \right| + c$   
(iii)  $\log \left| \frac{2 \log x}{3 \log x} \right| + c$ 

[Class XII : Maths]

(iv) 
$$\frac{1}{\sqrt{2}} \tan^{-1} \left\{ \frac{1}{\sqrt{2}} \left( x - \frac{1}{x} \right) \right\} + c$$
  
(v)  $2 \log \left| \sqrt{x - a} + \sqrt{x - b} \right| + c$   
(vi)  $\frac{5}{6} \log \left| 3x^2 + 2x + 1 \right| + \frac{-11}{3\sqrt{2}} \tan^{-1} \left( \frac{3x + 1}{\sqrt{2}} \right) + c$   
(vii)  $x - 3 \log \left| x^2 + 6x + 12 \right| + 2\sqrt{3} \tan^{-1} \left( \frac{x + 3}{\sqrt{3}} \right) + c$   
(viii)  $-\sqrt{4x - x^2} + 4 \sin^{-1} \left( \frac{x - 2}{2} \right) + c$   
(ix)  $-\frac{1}{3} (1 + x - x^2)^{3/2} + \frac{1}{8} (2x - 1)\sqrt{1 + x - x^2} + \frac{5}{16} \sin^{-1} \left( \frac{2x - 1}{\sqrt{5}} \right) + c$   
(x)  $\frac{\tan^5 x}{5} + \frac{\tan^7 x}{7} + c$   
(xi)  $-\log \left| \cos x + \frac{1}{2} + \sqrt{\cos^2 x + \cos x} \right| + c$   
(i)  $\frac{1}{7} \log \left| \frac{x^7}{x^7 + 1} \right| + c$   
(ii)  $\frac{1}{2} \log \left| \frac{x + 1}{x - 1} \right| - \frac{4}{x - 1} + c$   
(iii)  $-\frac{2}{3} \log \left| \cos \theta - 2 \right| - \frac{1}{3} \log \left| 1 + \cos \theta \right| + c$ 

(iv) 
$$\frac{1}{14} \log \left| \frac{x^2 + 3}{(2 - x)^2} \right| + \frac{2}{7\sqrt{3}} \tan^{-1} \left( \frac{x}{\sqrt{3}} \right) + c$$

(v) 
$$x + 4 \log \left| \frac{(x-2)^2}{x-1} \right| + c$$

(vi) 
$$x + \frac{2}{\sqrt{3}} \tan^{-1}\left(\frac{x}{\sqrt{3}}\right) - 3 \tan^{-1}\left(\frac{x}{2}\right) + c$$

3.

(vii) 
$$\frac{2}{17}\log|2x+1| - \frac{1}{17}\log|x^2+4| + \frac{1}{34}\tan^{-1}\frac{x}{2} + c$$

(viii) 
$$\frac{1}{2} \log \left| \frac{x^2 - x + 1}{x^2 + x + 1} \right| + c$$

(ix) 
$$\frac{1}{\sqrt{2}} \tan^{-1} \left( \frac{\tan x - 1}{\sqrt{2 \tan x}} \right) + \frac{1}{2\sqrt{2}} \log \left| \frac{\tan x - \sqrt{2 \tan x} + 1}{\tan x + \sqrt{2 \tan x} + 1} \right| + c$$

(x) 
$$-\frac{1}{2}\log|\cos x - 1| - \frac{1}{6}\log|\cos x + 1| + \frac{2}{3}\log|1 - 2\cos x| + c$$

4. (i) 
$$\frac{1}{3}[-x^3\cos x^3 + \sin x^3] + c$$

(ii) 
$$\frac{1}{2}[\sec x \tan x + \log|\sec x + \tan x|] + c$$

(iii) 
$$\frac{e^{ax}}{a^2+b^2}[a\cos(bx+c)+b\sin(bx+c)]+c$$

(iv) 
$$2x \tan^{-1} 3x - \frac{1}{3} \log |1 + 9x^2| + c$$

(v) 
$$2\left[\sqrt{x}\sin\sqrt{x} + \cos\sqrt{x}\right] + c$$

(vi) 
$$\left(\frac{x^4-1}{4}\right) \tan^{-1} x - \frac{x^3}{12} + \frac{x}{4} + c$$

(vii) 
$$\frac{1}{2}e^{2x}\tan x + c$$

(viii) 
$$\frac{x}{\log x} + c$$

(ix) 
$$\left(\frac{x-a}{2}\right)\sqrt{2ax-x^x} + \frac{a^2}{2}\sin^{-1}\left(\frac{x-a}{a}\right) + c$$

(x) 
$$e^{x}\left(\frac{x-1}{x+1}\right) + c$$

(xi) 
$$\frac{x^4}{4}\sin^{-1}\left(\frac{1}{x}\right) + \frac{x^2+2}{12}\sqrt{x^2-1} + c$$

(Xii) 
$$x \log |\log x| - \frac{x}{\log x} + c$$
  
(Xiii)  $-2(6 + x - x^2)^{\frac{3}{2}} + 8^{\left[\frac{2x-1}{3}\sqrt{6 + x - x^2} + \frac{25}{5}\sin^{-1}\right]}$ 

(Xiii) 
$$-2(6 + x - x^2)^{\frac{3}{2}} + 8\left[\frac{2x-1}{4}\sqrt{6 + x - x^2} + \frac{25}{8}\sin^{-1}\left(\frac{2x-1}{5}\right)\right] + c$$

(xiv) 
$$\frac{1}{3}\log|x+1| - \frac{1}{6}\log|x^2 - x+1| + \frac{1}{\sqrt{3}}\tan^{-1}\left(\frac{2x-1}{\sqrt{3}}\right) + c$$
  
(xv)  $x\tan^{-1}x - \frac{1}{2}\log|1 + x^2| - x\tan^{-1}5 + c$   
(xvi)  $\frac{2}{3}\tan^{-1}\left(\frac{1}{3}\tan\frac{x}{2}\right) + c$ 

5. (i) 
$$\frac{1}{20}\log 3$$

(ii) 
$$-\pi/4$$

- (iii)  $\frac{\pi}{4} \frac{1}{2}$
- (iv)  $\frac{\pi}{4} \frac{1}{2}\log 2$
- (v)  $\frac{\pi}{2}$
- (vi)  $\pi/4$
- (vii)  $\pi/2$
- (viii)  $\frac{3}{4} + \frac{3}{2}\log\frac{2}{3}$
- (ix)  $\frac{3}{2\pi} \frac{1}{\pi^2}$
- (x)  $2\pi + \frac{1}{2a}\sin 2a\pi \frac{1}{2b}\sin 2b\pi$
- 6. (i)  $\frac{1}{2}$ 
  - (ii)
  - (iii)  $e^{\pi/4} + e^{-\pi/4}$

π

- (iv)  $\frac{1}{4}\pi^2$ 
  - (v)  $5 \sqrt{3} \sqrt{2}$

$$\begin{array}{rcl} (\mathrm{vi}) & \frac{\pi^2}{16} & (\mathrm{vii}) & \frac{\pi^2}{2a} \\ (\mathrm{i}) & \frac{\pi}{12} & (\mathrm{i}) & 2 \\ (\mathrm{ii}) & \frac{\pi}{2} & (\mathrm{iv}) & \frac{\pi^2}{4} \\ (\mathrm{v}) & a\pi \\ \mathbf{8.} & (\mathrm{i}) & \frac{2(2x-1)}{\pi} \sin^{-1} \sqrt{x} + \frac{2\sqrt{x-x^2}}{\pi} - x + c \\ (\mathrm{ii}) & -2\sqrt{1-x} + \cos^{-1} \sqrt{x} + \sqrt{x-x^2} + c \\ (\mathrm{iii}) & \frac{x-2}{x+2} e^x + c \\ (\mathrm{iii}) & \frac{x+2}{x+2} e^x + c \\ (\mathrm{iv}) & \frac{\sin x - x \cos x}{x \sin x + \cos x} + c \\ (\mathrm{v}) & (x+a) \tan^{-1} \sqrt{\frac{x}{a}} - \sqrt{ax} + c \\ (\mathrm{vi}) & 2 \sin^{-1} \frac{\sqrt{3}-1}{2} \\ (\mathrm{vii}) & \frac{1}{8} \log \left| \frac{1-\sin x}{1+\sin x} \right| - \frac{1}{4\sqrt{2}} \log \left| \frac{1+\sqrt{2} \sin x}{1-\sqrt{2} \sin x} \right| + c \\ (\mathrm{viii}) & \frac{1}{\pi} + \frac{1}{\pi^2} \\ (\mathrm{ix}) & (\cos 2a)(x+a) - (\sin 2a) \log|\sin(x+a)| + c \\ (x) & -\frac{4}{5} \log|x^2 + 4| + \frac{9}{5} \log|x^2 + 9| + c \\ (xi) & -\left(\frac{1}{2} \sin 2x + \sin x\right) + c \\ \mathbf{9.} & (\mathrm{i}) & x - 4 \log|x| + \frac{5}{4} \log|x - 1| + \frac{3}{4} \log|x + 1| + \log|x^2 + 1| \\ (\mathrm{ii}) & \frac{-1}{2} \tan^{-1} x + c \\ (\mathrm{iii}) & \log \left| \frac{(e^x - 1)(e^x - 3)}{(e^x - 2)^2} \right| + c \end{array}$$

[Class XII : Maths]

	<i>(</i> ; )	$2^{1}$	<sup>81</sup> 1	27	
	(1V)	$2x - \frac{100}{8} x + 1  +$	$\frac{-100}{8}$ $\frac{-31}{-3}$	$\frac{1}{2(x-3)} + C$	
	(v)	$\frac{1}{4}\log\left \frac{1-\cos x}{1+\cos}\right  + \frac{1}{2(1-x)}$	$\frac{1}{1+\cos(x)}$ + $\tan(\frac{x}{2})$ + $\tan(\frac{x}{2})$ + $\cos(x)$	2	
	(vi)	$\frac{\pi}{\sqrt{2}}$	(vii)	$\frac{\pi-2}{4}$	
	(viii)	$\frac{\pi}{4} - \frac{1}{2}\log 2$			
10.	(i)	14	(ii)	$\frac{26}{3}$	
	(iii)	26	(v)	$\frac{1}{3}\left(e^2-\frac{1}{e}\right)$	
	(iv)	$\frac{1}{2}(127 + e^8)$	(vi)	3	
11.	(i)	$\frac{1}{5} \log \left  \frac{\tan x - 2}{2 \tan x + 1} \right  + c$	(ii)	$\frac{\pi}{8}\log 2$	
	(iii)	$\frac{\pi}{2}\log\frac{1}{2}$			
12.	$\frac{\pi^2}{16} - \frac{\pi}{4}$	$+\frac{1}{2}\log 2$			
13.	$\frac{-\pi}{2}\log$	2			
14.	log 2				
15.	$\frac{1}{\sqrt{2}}\log$	$\sqrt{2}+1$			
	SELF ASSESSMENT TEST-1				
1. (c)		2. (d)	3. (c)	4. (a)	

1. (c)	2. (d)	3. (c)	4. (a)	5. (d)			
SELF ASSESSMENT TEST-2							
1. (a)	2. (c)	3. (a)	4. (b)	5. (c)			

# **CHAPTER 8**

# **APPLICATIONS OF INTEGRALS**

In real life, integrations are used in various fields such as engineering, where engineers use integrals to find the shape of building. In Physics, used in the centre of gravity etc. In the field of graphical representation. Where three-dimensional models are demonstrated.

The PETRONAS TOWERS in KUALA LUMPUR experience high forces due to wind. Integration was used to create this design of building.



#### APPLICATIONS OF INTEGRALS

Topics to be covered as per C.B.S.E. revised syllabus (2024-25)

• Applications in finding the area under simple curves, especially lines, circles/ parabolas/ellipse (in standard form only)

### POINTS TO REMEMBER

#### AREAS OF BOUNDED REGIONS

Area bounded by the curve y = f(x), the x axis and between the ordinates, x = a and x = b is given by



[Class XII : Maths]

Area bounded by the curve x = f(y), the y-axis and between the abscissas, y
 = c and y = d is given by



• Area bounded by two curves y = f(x) and y = g(x) such that  $0 \le g(x) \le f(x)$ for all  $x \in [a, b]$  and between the ordinates x = a and x = b is given by



#### Illustration:

Using integration. Find the area of the region bounded by the line 2y + x = 8, the x-axis and the lines x = 2 and x = 4

Solution: Required area= Areaof PQRS

= Area bounded by the line 2y + x = 8, x-axis and ordinates x = 2, x = 4

$$= \int_{2}^{4} y \, dx = \int_{2}^{4} \frac{8 - x}{2} \, dx$$
  
=  $\frac{1}{2} \left[ 8x - \frac{x^2}{2} \right]_{2}^{4} = \frac{1}{2} \left[ (32 - 8) - (16 - 2) \right]$   
=  $\frac{1}{2} \left[ 24 - 14 \right] = \frac{1}{2} \times 10 = 5 \text{ sq. units}$ 

#### Illustration:

Draw a rough sketch of the curves  $y = \sin x$  and  $y = \cos x$  as x-varies from 0 to  $\pi/2$ . Find the area of the region enclosed by the curves and the x-axis.



#### Illustration:

Using integration, find the area of the region bounded by the parabola  $y^2 = 16x$  and the line x = 4.

Solution: Given curve  $y^2 = 16x$ line x = 4Area of shaded region = 2(area of AOC) $= 2\int_{1}^{4} y \, dx = 2\int_{1}^{4} 4\sqrt{x} \, dx$ 

 $= 8 \times \frac{2}{3} \left[ x^{3/2} \right]_{0}^{4} = \frac{16}{3} \left[ 8 \right] = \frac{128}{3}$  sq.units



[Class XII : Maths]



#### **ONE MARK QUESTIONS**

# Multiple Choice Questions (1 Mark Each)

#### Select the correct option out of the four given options:

 The area of the region bounded by the curve y = x<sup>2</sup>, x-axis and the lines x = -1, x = 1 is

(a) $\frac{1}{3}$ sq. units	(b) $\frac{2}{3}sq$ . units
(c) 1 sq. unit	(d) 2 sq. units

2. The area bounded by  $y = \sin 2x$ ,  $0 \le x \le \frac{\pi}{4}$  and coordinate axes is (a)  $\frac{1}{2}$  sq. units (b) 1 sq. unit (c)  $\frac{3}{2}$  sq. units (d) 2 sq. units 3. The area bounded by the line x + 2y = 8 and the lines x = 1 and x = 3 is (a) 16 sq. units (b) 8 sq. units (c) 12 sq. units (d) 6 sq. units

4. The area enclosed by the parabola  $y^2 = 8x$  and its latus rectum is

(a)	$\frac{16}{3}$ sq. units	(b) $\frac{64}{3}$ sq. units
(c)	$\frac{32}{3}$ sq. units	(d) $\frac{16\sqrt{2}}{3}$ sq. units

5. The area bounded by the curve  $y = \cos x$  and x-axis between x = 0 and  $x = \pi$  is

(a) 0 sq. units	(b) 1 sq. units			
(c) 2 sq. units	(d) 4 sq. units			

#### ASSERTION-REASON BASED QUESTIONS

In the following qustions a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices:

- (a) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (b) Both (A) and (R) are true, but (R) is not the correct explanation of (A)
- (c) (A) is true and (R) is false
- (d) (A) is false, but (R) is true
- 6. Assertion (A) : Area enclosed by the curve  $x^2 + y^2 = 4$  is given by  $4 \int_0^2 \sqrt{4 x^2} dx$

Reason (R) : The curve  $x^2 + y^2 = 4$  is symmetric about both the axes.

7. Assertion (A): Area of the region bounded by the parabola  $y^2 = 4x$  and its latus rectum

is given by  $2\int_{0}^{1} 2\sqrt{x} dx$ 

Reason (R) : Length of the latus rectum of the parabola  $y^2 = 4ax$  is 4a.

#### TWO MARKS QUESTIONS

1. Find the area of the circle  $x^2 + y^2 = 16$ .

**Using Integration:** 

- 2. Find the area of the parabols  $y^2 = 4a \times bounded by its latus rectum.$
- 3. Find the area bounded by the curve  $y^2 = x, x axis$  and the lines x = 0, x = 4.
- 4. Find the area bounded by the region  $\{(x, y): x^2 \le y \le |x|\}$ .
- 5. Find the area bounded by the region  $y = 9x^2$ , y = 1 and y = 4.
- 6. Find the area bounded by the curve  $y = \sin x$  between  $x = \frac{\pi}{2}$  and  $x = \frac{3\pi}{2}$
- 7. Find the area bounded by the lines y = 2x + 3, y = 0, x = 2 and x = 4.
- 8. Find the area of the region bounded by  $y^2 = 4x$ , x = 1, x = 4 and x-axis in the first quadrant.
- 9. Find the area bounded by the curves  $y^2 = 4ax$  and the lines y = 2a and y-axis.
- 10. Find the area of the triangle formed by the straight lines y = 2x, x = 0 and y = 2

#### THREE/FIVE MARKS QUESTIONS Using Integration

- 1. Find the area bounded by the curve  $4y = 3x^2$  and the line 3x 2y + 12 = 0.
- 2. Find the area bounded by the curve  $x = y^2$  and the line x + y = 2.
- 3. Find the area of the triangular region whose vertices are (1, 2), (2-2) and (4, 3).
- 4. Find the area bounded by the region { $(x, y): x^2 + y^2 \le 1 \le x + \frac{y}{2}$  }
- 5. Find the area of the region bounded by the lines x 2y = 1, 3x y 3 = 0 and 2x + y 12 = 0.
- 6. Prove that the curve  $y = x^2$  and,  $x = y^2$  divide the square bounded by x = 0, y = 0, x = 1, y = 1 into three equal parts.
- 7. Find the area of the smaller region enclosed between ellipse  $b^2x^2 + a^2y^2 = a^2b^2$  and the line bx + ay = ab.
- 8. Using integration, find the area of the triangle whose sides are given by 2x + y = 4, 3x 2y = 6 and x 3y + 5 = 0.
- 9. Using integration, find the area of the triangle whose vertices are (-1, 0), (1, 3) and (3, 2).

- 10. Find the area of the region  $\{(x, y) : x^2 + y^2 \le 1 \le x + y\}$ .
- 11. Find the area of the region bounded by the curve  $x^2 = 4y$  and the line x = 4y 2.
- 12. Using integration, find the area of the region bounded by the line x y + 2 = 0, the curve  $x^2 = y$  and y-axis.
- 13. Using integration, find the area of the region bounded by the curve y = 1 + |x + 1| and lines x = -3, x = 3, y = 0.
- 14. Find the area of the region enclosed between curves y = |x 1| and y = 3 |x|.
- 15. If the area bounded by the parabola  $y^2 = 16$  ax and the line y = 4 mx is  $\frac{a^2}{12}$  sq unit then using integration find the value of m.
- 16. Find the area bounded by the circle  $x^2 + y^2 = 16$  and the line y = x and x-axis in first quadrant.
- 17. Find the area bounded by the parabola  $y^2 = 4x$  and the straight line x + y = 3.
- 18. Find the area bounded by the parabola  $y^2 = 4x$  and the line y = 2x 4.

19. Find the area of region 
$$\left\{ (x,y): \frac{x^2}{9} + \frac{y^2}{4} \le 1 \le \frac{x}{3} + \frac{y}{2} \right\}$$

20. Using integration, find the area of the triangle ABC, whose vertices are A(2, 5), B(4, 7) and C(6, 2).

#### **SELF ASSESSMENT-1**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. Area of the region bounded by the curve  $y^2 = 4x$ , y-axis and the line y = 3 is

(a)	$\frac{9}{2}$ sq. units	(b)	$\frac{9}{3}$ sq. units
(c)	$\frac{9}{4}$ sq. units	(d)	$\frac{9}{5}$ sq. units

2. Area lying in first quadrant and bounded by the circle  $x^2 + y^2 = 4$  and the lines x = 0 and x = 2 is

(a) π	sq. units	(b)	$\frac{\pi}{3}$ sq. units
(c) $\frac{\pi}{2}$	sq. units	(d)	$\frac{\pi}{4}$ sq. units

3. The area of the region bounded by the curve y = x + 1 and the lines x = 2, x = 3 and x-axis is

(a) 
$$\frac{13}{2}$$
 sq.units (b)  $\frac{11}{2}$  sq.units

(c) 
$$\frac{9}{2}$$
 sq.units (d)  $\frac{7}{2}$  sq.units

4. The area bounded by the curve  $y^2 = x$  and the line x = 2y is

(a) 
$$\frac{1}{3}$$
 sq.units  
(b)  $\frac{2}{3}$  sq.units  
(c) 1 sq.unit  
(d)  $\frac{4}{3}$  sq.units

5. The area of the region bounded by the  $y = \sin x$ ,  $y = \cos x$  and y-axis,  $0 \le x \le \frac{\pi}{4}$  is (a)  $(\sqrt{2} + 1)$  sq.units (b)  $(\sqrt{2} - 1)$  sq.units

- (d)  $(2\sqrt{2} 1)$  sq.units (c)  $2\sqrt{2}$  sq.units

# **ANSWERS**

#### ONE MARKS QUESTION

1.	(b) $\frac{2}{3}$ square units	2.	(a)	$\frac{1}{2}$	square units.
3.	(d) 6 square units	4.	(c)	$\frac{32}{3}$	square units.
5.	(c) 2 square units	6.	(a)	-	
7.	(b)				
	TWO MARKS QUES	тю	NS		
1.	16 $\pi$ square units.				
2.	$\frac{8}{3}a^2$ square units.				
3.	$\frac{16}{3}$ square units.				
4.	$\frac{1}{3}$ square units.				
5.	28 9 square units.				
6.	2 square units.				
7.	18 square units.				
8.	$\frac{28}{3}$ square units.				
9.	$\frac{2}{3}a^2$ square units.				
10.	1 square units.				
	THREE/FIVE MARKS Q	UES	бтіо	NS	i

1. 27 square units.

2. 
$$\frac{9}{2}$$
 square units.

3. 
$$\frac{13}{2}$$
 square units.
4. 
$$\left(\frac{\pi}{4}-\frac{2}{5}-\frac{1}{2}\sin^{\prime}\frac{3}{5}\right)$$
 square units.

5. 10 square units.

7. 
$$\left(\frac{\pi-2}{4}\right)$$
ab square units.

- 8. 3.5 square units.
- 9. 4 square units.

10. 
$$\left(\pi - \frac{1}{2}\right)$$
 square units.

11. 
$$\frac{9}{8}$$
 square units.

12. 
$$\frac{10}{3}$$
 square units.

- 13. 16 square units.
- 14. 4 square units.
- 15. m = 2.
- 16.  $2\pi$  sq. units
- 17.  $\frac{64}{3}$  sq. units
- 18. 9 sq. units
- 19.  $\frac{3}{2}(\pi 2)$  sq. units
- 20. 7 sq. units

### **SELF ASSESSMENT TEST-1**

1.	(C)	2. (A)	3. (D)	4. (D)	5. (B)

### **CHAPTER-9**

### **DIFFERENTIAL EQUATIONS**

Sky diving is a method of transiting from a high point in the atmosphere to the surface of the Earth with the aid of graity. This involves the control of speed during the descent using a parachute. Once the sky diver jumps from an airplane, the net force experienced by the diver can be calculated using

#### DIFFERENTIAL EQUATIONS.

Another eg.



D.E. is

my'' = *mg* 

 $\Rightarrow$  y'' = g = constant

where y = distance travelled by the stone at any time t.

and g = acceleration due to gravity.

### TOPICS TO BE COVERED AS PER CBSE LATEST CURRICULUM 2023-24

- Definition, order and degree
- General and particular solutions of a D.E.
- Solutions of D.E. using method of separation of variables.
- Solutions of homogeneous differential equations of first order and first degree.
- Solutions of linear differential equations of the type.

$$\frac{dy}{dx} + py = q$$
, where p and q are functions of x or constants.

$$\frac{dx}{dy} + px = q$$
, where p and q are functions of y or constants.



**KEY POINTS:** 

- DIFFERENTIAL EQUATION : is an equation involving derivatives of the dependent variable w.r.t independent variables and the variables themselves.
- ORDINARY DIFFERENTIAL EQUATION (ODE): A.D.E. involving derivatives of the dependent variable w.r.t only one independent variable is an ordinary D.E.

In class XII ODE is referred to as D.E.

- PARTIAL DIFFERENTIAL EQUATION (PDE): A.D.E involving derivatives w.r.t more than one independent variables is called a partial D.E.
- ORDER of a DE : is the order of the highest order derivative occurring in the D.E.
- DEGREE of a D.E.: is the highest power of the highest order derivative occurring in the D.E provided D.E is a polynomial equation in its derivatives. It is always a whole no.
- SOLUTION OF THE D.E : A relation between involved variables, which satisfy the given D.E is called its solution.



Two Types of Solutions of DE

- FORMATION OF A DIFFERENTIAL EQUATION : We differentiate the function successively as many times as the arbitary constants in the given function and then eliminate the arbitiary constants from these equations.
- ORDER of A D.E : Is equal to the number of arbitrary constants in the general solution of a D.E.



- "VARIABLE SEPARABLE METHOD" : is used to solve D.E. in which variables can be separated completely i.e, terms containing x should remain with dx and terms containing y should remain with dy.
- "HOMOGENEOUS DIFFERENTIAL EQUATION : D.E. of the form  $\frac{dy}{dx} = F(x, y)$ where F(x, y) is a homogeneous function of degree 0

i.e.  $F(\lambda x, \lambda y) = \lambda^0 F(x, y)$ 

or  $F(\lambda x, \lambda y) = F(x, y)$  for some non-zero constant  $\lambda$ .

To solve this type put y = vx

To Solve homogenous D.E of the trype  $\frac{dx}{dy} = G(x, y)$ , we make substitution x = vy

• LINEAR DIFFERENTIAL EQUATION : A.D.E of the form  $\frac{dy}{dx} + Py = Q$  where P and Q are constants or functions of x only is known as first order linear differential equation.

Its solution

$$y.(IF) = \int Q \times (I.F.) dx + C$$
, where

I. F = Integrating factor =  $e^{\int Pdx}$ 

Another form of Linear Differential Equation is  $\frac{dx}{dy} + P_1 x = Q_1$ , where  $P_1$  and

 $Q_1$  are constants or functions of y only.

Its solution is given as

$$x.(I.F) = \int Q_1 X(I.F.) dy + C$$
, where  $I.F. = e^{\int P_1 dy}$ 

### Illustration:

Write the order and degree of the Differential Equation

$$\left[1+(y')^2\right]^{3/2}=ky''$$

**Solution:** Squaring both the sides

$$\left\lceil 1 + (y')^2 \right\rceil^3 = k^2 (y'')^2$$

∴ Order of D.E. = 2

and Degree of D.E. = 2

Illustration: Solve the differential equations  $(1+e^{2x})dy+e^{x}(1+y^{2})dx=0; y(0)=1$ **Solution:**  $\frac{dy}{dx} = \frac{-e^x(1+y^2)}{1+e^{2x}}$ Using Variables separables method,  $\frac{dy}{1+y^2} = \frac{-e^x}{1+e^{2x}}dx$ Integrating both sides we get  $\int \frac{1}{1+y^2} dy = -\int \frac{e^x}{1+e^{2x}} dx$  $\Rightarrow \tan^{-1}y = -\int \frac{dt}{1+t^2}; \text{ On putting } e^x = t$ = – tan⁻¹*t*  $\Rightarrow \tan^{-1}y = -\tan^{-1}(e^x) + C$  $\Rightarrow \tan^{-1}y + \tan^{-1}(e^x) = C$ At x = 0, y = 1 given :.  $tan^{-1}(1) + tan^{-1}(1) = C$  $\Rightarrow \frac{\pi}{4} \times 2 = C$  $\Rightarrow$  C =  $\frac{\pi}{2}$ :. Particular solution of D.E. is given by  $\tan^{-1}y + \tan^{-1}(e^x) = \frac{\pi}{2}$ . Illustration: Solve  $(x - y) \frac{dy}{dx} = x + 2y$ Solution:  $\frac{dy}{dx} = \frac{x+2y}{x-y} = f(x,y)$ Now  $f(\lambda x, \lambda y) = \frac{\lambda x + 2\lambda y}{\lambda x - \lambda y} = \frac{\lambda(x + 2y)}{\lambda(x - y)} = \lambda^{\circ} f(x, y)$ 

[Class XII : Maths]

165

Clearly, f is homogeneous function in x and y. So, given D.E. is homogenous D.E. Now, Put y = vx $\Rightarrow \frac{dy}{dx} = v + \frac{x \, dv}{dx}$  $\therefore v + \frac{x \, dv}{dx} = \frac{x + 2vx}{x - vx}$  $\Rightarrow v + \frac{x \, dv}{dx} = \frac{1 + 2v}{1 - v}$  $\Rightarrow \frac{x \, dv}{dx} = \frac{1 + 2v - v + v^2}{1 - v}$  $\Rightarrow \frac{x \, dv}{dx} = \frac{1 + v + v^2}{1 - v}$  $\Rightarrow \frac{(1-v)dv}{1+v+v^2} = \frac{dx}{x}$ Integrating both sides we get  $\Rightarrow -\frac{1}{2} \int \frac{2v - 2 + 1 - 1}{1 + v + v^2} dv = \log |x| + C$  $\Rightarrow -\frac{1}{2}\int \frac{2v+1}{1+v+v^2}dv + \frac{3}{2}\int \frac{1}{1+v+v^2}dv = \log |x| + C$  $\Rightarrow -\frac{1}{2}\log|1+v+v^2| + \frac{3}{2}\int \frac{1}{\left(v+\frac{1}{2}\right)^2 + \left(\frac{\sqrt{3}}{2}\right)^2} dv = \log|x| + C$  $\Rightarrow -\frac{1}{2}\log\left|1+\frac{y}{x}+\frac{y^2}{x^2}\right|+\left(\frac{3}{2}\right)\cdot\left(\frac{2}{\sqrt{3}}\right)\tan^{-1}\left(\frac{2\nu+1}{\sqrt{3}}\right) = \log|x|+C$  $\Rightarrow -\frac{1}{2}\log |x^2 + xy + y^2| + \sqrt{3}\tan^{-1}\left(\frac{2y + x}{\sqrt{3}x}\right) = C$ 

# Illustration: Find the particular solution of the differential equation $\frac{dx}{dy} + x \cot y = 2y + y^2 \cot y (y \neq 0)$ given that x = 0 when $y = \pi/2$ . Solution: Clearly, it is a Linear D.E. $\frac{dx}{dv} + Px = Q$ where $P = \cot y, Q = 2y + y^2 \cot y$ I.F. = $e^{\int^{Pdy}} = e^{\int^{\cot y \, dy}} = e^{\log(\sin y)} = \sin y$ ... solution of D.E. is given by x. (I.F) = $\int Q.IFdy + C$ ; C is arbitrary constant $\Rightarrow x. (\sin y) = \int (2y + y^2 \cot y) \sin y \, dy + C$ $= \int 2y \sin y \, dy + \int y^2 \cos y \, dy + C$ = $\int 2y \sin y \, dy + y^2 . \sin y - \int 2y \sin y \, dy + C$ $\Rightarrow$ x sin y = y<sup>2</sup> sin y + C Now, x = 0, when $y = \frac{\pi}{2}$ So, $0 = \frac{\pi^2}{4} + C \Rightarrow C = -\frac{\pi^2}{4}$ $\therefore \quad x \sin y = y^2 \sin y - \frac{\pi^2}{4}$ or $x = y^2 - \frac{\pi^2}{4}$ cosec y

### **ONE MARK QUESTIONS**

1. The general solution of the D.E.

y dx - xdy = 0; (Given x, y > 0), is of the form.

(a) xy = c (b)  $x = cy^2$ 

(c) 
$$y = cx$$
 (d)  $y = cx^{2}$ 

(Where 'c' is an orbitary positive constant of integration)

2. The differential equation  $\frac{dy}{dx} = \frac{\sqrt{1-y^2}}{y}$  determines a family of circles with

- (a) Variable radii and fixed centre (0, 1)
- (b) Variable radii and fixed centre (0, -1)
- (c) Fixed radius 1 and variable centre on x-axis
- (d) Fixed radius 1 and variable centre on y-axis

3. The solution of the D.E. 
$$\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$$
 is

(a) 
$$e^{x} = \frac{y^{3}}{3} + e^{y} + c$$
 (b)  $e^{y} = \frac{x^{2}}{3} + e^{x} + c$ 

(c) 
$$e^{y} = \frac{x^{3}}{3} + e^{x} + c$$
 (d) None f these

- 4. The order and degree of the D.E.  $\frac{d^4y}{dx^4} + \sin(y''') = 0$  are respectively
  - (a) 4 and 1 (b) 1 and 2
  - (c) 4 and 4 (d) 4 and not defined
- 5. A homogeneous differential equation of the type  $\frac{dx}{dy} = h\left(\frac{x}{y}\right)$  can be solved by making the substitution.

- (a) y = vx (b) v = yx
- (c) x = vy (d) x = v

6. Integrating factor of the D.E. 
$$\frac{dy}{dx} + y \tan x - \sec x = 0$$
 is  
(a)  $\cos x$  (b)  $\sec x$   
(c)  $e^{\cos x}$  (d)  $e^{\sec x}$ 

7. The order and degree of the D.E.  $\frac{d^2y}{dx^2} + \left(\frac{dy}{dx}\right)^{\frac{1}{4}} + x^{\frac{1}{6}} = 0$ , respectively are

(a)	2 and not defined	(b) 2 and 2
(c)	2 and 3	(c) 3 and 3

8. The order of the D.E. of a family of curves respresented by an equation containing four arbitrary constants, will be

(c) 6	(d) None of these
· · ·	

- 9. An equation which involves variable as well as dirivatives of the dependent variable w.r.t. the independent variable, is known as
  - (a) differential equation (b) integral equation
  - (c) linear equation (d) quadantic equation
- 10.  $\tan^{-1} x + \tan^{-1} y = c$  is general solution of the D.E.
- (a)  $\frac{dy}{dx} = \frac{1+y^2}{1+x^2}$ (b)  $\frac{dy}{dx} = \frac{1+x^2}{1+y^2}$ (c)  $(1+x^2)dy + (1+y^2)dx = 0$ (d)  $(1+x^2)dx + (1+y^2)dy = 0$ 11. The particular solution of  $\log\left(\frac{dy}{dx}\right) = 3x + 4y, y(0) = 0$  is (a)  $e^{3x} + 3e^{-4y} = 4$ (b)  $4e^{4x} + 3e^{-4y} = 3$ (c)  $3e^{3x} - 4e^{4y} = 7$ (d)  $4e^{3x} + 3e^{-4y} = 7$ 12. The solution of the equation  $\frac{dy}{dx} = \frac{3x - 4y - 2}{3x - 4y - 3}$  is (a)  $(x - y^2) + c = \log(3x - 4y + 1)$ (b)  $x - y + c = \log(3x - 4y + 4)$ (c)  $(x - y + c) = \log(3x - 4y - 3)$ (d)  $x - y + c = \log(3x - 4y + 1)$

13. If  $x \frac{dy}{dx} = y(\log y - \log x + 1)$ , then the solution of the equation is

(a) 
$$y \log\left(\frac{x}{y}\right) = cx$$
 (b)  $x \log\left(\frac{y}{x}\right) = cy$ 

(c) 
$$\log\left(\frac{y}{x}\right) = cx$$
 (d)  $\log\left(\frac{x}{y}\right) = cy$ 

- 14. Solution of D.E. xdy ydx = 0 respresents
  - (a) rectangular hyperbola (b) parabola whose vertex is at orgain
  - (c) circle whose centre is at origin (d) straight line passing through origin
- 15. Family  $y = bx + c^4$  of curves will correspond to a differential equation of order
  - (a) 3 (b) 2

16. The integrating factor of the differential equation  $(1 - y^2)\frac{dx}{dy} + yx = ay, (-1 < y < 1)$ 

is :

(a) 
$$\frac{1}{y^2 - 1}$$
 (b)  $\frac{1}{\sqrt{y^2 - 1}}$ 

(c) 
$$\frac{1}{1-y^2}$$
 (d)  $\frac{1}{\sqrt{1-y^2}}$ 

17. The general solution of the differential equation  $xdy - (1 + x^2)dx = 0$  is

(a) 
$$y = 2x + \frac{x^3}{3} + c$$
  
(b)  $y = 2\log x + \frac{x^3}{3} + c$   
(c)  $y = \frac{x^2}{2} + c$   
(d)  $y = \log x + \frac{x^2}{2} + c$ 

### ASSERTION REASON TYPE QUESTIONS

**Directions :** Each of these questions contains two statements, Assertion (A) and Reason (R) Choose the correct answer out of the following choices.

- (a) Both (A) and (R) are true and (R) is the correct explantion of (A)
- (b) Both (A) and (R) are true but (R) is not the correct explanation of (A)
- (c) (A) is true and (R) is false
- (d) (A) is false but (R) is true
- 18. Assertion (A): Order of the differential equation whose solution is  $y = c_1 e^{x+c_2} + c_3 e^{x+c_4}$  is 4.

Reason (R) : Order of the differential equation is equal to the number of independent arbitrary constant mentioned in the solution of differential equation.

19. Assertion (A): The degree of the differential equation  $\frac{d^2 y}{dx^2} + 3\left(\frac{dy}{dx}\right)^2 = x^2 \log\left(\frac{d^2 y}{dx^2}\right)$ 

is not defined.

Reason (R) : If the differential equation is a polynomial in terms of its derivatives, then its degree is defined.

20. Assertion (A):  $\frac{dy}{dx} = \frac{x^3 - xy^2 + y^3}{x^2y - x^3}$  is a homogeneous differential equation.

Reason (R): The function  $f(x,y) = \frac{x^3 - xy^2 + y^3}{x^2y - x^3}$  is homogeneous.

### **TWO MARKS QUESTIONS**

- 1. Write the general solution of the following D.Eqns.
  - (i)  $\frac{dy}{dx} = x^5 + x^2 \frac{2}{x}$  (ii)  $\frac{dy}{dx} = \frac{1 \cos 2x}{1 + \cos 2y}$ (iii)  $(e^x + e^{-x}) dy = (e^x - e^{-x}) dx$
- 2. Given that  $\frac{dy}{dx} = e^{-2y}$  and y = 0 when x = 5.

Find the value of x when y = 3.

- Name the curve for which the slope of the tangent at any point is equal to the ratio of the abbcissa to the ordinate of the point.
- 4. Solve  $\frac{xdy}{dx} + y = e^x$ .

### THREE MARKS QUESTIONS

1. (i) Show that  $y = e^{m \sin^{-1}x}$  is a solution of  $(1 - x^2) \frac{d^2 y}{dx^2} - \frac{x dy}{dx} - m^2 y = 0$ 

(ii) Show that  $y = a \cos(\log x) + b \sin(\log x)$  is a solution of

$$\frac{x^2d^2y}{dx^2} + \frac{xdy}{dx} + y = 0$$

(iii) Verify that y = log  $(x + \sqrt{x^2 + a^2})$  satisfies the D.E.

$$\left(a^2+x^2\right)y''+xy'=0$$

2. Solve the following differential equations.

(i) 
$$xdy - (y + 2x^{2})dx = 0$$
  
(ii)  $(1 + y^{2})tan^{-1}x dx + 2y(1 + x^{2})dy = 0$   
(iii)  $x^{2}\frac{dy}{dx} = x^{2} + xy + y^{2}$ 

(iv) 
$$\frac{dy}{dx} = 1 + x + y^2 + xy^2$$
,  $y = 0$  when  $x = 0$ 

(v) 
$$xdy - ydx = \sqrt{x^2 + y^2} dx, y = 0$$
 when  $x = 1$ 

3. Solve each of the following differential equations

(i) 
$$(1 + x^2)\frac{dy}{dx} + 2xy - 4x^2 = 0, y(0) = 0$$
  
(ii)  $(x + 1)\frac{dy}{dx} = 2e^{-y} - 1, y(0) = 0$   
(iii)  $e^x \tan y dx + (2 - e^x) \sec^2 y dy = 0, y(0) = \frac{\pi}{4}$   
(iv)  $(x^2 - y^2) dx + 2xy dy = 0$   
(v)  $(1 + x^2)\frac{dy}{dx} + 2xy = \frac{1}{1 + x^2}, y = 0$  when  $x = 1$ 

4. Solve the following differential equations (i) Find the particular solution of

$$2y e^{x/y} dx + (y - 2xe^{x/y}) dy = 0, x = 0$$
 if  $y = 1$ 

(ii) 
$$x \cos\left(\frac{y}{x}\right) \frac{dy}{dx} = y \cos\left(\frac{y}{x}\right) + x$$

(iii) 
$$\frac{dy}{dx} = \cos(x+y) + \sin(x+y)$$

[**Hint** : Put *x* + *y* = *z*]

(iv) Show that the Differential Equation  $\frac{dy}{dx} = \frac{y^2}{xy - x^2}$  is homogenous and also solve it.

(v) 
$$(x^2 - 1)\frac{dy}{dx} + 2xy = \frac{1}{x^2 - 1}, |x| \neq 1$$

### **FIVE MARKS QUESTIONS**

Q. 1 Solve 
$$y + \frac{d}{dx}(xy) = x(\sin x + \log x)$$
  
Q. 2 Solve  $(x \, dy - y dx)y \sin\left(\frac{y}{x}\right) = (y dx + x dy)x \cos\left(\frac{y}{x}\right)$ 

Q. 3 Find the particular solution of the D.E.  $(x - y)\frac{dy}{dx} = x + 2y$  given that y = 0 when x = 1.

Q. 4 Solve  $dy = \cos x (2 - y \cos ec x) dx$ , given that y = 2 when  $x = \pi/2$ 

Q. 5 Find the particular solution of the D.E.  $(1+y^2) + (x - e^{\tan^{-1}y})\frac{dy}{dx} = 0$ 

given that y = 0 when x = 1

### CASE STUDY QUESTIONS

1. An equation involving derivatives of the dependent variable w.r.t. the independent variables

is called a differential equation. A differential equation of the from  $\frac{dy}{dx} = f(x, y)$  is said

to be homogeneous if f(x, y) is a homogeneous function of degree zero, whereas a function f(x, y) is a homogeneous function of degree *n* if  $f(\lambda x, \lambda y) = \lambda^n f(x, y)$ . To solve a

homogeneous differential equation of the type  $\frac{dy}{dx} = f(x, y) = g\left(\frac{y}{x}\right)$  we make the

substitution y = vx and then separate the variables.

Based on the above, answer the following quations:

- (i) Show that  $(x^2 y^2)dx + 2xydy = 0$  is a differential equation of the type
  - $\frac{dy}{dx} = g\left(\frac{y}{x}\right)$
- (ii) Solve the above equation to find its general solution.

### **Self Assessment Test-1 Differential Equations**

Q. 1 The general solution of the D.E.

$$\log\left(\frac{dy}{dx}\right) = ax + by \text{ is}$$

(a) 
$$\frac{-e^{-by}}{b} = \frac{e^{ax}}{a} + C$$
 (b)  $e^{ax} - e^{-by} = C$ 

(c) 
$$be^{ax} + ae^{by} = C$$

Q. 2 The general solution of the DE

$$x^{2} \frac{dy}{dx} = x^{2} + xy + y^{2} \text{ is}$$
(a)  $\tan^{-1}\left(\frac{y}{x}\right) = \log x + c$ 
(b)  $\tan^{-1}\left(\frac{x}{y}\right) = \log x + c$ 

(c) 
$$\tan^{-1}\left(\frac{y}{x}\right) = \log y + c$$

Q. 3 The solution of the D.E.

$$dy = (4 + y^{2}) dx \text{ is}$$
(a)  $y = 2 \tan (x + c)$ 
(b)  $y = 2 \tan (2x + c)$ 
(c)  $2y = \tan (2x + c)$ 
(d)  $2y = 2 \tan (x + c)$ 

Q. 4 What is the degree of the D.E.

$$y = x \left(\frac{dy}{dx}\right)^3 + \left(\frac{dy}{dx}\right)^2$$

(a) 1 (b) 3 (c) -2 (d) Degree doesn't exist

- Q. 5 Solution of D.E. xdy ydx = 0 represents:
  - (a) a rectangular hyperbola
  - (b) a parabola whose vertex is at the origin
  - (c) a straight line passing through the origin
  - (d) a circle whose centre is at the origin.

### Self Assessment Test-2

Q. 1	The solution of the D.E.	$x\frac{dy}{dx}$ + 2y = $x^2$ is
------	--------------------------	----------------------------------

(a) 
$$y = \frac{x^2 + c}{4x^2}$$
 (b)  $y = \frac{x^2}{4} + c$ 

(c) 
$$y = \frac{x^2 + c}{x^2}$$
 (d)  $y = \frac{x^4 + c}{4x^2}$ 

Q. 2 The solution of the  $\frac{dy}{dx} + y = e^{-x}$ , y(0) = 0, is

(a) 
$$y = e^{-x}(x-1)$$
 (b)  $y = x e^{x}$   
(c)  $y = xe^{-x} + 1$  (d)  $y = x e^{-x}$ 

Q. 3 If $\frac{dy}{dx} = \frac{2^{x+y} - 2^x}{2^y}$ , $y(0) =$	1, then $y(1)$ is equal to	[JEE mains 2021]
(a) $\log_2(2 + e)$	(b) log <sub>2</sub> (1 + <i>e</i> )	
(c) log <sub>2</sub> (2e)	(d) $\log_2(1 + e^2)$	

Q. 4 If the solution curve of the D.E.  $(2x - 10y^3) dy + ydx = 0$  passess through the points (0, 1) and (2,  $\beta$ ), then  $\beta$  is a root of the equation

(a) 
$$y^5 - 2y - 2 = 0$$
(b)  $2y^5 - 2y - 1 = 0$ (c)  $2y^5 - y^2 - 2 = 0$ (d)  $y^5 - y^2 - 1 = 0$ [JEE mains 2021]

Q. 5 Consider a curve y = f(x) passing through the point (-2, 2) and the slope of the tangent to the curve at any point (*x*, *f*(*x*)) is given by

$$f(x) + xf'(x) = x^2,$$

(a) $x^3 + 2x f(x) - 12 = 0$	(b) $x^3 + xf(x) + 12 = 0$	
(c) $x^3 - 3x f(x) - 4 = 0$	(d) $x^2 + 2xf(x) + 4 = 0$	(HOTS)

#### Answers

#### **ONE MARK QUESTIONS**

1. (c) y = cx	2. (c)	3. (c)	4. (d)
5. (c) x = vy	6. (b)	7. (a)	8. (b)4
9. (a)	10. (c)	11. (d)	12. (d)

13.(c)	14. (d)	15. (b)2	16. (d) $\frac{1}{\sqrt{1-y^2}}$
17.(d)	18. (d)	19. (a)	20. (a)

### TWO MARKS QUESTIONS

1. (i)  $y = \frac{x^6}{6} + \frac{x^3}{3} - 2\log |x| + C$  (ii)  $2(y - x) + \sin 2y + \sin 2x = c$ (iii)  $y = \log_e |e^x + e^{-x}| + C$ 2.  $\frac{e^6 + 9}{2}$  3. Rectangular hyperbola 4.  $y \cdot x = e^x + c$ 

### THREE MARKS QUESTIONS

2. (i) 
$$y = 2x^{2} + cx$$
  
(ii)  $\frac{1}{2}(\tan^{-1}x)^{2} + \log(1+y^{2}) = c$   
(iii)  $\tan^{-1}\left(\frac{y}{x}\right) = \log |x| + c$   
(iv)  $y = tan\left(x + \frac{x^{2}}{2}\right)$   
(v)  $y + \sqrt{x^{2} + y^{2}} = x^{2}$   
3. (i)  $(1 + x^{2})y = \frac{4x^{3}}{3}$   
(ii)  $(2 - e^{y})(x + 1) = 1$   
(iii)  $\tan y = 2 - e^{x}$   
(iv)  $x^{2} + y^{2} = cx$   
(v)  $(1 + x^{2})y = tan^{-1}x - \pi/4$   
4. (i)  $e^{x/y} = \frac{-1}{2}\log |y| + 1$   
(ii)  $sin(y/x) = \log |x| + c$   
(iii)  $\log \left|1 + tan\left(\frac{x + y}{2}\right)\right| = x + c$   
(iv)  $\frac{y}{x} - \log |y| = c$   
(v)  $(x^{2} - 1)y = \frac{1}{2}\log \left|\frac{x - 1}{x + 1}\right| + c$ 

### FIVE MARKS QUESTIONS

1. 
$$y = -\cos x + \frac{2\sin x}{x} + \frac{2\cos x}{x^2} + \frac{x\log x}{3} - \frac{x}{9} + \frac{c}{x^2}$$
  
2.  $xy\cos\left(\frac{y}{x}\right) = c$ 

3. 
$$\sqrt{3}\tan^{-1}\left(\frac{2y+x}{\sqrt{3}x}\right) - \frac{1}{2}\log|x^2 + xy + y^2| = \frac{\sqrt{3}\pi}{6}$$
  
4.  $y\sin x = \frac{-1}{2}\cos(2x) + \frac{3}{2}$   
5.  $x = \frac{1}{2}e^{\tan^{-1}y} + \frac{1}{2}e^{-\tan^{-1}y}$ 

### CASE STUDY QUESTIONS

1. (iii)  $x^2 + y^2 = cx$ ; c is an arbitrary constant

### SELE ASSESSMENT TEST-1

1. (a)	2. (a)	3. (b)
4. (b)	5. (c)	

### SELE ASSESSMENT TEST-2

1. (d)	2. (d)	3. (b)
4. (d)	5. (c)	

### CHAPTER 10

## VECTORS

Vectors are probably the most important tool to learn in all of physics and engineering. Vectors are using in daily life following are few of the example.

- Navigating by air and by boat is generally done using vectors.
- Planes are given a vector to travel, and they use their speed to determine how far they need to go before turning or landing. Flight plans are made using a series of vectors.
- Sports instructions are based on using vectors.





- A quantity that has magnitude as well as direction is called a vector. It is denoted by a directed line segment.
- Two or more vectors which are parallel to same line are called collinear vectors.
- Position vector of a point P(a, b, c) w.r.t. origin (0, 0, 0) is denoted by  $\overrightarrow{OP}$  where  $\overrightarrow{OP} = a\hat{i} + b\hat{j} + c\hat{k}$  and  $|\overrightarrow{OP}| = \sqrt{a^2 + b^2 + c^2}$ .
- If  $A(x_1, y_1, z_1)$  and  $B(x_2, y_2, z_2)$  be any two points in space, then

 $\overrightarrow{AB} = (x_2 - x_1)\hat{i} + (y_2 - y_1)\hat{j} + (z_2 - z_1)\hat{k}$  and

$$\overline{AB} = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

- Any vector  $\vec{a}$  is called unit vector if  $|\vec{a}| = 1$  It is denoted by  $\hat{a}$
- If two vectors  $\vec{a}$  and  $\vec{b}$  are represented in magnitude and direction by the two sides of a triangle in order, then their sum  $\vec{a} + \vec{b}$  is represented in magnitude and direction by third side of a triangle taken in opposite order. This is called triangle law of addition of vectors.
- If  $\vec{a}$  is any vector and  $\lambda$  is a scalar, then  $\lambda \vec{a}$  is vector collinear with  $\vec{a}$  and  $|\lambda \vec{a}| = |\lambda| |\vec{a}|$ .
- If  $\vec{a}$  and  $\vec{b}$  are two collinear vectors, then  $\vec{a} = \lambda \vec{b}$  where  $\lambda$  is some non-zero scalar.

- Any vector  $\vec{a}$  can be written as  $\vec{a} = |\vec{a}|\hat{a}$  where  $\hat{a}$  is a unit vector in the direction of  $\vec{a}$ .
- If  $\vec{a}$  and  $\vec{b}$  be the position vectors of points A and B, and C is any point which divides  $\overrightarrow{AB}$  in ratio m:n internally then position vector  $\vec{c}$  of point C is given as  $\vec{c} = \frac{m\vec{b}+n\vec{a}}{m+n}$ . If C divides  $\overrightarrow{AB}$  in ratio m:n externally, then  $\vec{c} = \frac{m\vec{b}-n\vec{a}}{m-n}$ . If C is mid point then  $\vec{c} = \frac{\vec{a}+\vec{b}}{2}$
- The angles  $\alpha$ ,  $\beta$  and  $\gamma$  made by  $\vec{r} = a\hat{i} + b\hat{j} + c\hat{k}$  with positive direction of x, y and z-axis are called direction angles and cosines of these angles are called direction cosines of  $\vec{r}$  usually denoted as  $l = \cos \alpha$ ,  $m = \cos \beta$ ,  $n = \cos \gamma$ Also  $I = \frac{a}{|\vec{r}|}$ ,  $m = \frac{b}{|\vec{r}|}$ ,  $n = \frac{c}{|\vec{r}|}$  and  $l^2 + m^2 + n^2 = 1$ or  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma = 1$
- The numbers a, b, c proportional to I, m, n are called direction ratios.
- Scalar product or dot product of two vectors  $\vec{a}$  and  $\vec{b}$  is denoted as  $\vec{a} \cdot \vec{b}$  and is defined as  $\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$ ,  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b} \cdot (0 \le \theta \le \pi)$ .
- Dot product of two vectors is commutative i.e.  $\vec{a} \cdot \vec{b} = \vec{b} \cdot \vec{a}$
- $\vec{a} \cdot \vec{b} = 0 \iff \vec{a} = \vec{o} \text{ or } \vec{b} = \vec{o} \text{ or } \vec{a} \perp \vec{b}.$
- $\vec{a} \cdot \vec{a} = |\vec{a}|^2$ , so  $\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$
- If  $\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$  and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ , then

$$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3.$$

• Projection of 
$$\vec{a}$$
 on  $\vec{b} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$   
Projection vector of  $\vec{a}$  along  $\vec{b} = \left(\frac{(\vec{a} \cdot \vec{b})}{|\vec{b}|}\right)\hat{b}$ .

- Cross product or vector product of two vectors  $\vec{a}$  and  $\vec{b}$  is denoted as  $\vec{a} \times \vec{b}$  and is defined as  $\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{n}$ . where  $\theta$  is the angle between  $\vec{a}$  and  $\vec{b}$ . ( $0 \le \theta \le \pi$ ). And  $\hat{n}$  is a unit vector perpendicular to both  $\vec{a}$  and  $\vec{b}$  such that  $\vec{a} \cdot \vec{b}$  and  $\hat{n}$  form a right handed system.
- Cross product of two vectors is not commutative i.e.,  $\vec{a} \times \vec{b} \neq \vec{b} \times \vec{a}$ , but  $\vec{a} \times \vec{b} = -(\vec{b} \times \vec{a})$ .

• 
$$\vec{a} \times \vec{b} = \vec{0} \iff \vec{a} = \vec{0}, \vec{b} = \vec{0} \text{ or } \vec{a} \parallel \vec{b}.$$

• 
$$\hat{i} \times \hat{i} = \hat{j} \times \hat{j} = \hat{k} \times \hat{k} = \vec{0}.$$

• 
$$\hat{i} \times \hat{j} = \hat{k}, \hat{j} \times \hat{k} = \hat{i}, \hat{k} \times \hat{i} = \hat{j}$$
 and  $\hat{j} \times \hat{i} = -\hat{k}, \hat{k} \times \hat{j} = -\hat{i}, \hat{i} \times \hat{k} = -\hat{j}$ 

• If 
$$\vec{a} = a_1\hat{i} + a_2\hat{j} + a_3\hat{k}$$
 and  $\vec{b} = b_1\hat{i} + b_2\hat{j} + b_3\hat{k}$ ,then

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$$

• Unit vector perpendicular to both  $\vec{a}$  and  $\vec{b} = \pm \left( \frac{(\vec{a} \times \vec{b})}{|\vec{a} \times \vec{b}|} \right)$ .

- $|\vec{a} \times \vec{b}|$  is the area of parallelogram whose adjacent sides are  $\vec{a}$  and  $\vec{b}$
- $\frac{1}{2} |\vec{a} \times \vec{b}|$  is the area of parallelogram where diagonals are  $\vec{a}$  and  $\vec{b}$ .
- If  $\vec{a}, \vec{b}$  and  $\vec{c}$  form a triangle, then area of the triangle

• 
$$=\frac{1}{2}\left|\vec{a}\times\vec{b}\right| = \frac{1}{2}\left|\vec{b}\times\vec{c}\right| = \frac{1}{2}\left|\vec{c}\times\vec{a}\right|.$$

**Illustration:** Let  $\vec{a} = \hat{i} + 4\hat{j} + 2\hat{k}$ ,  $\vec{b} = 3\hat{i} - 2\hat{j} + 7\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$  Find a vector  $\vec{d}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and  $\vec{c} \cdot \vec{d} = 27$ Solution:

 $\therefore \vec{d}$  is perpendicular to  $\vec{a}$  and  $\vec{b}$  both

Let 
$$\vec{d} = \lambda (\vec{a} \times \vec{b}) = \lambda \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 4 & 2 \\ 3 & -2 & 7 \\ \vec{d} = \lambda (32\hat{i} - \hat{j} - 14\hat{k}) \end{vmatrix}$$

But  $\vec{c} \cdot \vec{d} = 27$  $\therefore (2\hat{i} - \hat{j} + 4\hat{k}) \cdot \lambda (32\hat{i} - \hat{j} - 14\hat{k}) = 27$  $\Rightarrow \lambda (64 + 1 - 56) = 27$  $\Rightarrow \qquad \lambda = 3$ and  $\vec{d} = 3(32\hat{i} - \hat{j} - 14\hat{k}) = 96\hat{i} - 3\hat{j} + 42\hat{k}$ 

### Illustration:

Vectors  $\vec{a}$ ,  $\vec{b}$  and  $\vec{b}$  are such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = 5$ ,  $|\vec{b}| = 7$  and  $|\vec{c}| = 3$ . Find the angle between  $\vec{a}$  and  $\vec{c}$ 

### Solution:

Given 
$$\vec{a} + \vec{b} + \vec{c} = \vec{0}$$
  
 $\vec{a} + \vec{c} = -\vec{b}$   
 $(\vec{a} + \vec{c}) \cdot (\vec{a} + \vec{c}) = (-\vec{b}) \cdot - (\vec{b})$   
 $\Rightarrow (\vec{a})^2 + \vec{a} \cdot \vec{c} + \vec{c} \cdot \vec{a} + (\vec{c})^2 = |\vec{b}|^2$  ( $\because \vec{a} \cdot \vec{a} = |\vec{a}|^2$ )  
 $\Rightarrow 2\vec{a} \cdot \vec{c} = |\vec{b}|^2 - |\vec{a}|^2 - |\vec{c}|^2$   
 $\Rightarrow 2|\vec{a}||\vec{c}|\cos\theta = |\vec{b}|^2 - |\vec{a}|^2 - |\vec{c}|^2$   
Where '\theta' be the angle between  $\vec{a}$  and  $\vec{c}$   
 $\Rightarrow 2 \times 5 \times 3 \cos\theta = 49 - 25 - 9$   
 $\Rightarrow \cos\theta = \frac{15}{30}$   
 $\Rightarrow \cos\theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ 

### Illustration:

 $\Rightarrow \cos \theta = -\frac{1}{2}$ 

 $\Rightarrow \theta = \frac{2\pi}{3}.$ 

Let  $\vec{a}$  and  $\vec{b}$  are two unit vectors and ' $\theta$ ' is the angle between them, then find

' $\theta$ 'if  $\vec{a} + \vec{b}$  is unit vector. Solution: Here  $|\vec{a}| = |\vec{b}| = 1$  and  $|\vec{a} + \vec{b}| = 1$   $\therefore |\vec{a} + \vec{b}|^2 = 1$   $\Rightarrow (\vec{a} + \vec{b}) \cdot (\vec{a} + \vec{b}) = 1$   $(\because \vec{a} \cdot \vec{a} = |\vec{a}|^2)$   $\Rightarrow (\vec{a})^2 + \vec{a} \cdot \vec{b} + \vec{b} \cdot \vec{a} + (\vec{b})^2 = 1$  $\Rightarrow 2 |\vec{a}| |\vec{b}| \cos \theta = -1$ 

### **ONE MARK QUESTIONS**

#### MULTIPLE CHOICE QUESTIONS (1 Mark Each)

Select the correct option out of the four given options:

1. If  $\overrightarrow{AB} = 3\hat{i} + 2\hat{j} - \hat{k}$  and the coordinate of A are (4, 1, 1), then the coordinate of B are.

(a) (1, -1, 2)	(b) (-7, -3, 0)
(c) (7, 3, 0)	(d) (-1, 1, -2)

2. Let  $\vec{a} = -2\hat{i} + \hat{j}$ ,  $\vec{b} = \hat{i} + 2\hat{j}$  and  $\vec{c} = 4\hat{i} + 3\hat{j}$ , then the values of x and y such that  $\vec{c} = x\vec{a} + y\vec{b}$ , are:

(a) 
$$x = 1, y = 2$$
  
(b)  $x = -1, y = 2$   
(c)  $x = -1, y = -2$   
(d)  $x = 1, y = -1$ 

3. A unit vector in the direction of the resultant of the vector  $\hat{i} - \hat{j} + 3\hat{k}$ ,  $2\hat{i} + \hat{j} - 2\hat{k}$  and  $\hat{i} + 2\hat{j} - 2\hat{k}$  is

(a) 
$$\frac{1}{\sqrt{21}}(4\hat{i}-2\hat{j}-\hat{k})$$
  
(b)  $\frac{1}{\sqrt{21}}(4\hat{i}-2\hat{j}+\hat{k})$   
(c)  $4\hat{i}-2\hat{j}-\hat{k}$   
(d)  $\frac{1}{\sqrt{21}}(4\hat{i}+2\hat{j}-\hat{k})$ 

4. If  $2\hat{i} + 3\hat{j} + \hat{k}$  and  $\hat{i} - 2\hat{j} - \hat{k}$  are two vectors, then a vector of magnitude 5 units parallel to the sum of given vectors

(a) 
$$\sqrt{\frac{5}{2}}(3\hat{i} + \hat{j})$$
  
(b)  $\frac{1}{\sqrt{30}}(\hat{i} + 5\hat{j} + 2\hat{k})$   
(c)  $\frac{1}{\sqrt{10}}(3\hat{i} + \hat{j})$   
(d)  $5(3\hat{i} + \hat{j})$ 

5. If 
$$\vec{a} = \lambda \hat{i} + 2\hat{j} + \hat{k}$$
 and  $\vec{b} = 5\hat{i} - 9\hat{j} + 2\hat{k}$  are perpendicular, then the value of ' $\lambda$ ' is

(a) 
$$\lambda = \frac{16}{5}$$
 (b)  $\lambda = -\frac{16}{5}$ 

(c) 
$$\lambda = 4$$
 (d)  $\lambda = \frac{10}{9}$ 

6. The value of p for which  $3\hat{i} + 2\hat{j} + 9\hat{k}$  and  $\hat{i} + p\hat{j} + 3\hat{k}$  are parallel vector is

(a) 
$$p = -\frac{30}{2}$$
 (b)  $p = 15$   
(c)  $p = \frac{2}{3}$  (d)  $p = \frac{3}{2}$ 

7. If  $(2\hat{i} + 6\hat{j} + 27\hat{k}) \times (\hat{i} + 3\hat{j} + p\hat{k}) = \vec{0}$ , then the value of 'p' is

- (a)  $p = -\frac{20}{27}$  (b)  $p = \frac{27}{2}$
- (c) p = 0 (d)  $p = -\frac{27}{2}$

### 8. Value of $\hat{i}.(\hat{j} \times \hat{k}) + (\hat{i} \times \hat{k}).\hat{j}$ is (a) 2 (b) 1 (c) 0 (d) -2

10.	If vector $\lambda \hat{i} + 3\hat{j}$ and $4\hat{i} + \lambda\hat{j}$ are collinea (a) $\lambda = 0$	ar, then the value of ' $\lambda$ ' is (b) $\lambda = 4$
	(c) $\lambda = 3$	(d) $\lambda = \pm 2\sqrt{3}$
11.	A unit vector perpendicular to $2\hat{i} - \hat{j} + 2$	$\hat{k}$ and $4\hat{i}-\hat{j}+3\hat{k}$ is
	(a) $\frac{1}{3}(-\hat{i}+2\hat{j}+2\hat{k})$	(b) $\frac{1}{3}(\hat{i}-2\hat{j}+2\hat{k})$
	(c) $\frac{1}{3}(\hat{i}+2\hat{j}+2\hat{k})$	(d) $\frac{1}{3}(\hat{i}+2\hat{j}-3\hat{k})$
12.	If $\vec{a}$ and $\vec{b}$ are two vectors such that $ \vec{a} $	$ \vec{b}  = \vec{a}.\vec{b}$ , then the angle between $\vec{a}$ and $\vec{b}$ is
	(a) 30°	(b) 45°
	(c) 60°	(d) 90°
13.	If $3i + j - 2k$ and $i - 3j + 4k$ are the dia the parallelogram is	agonals of a parallelagram, then the area of
	(a) 8 sq. units	(b) √91 sq. units
	(c) $5\sqrt{3}$ sq. units	(d) $10\sqrt{3}$ sq. units
14.	If scalar projection of $\lambda \hat{i} + \hat{j} + 4\hat{k}$ on $2\hat{i} + \hat{j}$	$-6\hat{j}+3\hat{k}$ is 4 units, then the value of $\lambda$ is
	(a) $\lambda = 5$	(b) $\lambda = -5$
	(c) $\lambda = -9$	(d) $\lambda = 9$
15.	If $\vec{a} \cdot \vec{b} = 3$ and $ \vec{a} \times \vec{b}  = 3\sqrt{3}$ , then the a	angle between $\vec{a}$ and $\vec{b}$ is
	(a) 30°	(b) 60°
	(c) 120°	(d) 45°
16.	If $ a  = 4$ and $-3 \le k \le 2$ , than the range	e of $ k\bar{a} $ is
	(a) [0, 12] (c) [0, 8]	(b) [8, 12] (d) [-12, 8]
17	$[f \mid \vec{a} \mid -4 \mid \vec{b} \mid -2 \text{ and } \vec{a} \mid \vec{b} \mid -10 \text{ than}$	the value of $ \vec{a}, \vec{b} ^2$ is
17.	(a) = 22	
	(a) 22 (c) 88	(d) None of these
18.	If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , $ \vec{a}  = 3$ , $ \vec{b}  = 4$ and $ \vec{c}  = 4$	$\sqrt{37}$ , then the angle between $ec{a}$ and $ec{b}$ is
	(a) $\frac{\pi}{4}$	(b) $\frac{\pi}{6}$
	(c) $\frac{\pi}{3}$	(d) $\frac{\pi}{2}$

19. If  $(\vec{a} + \vec{b}) \perp \vec{b}$  and  $(\vec{a} + 2\vec{b}) \perp \vec{a}$ , then

(a) $(\vec{a}) = 2   \vec{b}  $	(b) $2 \vec{a}  = \vec{b}$
(c) $(\vec{a}) = (\vec{b})$	(d) $ \vec{a}  = \sqrt{2}  \vec{b} $

20. If 
$$|\vec{a}| = |\vec{b}| = |\vec{a} + \vec{b}| = 1$$
, then the value of  $|\vec{a} - \vec{b}|$  is

(a) 0	(b) 1

(c)  $\sqrt{3}$  (d) 2

#### Assertion-Reason Based Questions

In the following questions a statement of Assertion (A) is followed by a statement of Reason (R) Choose the correct answer out of the following couces:

- (a) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (b) Both (A) and (R) are true, but (R) is not the correct explanation of (A)
- (c) (A) is true and (R) is false
- (d) (A) is false, but (R) is true
- 21. Assertion (A) : If  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$  and  $\vec{a} \cdot \vec{b} = 10$ ,

 $|\vec{a} \times \vec{b}|^2 = 125$ 

Reason (R) :  $|\vec{a} \times \vec{b}|^2 - (\vec{a} \cdot \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2$ 

22. Assertion (A) : If  $\vec{a}$  and  $\vec{b}$  are unit vector such that  $|\vec{a} + \vec{b}| = \sqrt{3}$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is  $\frac{\pi}{2}$ 

Reason (R) : Angle between vectors  $\vec{a}$  and  $\vec{b}$  is given by  $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}||\vec{b}|}$ 

23. Assertion (A) : If  $|\vec{a}| = 4$ ,  $|\vec{b}| = 5$  and  $|\vec{a} \times \vec{b}| = 20$ , then  $\vec{a} \perp \vec{b}$ 

Reason (R): Two non zero vector  $\vec{a}$  and  $\vec{b}$  are perpandicular if  $|\vec{a} \times \vec{b}| = |\vec{a}||\vec{b}|$ 

24. Assertion (A) : If  $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = 12$  and  $\vec{a} = 2 |\vec{b}|$ , then  $|\vec{a}| = 4$  and  $|\vec{b}| = 2$ 

Reason (R): If  $\vec{a}$  and  $\vec{b}$  are two vectors, then  $(\vec{a} + \vec{b}) \cdot (\vec{a} - \vec{b}) = |\vec{a}|^2 - |b|^2$ 

25. Assertion (A) : If  $|2\vec{a} + \vec{b}| = |2\vec{a} - \vec{b}|$ , than  $\vec{a}$  parellel to  $\vec{b}$ 

Reason (B) : Two non zero vector  $\vec{a}$  and  $\vec{b}$  are perpendicular if  $\vec{a} \cdot \vec{b} = 0$ .

### **TWO MARK QUESTIONS**

- 1. A vector  $\vec{r}$  is inclined to x axis at 45° and y-axis at 60° if  $|\vec{r}| = 8$  units. find  $\vec{r}$ .
- 2. if  $|\vec{a} + \vec{b}| = 60$ ,  $|\vec{a} \vec{b}| = 40$  and  $|\vec{b}| = 46$  find  $|\vec{a}|$
- 3. Write the projection of  $\vec{b} + \vec{c}$  on  $\vec{a}$  where

 $\vec{a} = 2\hat{i} - 2\hat{j} + \hat{k}, \vec{b} = \hat{i} + 2\hat{j} - 2\hat{k}$  and  $\vec{c} = 2\hat{i} - \hat{j} + 4\hat{k}$ 

- 4. If the points (-1, -1, 2), (2, m, 5) and (3, 11, 6) are collinear, find the value of *m*.
- 5. For any three vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$  write value of the following.  $\vec{a} \times (\vec{b} + \vec{c}) + \vec{b} \times (\vec{c} + \vec{a}) + \vec{c} \times (\vec{a} + \vec{b})$
- 6. If  $(\vec{a} \times \vec{b})^2 + (\vec{a} \cdot \vec{b})^2 = 144$  and  $|\vec{a}| = 4$ . Find the value of  $|\vec{b}|$ .
- 7. If for any two vectors  $\vec{a}$  and  $\vec{b}$ ,

 $(\vec{a} + \vec{b})^2 + (\vec{a} - \vec{b})^2 = \lambda \left[ (\vec{a})^2 + (\vec{b})^2 \right]$  then write the value of  $\lambda$ .

8. if  $\vec{a}, \vec{b}$  are two vectors such that  $|(\vec{a} + \vec{b})| = |\vec{a}|$  then prove that  $2\vec{a} + \vec{b}$  is perpendicular to  $\vec{b}$ .

9. Show that vectors 
$$\vec{a} = 3\hat{i} - 2\hat{j} + \hat{k}$$

 $\vec{b} = \hat{i} - 3\hat{j} + 5\hat{k}, \vec{c} = 2\hat{i} + \hat{j} - 4\hat{k}$  form a right angle triangle.

- 10. If  $\vec{a}, \vec{b}, \vec{c}$  are three vectors such that  $\vec{a} + \vec{b} + \vec{c} = 0$  and  $|\vec{a}| = 5$ ,  $|\vec{b}| = 12$ ,  $|\vec{c}| = 13$ , then find  $\vec{a}.\vec{b} + \vec{b}.\vec{c} + \vec{c}.\vec{a}$
- 11. The two vectors  $\hat{i} + \hat{j}$  and  $3\hat{i} \hat{j} + 4\hat{k}$  represents the two sides AB and AC respectively of  $\triangle$  ABC, find the length of median through A.

- 12. If position vectors of the points *A*, *B* and *C* are  $\vec{a}$ ,  $\vec{b}$  and  $4\vec{a} 3\vec{b}$  respectively, then find vectors  $\overrightarrow{AC}$  and  $\overrightarrow{BC}$ .
- 13. If position vectors of three points A, B and C are  $-2\vec{a} + 3\vec{b} + 5\vec{c}$ ,  $\vec{a} + 2\vec{b} + 3\vec{c}$  and  $7\vec{a} \vec{c}$  respectively. Then prove that A, B and C are collinear.
- 14. If the vector  $\hat{i} + p\hat{j} + 3\hat{k}$  is rotated through an angle  $\theta$  and is doubled in magnitude, then it becomes  $4\hat{i} + (4p-2)\hat{j} + 2\hat{k}$ . Find the value of *p*.
- 15. If  $\overrightarrow{AB} = 5\hat{i} 2\hat{j} + 4\hat{k}$  and  $\overrightarrow{AC} = 3\hat{i} + 4\hat{k}$  are sides of the triangle *ABC*. Find the length of median through *A*.
- 16. Find scalar projection of the vector  $7\hat{i} + \hat{j} + 4\hat{k}$  on the vector  $2\hat{i} + 6\hat{j} + 3\hat{k}$ . Also find vector porojection
- 17. Let  $\vec{a} = 3\hat{i} + x\hat{j} \hat{k}$  and  $\vec{b} = 2\hat{i} + \hat{j} + y\hat{k}$  are mutually perpendicular and  $|\vec{a}| = |\vec{b}|$ . Find x and y.
- 18. If  $\vec{a}$  and  $\vec{b}$  are unit vectors, find the angle between  $\vec{a}$  and  $\vec{b}$  so that  $\vec{a} \sqrt{2} b$  is a unit vector.
- 19. If  $\vec{a} = 2\hat{i} 2\hat{j} + 3\hat{k}$  and  $\vec{b} = 2\hat{i} + 3\hat{j} 5\hat{k}$ . Find the angle between  $\vec{a}$  and  $\vec{a} \times \vec{b}$ .
- 20. Using vectors, prove that angle in a semi circle is 90°.

### THREE MARKS QUESTIONS

- 1. The points A,B and C with position vectors  $3\hat{i} y\hat{j} + 2\hat{k}$ ,  $5\hat{i} \hat{j} + \hat{k}$  and  $3x\hat{i} + 3\hat{j} \hat{k}$  are collinear. Find the values of x and y and also the ratio in which the point B divides AC.
- 2. If sum of two unit vectors is a unit vector, prove that the magnitude of their difference is  $\sqrt{3}$ .
- 3. Let  $\vec{a} = 4\hat{\imath} + 5\hat{\jmath} \hat{k}$ ,  $\vec{b} = \hat{\imath} 4\hat{\jmath} + 5\hat{k}$  and  $\vec{c} = 3\hat{\imath} + \hat{\jmath} \hat{k}$ . Find a vector  $\vec{d}$  which is perpendicular to both  $\vec{a}$  and  $\vec{b}$  and satisfying  $\vec{d} \cdot \vec{c} = 21$
- 4. If  $\hat{a}$  and  $\hat{b}$  are unit vectors inclined at an angle $\theta$  then proved that
  - (i)  $\cos\frac{\theta}{2} = \frac{1}{2} \left| \hat{a} + \hat{b} \right|$

(ii) 
$$\sin\frac{\theta}{2} = \frac{1}{2}|\hat{a} - \hat{b}|$$
  
(iii)  $\tan\frac{\theta}{2} = \left|\frac{\hat{a} - \hat{b}}{\hat{a} - \hat{b}}\right|$ 

- 5. If  $\vec{a}, \vec{b}, \vec{c}$  are three mutually perpendicular vectors of equal magnitude. Prove that  $\vec{a}+\vec{b}+\vec{c}$  is equally inclined with vectors  $\vec{a}, \vec{b}$  and  $\vec{c}$ . Also find angle.
- 6. For any vector  $\vec{a}$  prove that  $|\vec{a} \times \hat{\iota}|^2 + |\vec{a} \times \hat{j}|^2 + |\vec{a} \times \hat{k}|^2 = 2|\vec{a}|^2$
- 7. Show that  $(\vec{a} \times \vec{b})^2 = |\vec{a}|^2 |\vec{b}|^2 (\vec{a}.\vec{b})^2 = \begin{vmatrix} \vec{a}.\vec{a} & \vec{a}.\vec{b} \\ \vec{a}.\vec{b} & \vec{b}.\vec{b} \end{vmatrix}$
- 8. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are the position vectors of vertices A,B,C of a  $\Delta$  ABC, show that the area of triangle ABC is  $\frac{1}{2} |\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|$ . Deduce the condition for points  $\vec{a}, \vec{b}$  and  $\vec{c}$  to be collinear.

- 9. Let  $\vec{a}, \vec{b}$  and  $\vec{c}$  be unit vectors such that  $\vec{a}, \vec{b} = \vec{a}, \vec{c} = 0$  and the angle between b and c is  $\pi/6$ , prove that  $\vec{a} = \pm 2(\vec{b} \times \vec{c})$ .
- 10. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ , then prove that  $\vec{a} \times \vec{b} = \vec{b} \times \vec{c} = \vec{c} \times \vec{a}$ .
- 11. If  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{c} = \hat{j} \hat{k}$  are given vectors, then find a vector  $\vec{b}$  satisfying the equations  $\vec{a} \times \vec{b} = \vec{c}$  and  $\vec{a} \cdot \vec{b} = 3$ .
- 12. Find the altitude of a parallelepiped determined by the vectors  $\vec{a}, \vec{b} and \vec{c}$  if the base is taken as parallelogram determined by  $\vec{a} and \vec{b}$  and if  $\vec{a} = \hat{i} + \hat{j} + \hat{k}$ ,  $\vec{b} = 2\hat{i} + 4\hat{j} \hat{k}$  and  $\vec{c} = \hat{i} + \hat{j} + 3\hat{k}$ .
- 13. If  $|\vec{a}| = 3$ ,  $|\vec{b}| = 4$  and  $|\vec{c}| = 5$  such that each is perpendicular to sum of the other two, find  $|\vec{a} + \vec{b} + \vec{c}|$
- 14. Decompose the vector  $6\hat{i} 3\hat{j} 6\hat{k}$  in two vectors which are parallel and perpendicular to the vector  $\hat{i} + \hat{j} + \hat{k}$  respectively.
- 15. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are vectors such that  $\vec{a}, \vec{b} = \vec{a}, \vec{c}, \vec{a} \times \vec{b} = \vec{a} \times \vec{c}, a \neq 0$ , then show that  $\vec{b} = \vec{c}$ .
- 16. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three non zero vectors such that  $\vec{a} \times \vec{b} = \vec{c}$  and  $\vec{b} \times \vec{c} = \vec{a}$ . Prove that  $\vec{a}, \vec{b}$  and  $\vec{c}$  are mutually at right angles and  $|\vec{b}| = 1$  and  $|\vec{c}| = |\vec{a}|$
- 17. Simplify  $(\vec{a} \vec{b}) \cdot \{(\vec{b} \vec{c}) \times (\vec{c} \vec{a})\}$

- 18. If  $\vec{r} = x\hat{i} + y\hat{j} + z\hat{k}$ , find the value of  $(\vec{r} \times \hat{i}) \cdot (\vec{r} \times \hat{j}) + xy$
- 19. If  $\vec{a}, \vec{b}$  and  $\vec{c}$  are three vectors such that  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$  and  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$ ,  $|\vec{c}| = 7$ , find the angle between  $\vec{a}$  and  $\vec{b}$ .
- 20. The magnitude of the vector product of the vector  $\hat{i} + \hat{j} + \hat{k}$  with a unit vector along the sum of the vector  $2\hat{i} + 4\hat{j} 5\hat{k}$  and  $\lambda\hat{i} + 2\hat{j} + 3\hat{k}$  is equal to  $\sqrt{2}$ . Find the value of  $\lambda$ .
- 21. If  $\vec{a} \times \vec{b} = \vec{c} \times \vec{d}$  and  $\vec{a} \times \vec{c} = \vec{b} \times \vec{d}$ , prove that  $(\vec{a} \vec{d})$  is parallel to  $(\vec{b} \vec{c})$ , where  $\vec{a} \neq \vec{d}$  and  $\vec{b} \neq \vec{c}$ .
- 22. Find a vector of magnitude  $\sqrt{171}$  which is perpendicular to both of the vectors  $\vec{a} = \hat{i} + 2\hat{j} 3\hat{k}$  and  $\vec{b} = 3\hat{i} \hat{j} + 2\hat{k}$ .
- 23. Prove that the angle betwen two diagonals of a cube is  $\cos^{-1}\left(\frac{1}{3}\right)$ .
- 24. If  $\vec{\alpha} = 3\hat{i} \hat{j}$  and  $\vec{\beta} = 2\hat{i} + \hat{j} + 3\hat{k}$  then express  $\vec{\beta}$  in the form of  $\vec{\beta} = \vec{\beta}_1 + \vec{\beta}_2$ , where  $\vec{\beta}_1$  is parallel to  $\vec{\alpha}$  and  $\vec{\beta}_2$  is perpendicular to  $\vec{\alpha}$ .
- 25. Find a unit vector perpendicular to plane ABC, when position vectors of A,B,C are  $\widehat{3\iota} \hat{j} + 2\hat{k}, \hat{\iota} \hat{j} 3\hat{k}$  and  $4\hat{\iota} 3\hat{j} + \hat{k}$  respectively.
- 26. Find a unit vector in XY plane which makes an angle  $45^{\circ}$  with the vector  $\hat{i} + \hat{j}$  and angle of  $60^{\circ}$  with the vector  $3\hat{i} 4\hat{j}$ .

- 27. Suppose  $\vec{a} = \lambda \hat{i} 7\hat{j} + 3\hat{k}$ ,  $\vec{b} = \lambda \hat{i} + \hat{j} + 2\lambda \hat{k}$ . If the angle between  $\vec{a}$  and  $\vec{b}$  is greater than 90°, then prove that  $\lambda$  satisfies the inequality-7 <  $\lambda$  < 1.
- 28. If  $\vec{a}$  and  $\vec{b}$  are two unit vectors such that  $|\vec{a} + \vec{b}| = \sqrt{3}$  then find the value of  $(2\vec{a} 5\vec{b})$ .  $(3\vec{a} + \vec{b})$ .
- 29. Let  $\vec{a} = 2\hat{i} + \hat{j} 3\hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j} + \hat{k}$  and  $\vec{c} = \hat{i} \hat{j} + \hat{k}$ . Find a vector  $\vec{d}$  such that  $\vec{a} \cdot \vec{d} = 0$ ,  $\vec{b} \cdot \vec{d} = 2$  and  $\vec{c} \cdot \vec{d} = 4$ .

### **Case Study Questions (4 Marks Each)**

1. A farmer moves along the boundary of a triangular field PQR. Three vertices of the triangular field are P(2, 1, -2), Q(-1, 2, 1) and R(1, -4, -2) respectively.



On the basis of above information, answer the following questions:

- (i) Find the length of PQ.
- (ii) Find the  $\angle PQR$
- (iii) Find the area of the  $\triangle PQR$

OR

(iii) Find projection of QP on QR.

### **SELF ASSESSMENT-1**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECTION CHOOSE THE CORRECT OPTION.

1. A unit vector perpendicular to both  $\hat{i} + \hat{j}$  and  $\hat{j} + \hat{k}$  is

(A) 
$$\hat{i} + \hat{j} + \hat{k}$$
 (B)  $\hat{i} - \hat{j} + \hat{k}$   
(C)  $\frac{1}{\sqrt{3}} (\hat{i} - \hat{j} + \hat{k})$  (D)  $\frac{1}{\sqrt{3}} (\hat{i} + \hat{j} + \hat{k})$ 

2. If 
$$|\vec{a} \cdot \vec{b}| = 2$$
,  $|\vec{a} \times \vec{b}| = 4$ , then the value of  $|\vec{a}|^2 |\vec{b}|^2$  is  
(A) 2 (B) 6  
(C) 8 (D) 20

3. The projection of vector  $\vec{a} = \hat{i} - 2\hat{j} + \hat{k}$  on vector  $\vec{b} = 4\hat{i} - 4\hat{j} + 7\hat{k}$  is (A)  $\frac{9}{19}$  (B)  $\frac{9}{\sqrt{19}}$ (C)  $\frac{9}{\sqrt{6}}$  (D)  $\frac{19}{9}$ 

4. If  $\vec{a}$  is any vector, then the value of  $(\vec{a} \times \hat{i})^2 + (\vec{a} \times \hat{j})^2 + (\vec{a} \times \hat{k})^2$  is (A)  $|\vec{a}|^2$ (B)  $2|\vec{a}|^2$ (B)  $2|\vec{a}|^2$ (D)  $4|\vec{a}|^2$ 

5. If  $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ ,  $|\vec{a}| = 3$ ,  $|\vec{b}| = 5$ ,  $|\vec{c}| = 7$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is

(A) 
$$\frac{\pi}{6}$$
 (B)  $\frac{\pi}{3}$   
(C)  $\frac{2\pi}{3}$  (D)  $\frac{5\pi}{3}$ 

### **SELF ASSESSMENT-2**

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECTION CHOOSE THE CORRECT OPTION.

- 1. If  $\vec{a}$ ,  $\vec{b}$  and  $\vec{a} + \vec{b}$  are unit vectors. Then the value of  $|\vec{a} \vec{b}|$  is (A) 0 (B) 1 (C)  $\sqrt{2}$  (D)  $\sqrt{3}$
- 2. If  $\vec{a}$  and  $\vec{b}$  are two vectors such that  $|\vec{a}| = 2$ ,  $|\vec{b}| = 1$  and  $\vec{a} \cdot \vec{b} = 1$ , then the value of  $(3\vec{a} 5\vec{b}) \cdot (2\vec{a} + 7\vec{b})$  is (A) 0 (B) 41 (C) 29 (D) 7
- 3. If  $\vec{c} \cdot (\hat{i} + \hat{j}) = 2$ ,  $\vec{c} \cdot (\hat{i} \hat{j}) = 3$  and  $\vec{c} \cdot \hat{k} = 0$ , then the vector  $\vec{c}$  is (A)  $\frac{1}{2}(5\hat{i} + \hat{j})$  (B)  $\frac{1}{2}(5\hat{i} - \hat{j})$ (C)  $\frac{1}{2}(\hat{i} - 5\hat{j})$  (D)  $\frac{1}{2}(\hat{i} + 5\hat{j})$
- 4. If the projection of  $3\hat{i} + \lambda\hat{j} + \hat{k}$  on  $\hat{i} + \hat{j}$  is  $\sqrt{2}$  units, then the value  $\lambda$  is (A) 1 (B) -1 (C) 0 (D) 2
- 5. If  $|\vec{a}| = 2$ ,  $|\vec{b}| = 7$  and  $\vec{a} \times \vec{b} = 3\hat{i} + 2\hat{j} 6\hat{k}$ , then the angle between  $\vec{a}$  and  $\vec{b}$  is (A)  $\frac{\pi}{6}$  (B)  $\frac{\pi}{4}$  (C)  $\frac{\pi}{3}$  (D)  $\frac{\pi}{2}$

### <u>Answers</u>

### **ONE MARK QUESTIONS**

### MCQ (1 Mark Each)

1.	(c) (7, 3, 0)
2.	(b) x = -1, y = 2
3.	(d) $\frac{1}{\sqrt{21}}(4\hat{i}+2\hat{j}-\hat{k})$
4.	(a) $\sqrt{\frac{5}{2}}(3\hat{i}+\hat{j})$
5.	(a) $\lambda = \frac{16}{5}$
6.	(c) 2/3
7.	(b) $p = \frac{27}{2}$
8.	(c) 0
9.	(a) −5
10.	(d) $\lambda = \pm 2\sqrt{3}$
<b>1</b> 1.	(a) $\frac{1}{3}(-\hat{i}+2\hat{j}+2\hat{k})$
12.	(b) 45°
13.	(c) $5\sqrt{3}$ sq. units
14	(a) $\lambda = 5$
15.	(b) 60°
16.	(a) [0, 12]
17.	(b) 44
18.	(c) $\frac{\pi}{3}$
19.	(d) $ \vec{a}  = \sqrt{2}  \vec{b} $
20.	(c) $\sqrt{3}$
21.	(C)
22.	(b)
23.	(a)
24.	(a)
25.	(d)
# **TWO MARK QUESTIONS**

1.  $4(\sqrt{2i} + j + k)$ 2. 22 3. 2 4. m = 85. 0 6. 3 7.  $\lambda = 2$ 10. -169 11.  $2\sqrt{2}$ 12.  $\overrightarrow{AC} = 3(\overrightarrow{a} - \overrightarrow{b}), \overrightarrow{BC} = 4(\overrightarrow{a} - \overrightarrow{b})$ 14.  $p = -\frac{2}{3}, 2$ 15.  $\sqrt{33}$ 16.  $\frac{32}{7}, \frac{32}{49}(2i + 6j + 3k)$ 17.  $x = -\frac{31}{12}, y = \frac{41}{12}$ 18.  $\frac{\pi}{4}$ 19.  $\frac{\pi}{2}$ 

# THREE MARKS QUESTIONS

1.	x = 3, y = 3, 1:2	26. $\frac{13}{\sqrt{170}}\hat{i} + \frac{1}{\sqrt{170}}\hat{j}$
3.	$\vec{d} = 7\hat{\imath} - 7\hat{\jmath} - 7\hat{k}$	28. $-\frac{11}{2}$
5.	$\cos^{-1}\frac{1}{\sqrt{3}}$	$\vec{d} = 2\hat{i} - \hat{j} + \hat{k}$
8.	$\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a} = \vec{0}$	Case Study Questions
11.	$\vec{b} = \frac{5}{3}\hat{\imath} + \frac{2}{3}\hat{\jmath} + \frac{2}{3}\hat{k}$	(i) $\sqrt{19}$ units
12.	$\frac{4}{\sqrt{38}}$ units	(ii) $\cos^{-1}\left(\frac{3}{\sqrt{19}}\right)$
13.	$5\sqrt{2}$	(iii) $\frac{7}{2}\sqrt{10}$ square units
14.	$\left(-\hat{\iota}-\hat{j}-\hat{k}\right)+\left(7\hat{\iota}-2\hat{j}-5\hat{k}\right)$	OR (iii) 3 units
17.	0	
18.	0	<b>SELF ASSESSMENT-1</b> 1. (C) 2. (D)
19.	60°	3. (D) 4. (B) 5. (B)
20.	$\lambda = 1$	SELF ASSESSMENT-2
22.	$\hat{i} - 11\hat{j} - 7\hat{k}$	1. (D) 2. (A) 3. (B) 4. (B) 5. (A)
24.	$\vec{\beta} = \left(\frac{3}{2}\hat{\imath} - \frac{1}{2}\hat{\jmath}\right) + \left(\frac{1}{2}\hat{\imath} + \frac{3}{2}\hat{\jmath} + 3\hat{k}\right)$	
25.	$\frac{-1}{\sqrt{165}} \left( 10\hat{\imath} + 7\hat{\jmath} - 4\hat{k} \right)$	

# CHAPTER 11

# THREE-DIMENSIONAL GEOMETRY

In the real world, everything you see is in a threedimensional shape, it has length, breadth, and height. Just simply look around and observe. Even a thin sheet of paper has some thickness.





Applications of geometry in the real world include the computer-aided design (CAD) for construction blueprints, the design of assembly systems in manufacturing such as automobiles, nanotechnology, computer graphics, visual graphs, video game programming, and virtual reality creation.

The next time you play a mobile game, thank three-dimension geometry for the realistic look to the landscape and the characters that exhibit the game's virtual world.



Topics to be covered as per C.B.S.E. revised syllabus (2024-25)

- Direction cosines and direction ratios of a line joining two points.
- Cartesian equation and vector equation of a line.
- Skew lines
- Shortest distance between two lines.
- Angle between two lines.

• **Distance Formula:** Distance (d) between two points( $x_1$ ,  $y_1$ ,  $z_1$ )and( $x_2$ ,  $y_2$ ,  $z_2$ )

$$d = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2 + (z_2 - z_1)^2}$$

• Section Formula: line segment AB is divided by P (x, y, z) in ratio m:n

(a) Internally	(b) Externally		
$\left(\frac{m x_2 + n x_1}{m + n}, \frac{m y_2 + n y_1}{m + n}, \frac{m z_2 + n z_1}{m + n}\right)$	$\left(\frac{mx_2 - nx_1}{m - n}, \frac{my_2 - ny_1}{m - n}, \frac{mz_2 - nz_1}{m - n}\right)$		

- Direction ratio of a line through  $(x_1, y_1, z_1)$  and  $(x_2, y_2, z_2)$  are  $x_2 x_1, y_2 y_1, z_2 z_1$
- **Direction cosines** of a line having direction ratios as a, b, c are:

$$l = \pm \frac{a}{\sqrt{a^2 + b^2 + c^2}}$$
,  $m = \pm \frac{b}{\sqrt{a^2 + b^2 + c^2}}$ ,  $n = \pm \frac{c}{\sqrt{a^2 + b^2 + c^2}}$ 

• Equation of line in space:

	Vector form		Cartesian form
(i)	Passing through point $ec{a}$ and	(i)	Passing through point
	parallel to vector $\vec{b}$ ; $\vec{r} = \vec{a} + \lambda \vec{b}$		$(x_1, y_1, z_1)$ and having direction ratios a, b, c;

				$\frac{x-x_1}{a}$	$=\frac{y-y_1}{b}=$	$=\frac{z-z_1}{c}$	
(ii)Passing	through	two	points	(ii) Passing	through	two	points
$\vec{a}$ and $\vec{b}$ ; $\vec{r} = \vec{a} + \lambda \ (\vec{b} - \vec{a})$			$(x_1, y_1z_1)$ and $(x_2, y_2z_2)$ ;				
				$\frac{x-x_1}{x_2-x}$	$\frac{1}{x_1} = \frac{y - y_1}{y_2 - y_1}$	$=\frac{z-z}{z_2-z}$	$\frac{z_1}{z_1}$

### • Angle between two lines:

Vector form	Cartesian form
(i) For lines $\vec{r} = \vec{a_1} + \lambda \vec{b_1}$ and	(ii) For lines $\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1}$ and
$\vec{r} = \vec{a_2} + \mu \vec{b_2}$ , $\cos\theta = \frac{ \vec{b_1} \cdot \vec{b_2} }{ \vec{b_1}  \vec{b_2} }$	$\frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$
where ' $\theta$ ' is the angle	$\cos \theta$
between two lines.	$= \frac{ a_1a_2 + b_1b_2 + c_1c_2 }{\sqrt{a_1^2 + b_1^2 + c_1^2}\sqrt{a_2^2 + b_2^2 + c_2^2}}$
(iii) Lines are perpendicular if $\vec{b}_1 \cdot \vec{b}_2 = 0$	(ii) Lines are perpendicular if $a_1a_2 + b_1b_2 + c_1c_2 = 0$
(iv) Lines are parallel if $\vec{b}_1 = k\vec{b}_2$ ;	(i) Lines are parallel if
$k \neq 0$	$\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

### Shortest distance between two skew lines

The shortest distance between  
two skew lines  

$$\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$$
 and  $\vec{r} = \vec{a}_2 + \mu \vec{b}_2$  is  

$$d = \left| \frac{(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)}{|\vec{b}_1 \times \vec{b}_2|} \right|$$
If  $d = 0$ , lines are intersecting  
If  $d = 0$ , lines are intersecting  

$$D = \{(a_1b_2 - a_2b_1)^2 + (b_1c_2 - b_2c_1)^2 + (c_1a_2 - c_2a_1)^2\}$$

### • Shortest distance between two parallel lines

Let  $\vec{r} = \vec{a}_1 + \lambda \vec{b}$  and  $\vec{r} = \vec{a}_2 + \mu \vec{b}$  are parallel lines then shortest distance between those lines  $d = \left| \frac{\vec{b} \times (\vec{a}_2 - \vec{a}_1)}{\vec{a}_1} \right|$  units

$$(b)$$
 If  $d = 0$ , then lines coincident.

Illustration 1:

Are the following lines interesting?

$$\vec{r} = 3\hat{i} + 2\hat{j} - 4\hat{k} + \lambda(\hat{i} + 2\hat{j} + 2\hat{k})$$
  
and  $\vec{r} = 5\hat{i} - 2\hat{j} + \mu(3\hat{i} + 2\hat{j} + 6\hat{k})$ 

If yes, find point of intersection. **Solution:** 

We can write the equations in cartesian form

and 
$$\frac{x-3}{1} = \frac{y-2}{2} = \frac{z+4}{2} = 1$$
 ...(*i*)  
 $\frac{x-5}{3} = \frac{y+2}{2} = \frac{z}{6} = m$  ...(*ii*)

Any point on line (i)  $P(\lambda + 3, 2\lambda + 2, 2\lambda - 4)$ Any point on line (ii)  $Q(3\mu + 5, 2\mu - 2, 6\mu)$ Comparing *x*, *y* and *z* coordinate respectively  $\lambda + 3 = 3\mu + 5, 2\lambda + 2 = 2\mu - 2, 2\lambda - 4 = 6\mu$ or  $\lambda - 3\mu = 2, 2\lambda - 2\mu = -4, 2\lambda - 6\mu = 4$ or  $\lambda - 3\mu = 2, \lambda - \mu = -2, \lambda - 3\mu = 2$ Solving first two, we get  $\lambda = -4, \mu = -2$  $\therefore \lambda = -4, \mu = -2$ , Satisfies  $\lambda - 3\mu = 2$  $\therefore$  lines are intersecting and point of interesting (-1, -6, -12) Or Using distance formula If  $tr(\vec{b}_1 \times \vec{b}_2) \cdot (\vec{a}_2 - \vec{a}_1) = 0$ 

Illustration 2:

Find the foot of perpendicular from the point P(1, 2, -3) to the line  $\frac{x+1}{2} = \frac{y-3}{-2} = \frac{z}{-1}$ . Also find the length of the perpendicular and image of P in the given lines. Solution: We have  $\frac{x+1}{2} = \frac{y-3}{-2} = \frac{z}{-1} = 1$ (say)  $\therefore x = 2\lambda - 1, y = -2\lambda + 3, z = -\lambda$ P(1, 2, -3)Let M( $2\lambda - 1$ ,  $-2\lambda + 3$ ,  $-\lambda$ ) be the foot of perpendicular. DR's of PM are  $< 2\lambda - 1 - 1$ ,  $-2\lambda + 3 - 2$ ,  $-\lambda + 3 >$ or  $< 2\lambda - 2, -2\lambda + 1, -\lambda + 3 >$ Line ΜΓ : PM is perpendicular to the line  $\therefore 2(2\lambda - 2) - 2(-2\lambda + 1) - 1(-\lambda + 3) = 0$  $4\lambda - 4 + 4\lambda - 2 + \lambda - 3 = 0$ Q(a, b, c)  $9\lambda - 9 = 0$  $\Rightarrow \lambda = 1$ 

[Class XII : Maths]

 $\therefore \text{ Foot of the perpendicular M} = (1, 1 - 1)$ and PM =  $\sqrt{(1-1)^2 + (2-1)^2 + (-3+1)^2} = \sqrt{0+1+4} = \sqrt{5}$ Let Q(a, b, c) be the image of P As M be the mid point of PQ. (As line is plane mirror)  $\therefore \quad \frac{a+1}{2} = 1 \quad P \quad a = 1$  $\frac{b+2}{2} = 1 \quad P \quad b = 0$  $\frac{c-3}{2} = -1 \quad P \quad c = 1$  $\therefore \text{ image of P is (1, 0, 1)}$ 

### **ONE MARK QUESTIONS**

Multiple Choice Questions (1 Mark Each)

Select the correct option out of the four given options:

1. Distance of the point (a, b, c) from x-axis is

	(a) $\sqrt{b^2 + c^2}$	(b) $\sqrt{c^2 + a^2}$
	(c) $\sqrt{a^2 + b^2}$	(d) $\sqrt{a^2 + b^2 + c^2}$
2.	Angle between the lines $2x = 3y = -z$ a	and $6x = -y = -4z$ is
	(a) 45°	(b) 60°
	(c) 90°	(d) 30°
3.	Equation of the line passing through (2,	-3, 5) and parallel to
	$\frac{x-1}{3} = \frac{y-2}{4} = \frac{z+1}{-1}$ is	
	(a) $\frac{x+2}{3} = \frac{y-3}{4} = \frac{z+5}{-1}$	(b) $\frac{x-2}{3} = \frac{y-3}{4} = \frac{z-5}{1}$
	(c) $\frac{x-2}{3} = \frac{y+3}{4} = \frac{5-z}{1}$	(d) $\frac{x-2}{-3} = \frac{y+3}{-4} = \frac{z-5}{2}$
4.	If the lines $\frac{x-1}{2} = \frac{z-3}{5} = \frac{z-1}{\lambda}$ and $\frac{z-2}{3}$	$\frac{2}{z} = \frac{y+1}{-2} = \frac{z}{2}$ are perpendicular, then the value
	of 'λ' is	
	(a) $\lambda = -2$	(b) $\lambda = 2$
	(c) $\lambda = 1$	(d) $\lambda = -1$

Cartesian form of line  $\vec{r} = (\hat{i} - \hat{j}) + \lambda(2\hat{j} - \hat{k})$  is 5. (a)  $\frac{x-1}{0} = \frac{y+1}{2} = \frac{z}{-1}$ (b)  $\frac{x+1}{1} = \frac{y-1}{2} = \frac{z}{-1}$ (d)  $\frac{x+1}{2} = \frac{y+1}{1} = \frac{z}{0}$ (c)  $\frac{x-1}{2} = \frac{y-1}{-1} = \frac{z}{0}$ The coordinates of the foot of the parpendicular drwan from the point (-2, 87) on the 6. xz plane is (a) (0, 8, 0) (b) (-2, 0, 7) (c) (2, 8, -7) (d) (-2, -8, 7) The length of perpendicular from the point (4, -7, 3) on the y-axis is 7. (b) 4 units (a) 3 units (c) 5 units (d) 7 units If  $\cos\alpha$ ,  $\cos\beta$  and  $\cos\gamma$  are direction cosimes of a line, then the value of  $\cos 2\alpha$  + 8.  $\cos 2\beta + \cos 2\gamma$  is (a) 1 (b) -1 (d) -2(c) 2 9. If two lines x = ay + b, z = cy + d and x = a'y + b, z = c'y + d are perpendicular, then (a) aa' + cc' = 1 (b) aa' + cc' + 1 = 0 (c)  $\frac{a}{a'} + \frac{c}{c'} = 1$ (d)  $\frac{a}{a'} + \frac{c}{c'} + 1 = 0$ 10. A point P lines on the line segment joining the points (-1, 3, 2) and (5, 0, 6), if xcoordinate of P is 2, then its z coordinate is (a) 8 (b) 4 (c) 3 (d) -1 ASSERTION-REASON BASED QUESTIONS In the following questions a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices: (a) Both (A) and (R) are true and (R) is the correct explanation of (A) (b) Both (A) and (R) are true but (R) is not the correct explanation of (A) (c) (A) is true and (R) is false (d) (A) is false, but (R) is true 11. Assertion (A) : The vector equation of a line passing through the points (3, 1, 2) and (4, 2, 5) is  $\vec{r} = 3\hat{i} + \hat{j} + 2\hat{k} + \lambda(\hat{i} + \hat{j} + 3\hat{k})$ ) Reason (R) : The vector equation of a line passing through the points with position vector  $\vec{a}$  and  $\vec{b}$  is  $\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$ 

Assertion (A) : If a line joining the points (1, 0, 4) and (3, λ, 7) is perpendicular to the line joining the points (1, 2, -1) and (2, 3, 0), then λ = -5

Reason (R) : Two lines with direction ratios  $(a_1, b_1, c_1)$  and  $\langle a_2, b_2, c_2 \rangle$  are parallel if

 $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$ 

13. Assertion (A) : The coordinates of the point where the line

$$\vec{r} = (3\hat{i} + \hat{j} - \hat{k}) + \lambda(-\hat{i} + 2\hat{j} + 3\hat{k})$$
 cuts xy-plane and  $\left(\frac{8}{3}, \frac{-5}{3}, 0\right)$ 

Reason (R): The z-coordinate of any point on xy-plane is 0.

14. Assertion (A) : Lines  $\frac{x+1}{-1} = \frac{2-y}{-2} = \frac{z-3}{3}$  and  $\frac{2-x}{-3} = \frac{y-1}{4} = \frac{z+2}{-1}$  intersect at a point.

Reason (R) : Two lines  $\vec{r} = \vec{a}_1 + \lambda \vec{b}_1$  and  $\vec{r} = \vec{a}_2 + \mu \vec{b}_2$  are intersecting if  $(\vec{b}_1 \times \vec{b}_2).(\vec{a}_2 - \vec{a}_1) \neq 0$ .

### **TWO MARKS QUESTIONS**

- 1. Find the equation of a line passing though (2, 0, 5) and which is parallel to line 6x 2 = 3y + 1 = 2z 2
- 2. The equation of a line are 5x 3 = 15y + 7 = 3 10 z. Write the direction cosines of the line
- 3. If a line makes angle  $\alpha$ ,  $\beta$ ,  $\gamma$  with Co-ordinate axis then what is the value of  $\sin^2 \alpha + \sin^2 \beta + \sin^2 \gamma$
- 4. Find the equation of a line passing through the point (2, 0, 1) and parallel to the line whose equation is  $\vec{r} = (2\lambda + 3)\hat{i} + (7\lambda 1)\hat{j} + (-3\lambda + 2)\hat{k}$
- 5. Find the condition that the lines x = ay + b, z = cy + d and x = a'y + b', z = c'y + d'may be perpendicular to each other.
- 6. Show that the lines x = -y = 2z and x + 2 = 2y 1 = -z + 1 are perpendicular to each other.

- 7. Find the equation of the line through (2, 1, 3) and parallel to the line  $\frac{2x-1}{2} = \frac{4-y}{7} = \frac{z+1}{2}$  in cartesian and vector form.
- 8. Find the cartesian and vector equation of the line through the points (2, -3, 1)and (3, -4, -5)
- 9. For what value of  $\lambda$  and  $\mu$  the line joining the points (7,  $\lambda$ , 2), ( $\mu$ , -2, 5) is parallel to the line joining the points (2, -3, 5), (-6, -15, 11)?
- 10. If the points (-1, 3, 2), (-4, 2, -2) and  $(5, 5, \lambda)$  are Collinear, find the value of  $\lambda$ .

# **THREE/FIVE MARKS QUESTIONS**

- 1. Find vector and Cartesian equation of a line passing through a point with position vector  $2\hat{i} - \hat{j} + \hat{k}$  and which is parallel to the line joining the points with position vectors  $-\hat{i} + 4\hat{j} + \hat{k}$  and  $\hat{i} + 2\hat{j} + 2\hat{k}$ .
- 2. Find image (reflection) of the point (7, 4, -3) in the line  $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$ .

3. Show that the lines  $\lim \frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$  and  $\lim \frac{x-2}{1} = \frac{y-4}{3} = \frac{z-6}{5}$  intersect each other. Find the point of intersection.

4. Find the shortest distance between the lines:

$$\overline{r} = \hat{\iota} + 2\hat{\jmath} + 3\hat{k} + \mu(2\hat{\iota} + 3\hat{\jmath} + 4\hat{k})$$
 and

$$\overline{r} = (2\hat{\imath} + 4\hat{\jmath} + 5\hat{k}) + \lambda(3\hat{\imath} + 4\hat{\jmath} + 5\hat{k}).$$

5. Find shortest distance between the lines:

$$\frac{x+1}{7} = \frac{y+1}{-6} = \frac{z+1}{1} \text{ and } \frac{x-3}{1} = \frac{5-y}{2} = \frac{z-7}{1}$$

6. Find the shortest distance between the lines:

$$\overline{r} = (1 - \lambda)\hat{\imath} + (\lambda - 2)\hat{\jmath} + (3 - 2\lambda)\hat{k}$$
$$\overline{r} = (\mu + 1)\hat{\imath} + (2\mu - 1)\hat{\jmath} - (2\mu + 1)\hat{k}$$

- 7. Find the foot of perpendicular from the point  $2\hat{i} \hat{j} + 5\hat{k}$  on the line  $\overline{r} = (11\hat{i} 2\hat{j} 8\hat{k}) + \lambda(10\hat{i} 4\hat{j} 11\hat{k})$ . Also find the length of the perpendicular.
- 8. A line makes angles  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$  with the four diagonal of a cube. Prove that  $\cos^2 \alpha + \cos^2 \beta + \cos^2 \gamma + \cos^2 \delta = \frac{4}{3}$
- 9. Find the length and the equations of the line of shortest distance between the lines  $\frac{x-8}{3} = \frac{y+9}{-16} = \frac{z-10}{7}$  and  $\frac{x-15}{3} = \frac{y-29}{8} = \frac{z-5}{-5}$ .
- 10. Show that  $\frac{x-1}{2} = \frac{y+1}{3} = z$  and  $\frac{x+1}{5} = \frac{y-2}{2}$ , z = 2. do not intersect each other.
- 11. If the line  $\frac{x+2}{2} = \frac{y+1}{3} = \frac{z-1}{4}$  and  $\frac{x-3}{1} = \frac{y-k}{2} = \frac{z}{1}$  intersect, then find the value of *k*.
- 12. Find the equation of the line which intersects the lines  $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{4}$ and  $\frac{x+2}{1} = \frac{y-3}{2} = \frac{z+1}{4}$  and passes through the point (1, 1, 1).
- 13. Find the equations of the two lines through the origin which intersect the line  $\frac{x-3}{2} = \frac{y-3}{1} = \frac{z}{1}$  at angle of  $\pi/3$ .

14. Find the foot of perpendicular drawn from the point (2, -1, 5) to the line  $\vec{r} = (11\hat{i} - 2\hat{j} - 8\hat{k}) + \lambda(10\hat{i} - 4\hat{j} - 11\hat{k})$ 

Also find the length of the perpendicular. Hence find the image of the point (2, -1, 5) in the given line.

- 15. Find the image of the point P(2, -1, 11) in the line  $\vec{r} = (2\hat{i} + 3\hat{k}) + \lambda(2\hat{i} + 3\hat{j} + 4\hat{k})$
- 16. Find the point(s) on the line through the point P(3, 5, 9) and Q(1, 2, 3) at a distance 14 units from the mid-point of segment PQ.
- 17. Find the shortest distance between the following pair of lines

 $\frac{x-1}{2} = \frac{y+1}{3} = z \text{ and } \frac{x+1}{5} = \frac{y-2}{1}; z = 2$ 

Hence write whether the lines are intersecting or not.

18. Find the foot of perpendicular from the point (1, 2, 3) to the line  $\vec{r} = (6\hat{i} + 7\hat{j} + 7\hat{k}) + \lambda(3\hat{i} + 2\hat{j} - 2\hat{k})$ 

Also find the equation of the perpendicular and length of perpendicular.

- 19. Find the equation of the line passing through (-1, 3, -2) and perpendicular to the lines  $\frac{x+1}{1} = \frac{y-2}{2} = \frac{z+5}{3}$  and  $\frac{x-2}{-3} = \frac{y}{2} = \frac{z+1}{5}$
- 20. Find the points on the line  $\frac{x+2}{3} = \frac{y+1}{2} = \frac{3-z}{-2}$  at a distance  $3\sqrt{2}$  from the point (1, 2, 3)
- 21. The points P(4, 5, 10), Q(2, 3, 4) and R(1, 2, -1) are three vertices of a parallelogram PQRS. Find the vector equations of the sides PQ and QR and also find the coordinates of point R.
- 22. Find the equation of perpendicular from the point (3, -1, 11) to the line

$$\frac{x}{2} = \frac{2y-4}{6} = \frac{3-z}{-4}.$$

Also find the foot of the perpendicular and the length of the perpendicular.

23. Show that the lines  $\frac{1-x}{-2} = \frac{y-2}{3} = \frac{z-3}{4}$  and  $\frac{x-4}{5} = \frac{1+y}{2} = z$  are intersecting. Also find the point of intersection.

24. For what value of ' $\lambda$ ', the following are Skew lines?  $\frac{x-4}{5} = \frac{1+y}{2} = z, \ \frac{x-1}{2} = \frac{y-2}{3} = \frac{z-\lambda}{4}$ 

25. |Find the vector equation of the line passing through (2, 1, -1) and parallel to the line  $\vec{r} = (\hat{i} + \hat{j}) + \lambda(2\hat{i} - \hat{j} + \hat{k})$ . Also find the distance between these two lines.

#### **SELF ASSESSMENT-1**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. The foot of perpendicular drawn from the point (2, -1, 5) to the line  $\frac{x-11}{10} = \frac{y+2}{-4} = \frac{z+8}{-11}$  is

(a)	(2, 1, 3)	(b)	(3, 1, 2)
(c)	(1,2,3)	(d)	(3, 2, 1)

2. The shortest distance between the lines  $\vec{r} = (6\hat{i} + 2\hat{j} + 2\hat{k}) + \lambda(\hat{i} - 2\hat{j} + 2\hat{k})$ 

and 
$$\vec{k} = (-4\hat{i} - 4\hat{k}) + \mu(3\hat{i} - 2\hat{j} - 2\hat{k})$$
 is

- (a) 10 units (b) 9 units
- (c) 12 units (d) 9/2 units
- 3. If the x-coordinate of a point A on the join of B(2, 2, 1) and C(5, 1, −2) is 4 then its z-coordinate is

(a)	+2	(b)	-1
(c)	1	(d)	2

4. The distance of the point M(a, b, c) from the x-axis is

(a)	$\sqrt{b^2 + c^2}$	(b)	$\sqrt{c^2 + a^2}$
©	$\sqrt{a^2+b^2}$	(d)	$\sqrt{a^2-b^2+c^2}$

5. The straight line  $\frac{x-3}{3} = \frac{y-2}{1} = \frac{z-1}{0}$  is

(c) parallel to z-axis (d) perpendicular to z-axis

[Class XII : Maths]

#### **SELF ASSESSMENT-2**

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. The shortest distance between the line  $\frac{x-3}{3} = \frac{y}{0} = \frac{z}{-4}$  and y-axis is (a)  $\frac{12}{5}$  units (b)  $\frac{1}{5}$  units
  - (c) 0 units (d) 3 units
- 2. The point of intersection of the lines  $\frac{x+1}{3} = \frac{y+3}{5} = \frac{z+5}{7}$ and  $\frac{x-2}{1} = \frac{y-4}{3} = \frac{z-6}{5}$  is (a)  $\left(\frac{1}{3}, \frac{-1}{3}, -\frac{2}{3}\right)$  (b)  $\left(\frac{1}{3}, \frac{1}{3}, \frac{1}{3}\right)$ (c)  $\left(\frac{1}{2}, \frac{1}{2}, \frac{1}{2}\right)$  (d)  $\left(\frac{1}{2}, \frac{-1}{2}, \frac{-3}{2}\right)$
- 3. If a line makes the same angle  $\alpha$ , with each of the *x* and *z* axes and the angle  $\beta$  with y-axis such that  $3\sin^2\alpha = \sin^2\beta$ , then the value of  $\cos^2\alpha$  is
- (a)  $\frac{1}{5}$ (b)  $\frac{2}{5}$ (c)  $\frac{3}{5}$ (d)  $\frac{2}{3}$ 4. If the lines  $\frac{x+3}{k-5} = \frac{y-1}{1} = \frac{5-z}{-2k-1}$  and  $\frac{x+2}{-1} = \frac{2-y}{-k} = \frac{z}{5}$  are perpendicular, then the value of k is (a) 1 (b) -1 (c) 2 (d) -2

5. The image of the point P(-1, 8, 4) to the line  $\frac{x}{2} = \frac{y+1}{-2} = \frac{z-3}{-4}$  is (a) (5, 4, 4) (b) (5, 0, 4)

(a)	(5, 4, 4)	(b)	(5, 0, 4)
(c)	(-3, -6, 10)	(d)	(1, 8, 4)

# **Case Study Based Questions**

1. Two birds are flying in the space along straight path  $L_1$  and  $L_2$ 

(Neither parallel nor intersecting) where,



$$L_1: \frac{3}{3} = \frac{-1}{-1} = \frac{-1}{1}$$
$$L_2: \frac{x+3}{-3} = \frac{y+7}{2} = \frac{z-6}{4}$$

P and Q are the points on the path L1 and L2 respectively such that PQ is perpendicular on both paths L1 and L2. On the basis of above information, answer the following questions

(i) Find the length PQ

- (ii) Find the equation of PQ
- 2. A carpenter designed a Cuba of side a units and put it in 3 dimensional system such that one vertex at origin and adjacent sides on three coordinate axes as shown in figure



Based on the above information, answer the following questions:

(i) Write the coordinates of the vertices D, E, F and G.

(ii) Find the direction ratios of the diagonal OG.

(iii) Find the direction cosines of the diagonals CE and DB

OR

(iii) Find the angle between CE and DB.

# ANSWERS ONE MARK QUESTIONS

1.	(a)	$\sqrt{b^2 + c^2}$	8.	(b)	—1
2.	(c)	90°	9	(h)	aa' + cc' + 1 = 0
3.	(c)	$\frac{x-2}{3} = \frac{y+3}{4} = \frac{z-5}{-1}$	10	(0)	
4	(b)	$\lambda = 2$	10.	(b)	4
	(U)	$\mathcal{N} = \mathcal{Z}$	11.	(a)	
5.	(a)	$\frac{x-1}{0} = \frac{y+1}{2} = \frac{z}{-1}$	12.	(b)	
6.	(b)	(-2, 0, 7)	13.	(d)	
7.	(c)	5 units	14.	(c)	

# **TWO MARK QUESTIONS**

1.	$\frac{x-2}{1} = \frac{y}{2} = \frac{z-5}{3}$	7.	$\frac{x-2}{1} = \frac{y-1}{-7} = \frac{z-3}{2},$
2.	$\frac{6}{7}, \frac{2}{7}, \frac{-3}{7}$		$\vec{r} = (2\hat{i} + \hat{j} + 3\hat{k}) + \lambda(\hat{i} - 7\hat{j} + 2\hat{k})$
3.	2	8.	$\frac{x-2}{1} = \frac{y+3}{-1} = \frac{z-1}{-6},$
4.	$\vec{\mathbf{r}} = (2\hat{\mathbf{i}} + \hat{\mathbf{k}}) + \lambda(2\hat{\mathbf{i}} + 7\hat{\mathbf{j}} - 3\hat{\mathbf{k}})$		$r = (2I - 3J + K) + \lambda(I - J - 6K)$
5.	aa' + cc' + 1 = 0	9.	$\lambda = 4$ $\mu = 3$
		10.	$\lambda = 10$

# **THREE/FIVE MARK QUESTIONS**

1.	$\overline{r} = (2\hat{\imath} - \hat{\jmath} + \hat{k}) + \lambda(2\hat{\imath} - 2\hat{\jmath} + \hat{k}) \text{ and } \frac{x-2}{2} = \frac{y+1}{-2} = \frac{z-1}{1}$
2.	$\left(-\frac{51}{7},-\frac{18}{7},\frac{43}{7}\right)$
3.	$\left(\frac{1}{2},-\frac{1}{2},-\frac{3}{2}\right)$
4.	$\frac{1}{\sqrt{6}}$
5.	$2\sqrt{29}$ units
6.	$\frac{8}{\sqrt{29}}$
7.	$(1, 2, 3), \sqrt{14}$
9.	$SD = 14 \text{ units}, \ \frac{x-5}{2} = \frac{y-7}{3} = \frac{z-3}{6}$
11.	K = 12
12.	$\frac{x-1}{3} = \frac{y-1}{10} = \frac{z-1}{17}$
13.	$\frac{x}{1} = \frac{y}{2} = \frac{z}{-1}$ and $\frac{x}{-1} = \frac{y}{1} = \frac{z}{-2}$
14.	(1, 2, 3), $\sqrt{14}$ , (0, 5, 1)
15.	(6, 7, 3)
16.	$\left(6, \frac{19}{2}, 18\right), \left(-2, \frac{-5}{2}, -6\right)$
17.	$\frac{9}{\sqrt{195}}$ , Not intersecting
18.	(3, 5, 9), $\frac{x-1}{2} = \frac{y-2}{3} = \frac{z-3}{6}$ , 7 units
19.	$\frac{x+1}{2} = \frac{y-3}{-7} = \frac{z+2}{4}$

20. 
$$(-2, -1, 3), \left(\frac{56}{17}, \frac{43}{17}, \frac{11}{17}\right)$$
  
21.  $PQ: \vec{r} = (4\hat{i} + 5\hat{j} + 10\hat{k}) + \lambda(2\hat{i} + 2\hat{j} + 6\hat{k})$   
 $QR: \vec{r} = (2\hat{i} + 3\hat{j} + 4\hat{k}) + \mu(\hat{i} + \hat{j} + 5\hat{k}), \text{Point R}(3, 4, 5)$   
22.  $\frac{x-3}{1} = \frac{y+1}{-6} = \frac{z-11}{4}, (2, 5, 7), \sqrt{13} \text{ units}$   
23.  $(-1, -1, -1)$   
24.  $\lambda \neq 69/11$   
25.  $\vec{r} = (2\hat{i} + \hat{j} - \hat{k}) + \mu(2\hat{i} - \hat{j} + \hat{k}), \sqrt{\frac{11}{6}} \text{ units}$   
**SELF ASSESSMENT TEST-1**  
1. (C) 2. (B) 3. (B) 4. (A) 5. (D)  
**SELF ASSESSMENT TEST-2**  
1. (A) 2. (D) 3. (C) 4. (B) 5. (C)

**CASE STUDY BASED QUESTIONS**  
1. (i) 
$$3\sqrt{30}$$
 units (ii)  $\vec{r} = (3\hat{i} + 8\hat{j} + 3\hat{k}) + \lambda(2\hat{i} + 5\hat{j} - \hat{k})$   
2. (i)  $D(a, 0, a), E(a, a, 0), F(0, a, a) \& G(a, a, a)$   
(ii) <1, 1, 1>  
(iii) Direction cosines of CE are  $\left\langle \frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, -\frac{1}{\sqrt{3}} \right\rangle$   
DB are  $\left\langle -\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}, -\frac{1}{\sqrt{3}} \right\rangle$   
OR  
Angle between CE and DB =  $\cos^{-1}\left(\frac{1}{3}\right)$ 

# CHAPTER-12

# LINEAR PROGRAMMING

Linear programming is used to obtain optimal solutions for operations research. Using LPP, researchers find the **best**, most economical **solution** to a problem within all of its **limitations**, or constraints.

Few examples of applications of LPP

- (i) Food and Agriculture: In nutrition, Linear programming provides a powerful tool to aid in planning for dietary needs. Here, we determine the different kinds of foods which should be included in a diet so as to minimize the cost of the desired diet such that it contains the minimum amount of each nutrient.
- (ii) **Transportation:** Systems rely upon linear programming for cost and time efficiency.

**Airlines** use linear programming to optimize their profits according to different seat prices and customer demand. Because of this only, efficiency of airlines increases and expenses are decreased.

### TOPICS TO BE COVERED AS PER CBSE LATEST CURRICULUM 2024-25

- Introduction, constraints, objective function, optimization.
- Graphical method of solution for problems in two variables.
- · Feasible and infeasioble region (bounded or unbounded)
- · Feasible and infeasible solutions.
- Optimal feasible solutions (upto three non-trival constraints)

### **KEY POINTS :**

- **OPTIMISATION PROBLEM :** is a problem which seeks to maximize or minimize a function. An optimisation problem may involve maximization of profit, minimization of transportation cost etc, from available resources.
- A LINEAR PROGRAMMING PROBLEM (LPP) : LPP deals with the optimisation (maximisation/minimisation) of a linear function of two variables (say x and y) known as objective function subject to the conditions that the variables are non negative and satisfy a set of linear inequalities (called linear constraints). A LPP is a special type of optimisation problem.
- **OBJECTIVE FUNCTION**: Linear function *z* = *ax* + *by* where a and b are constants which has to be maximised or minimised is called a linear objective function.
- **DECISION VARIABLES** : In the objective function *z* = *ax* + *by*, *x* and *y* are called decision variables.
- **CONSTRAINTS** : The linear inequalities or restrictions on the variables of an LPP are called constraints.

The conditions  $x \ge 0$ ,  $y \ge 0$  are called non-negative constraints.

- FEASIBLE REGION : The common region determined by all the constraints including non-negative constraints x ≥ 0, y ≥ 0 of a LPP is called the feasible region for the problem.
- **FEASIBLE SOLUTION :** Points within and on the boundary of the feasible region for a LPP represent feasible solutions.
- **INFEASIBLE SOLUTIONS** : Any point outside the feasible region is called an infeasible solution.
- **OPTIMAL (FEASIBLE) SOLUTION :** Any point in the feasible region that gives the optimal value (maximum or minimum) of the objective function is called an optimal solution.
- **THEOREM 1**: Let R be the feasible region (convex polygon) for a LPP and let z = ax + by be the objective function. When z has an optimal value (maximum or minimum), where x and y are subject to constraints described by linear inequalities, this optimal value must occur at a corner point (vertex) of the feasible region.
- **THEOREM 2**: Let R be the feasible region for a LPP. & let z = ax + by be the objective function. If R is bounded, then the objective function z has both a maximum and a minimum value on R and each of these occur at a corner point of R.

If the feasible region R is unbounded, then a maximum or minimum value of the objective function may or not exist. However, if it exists it must occur at a corner point of R.

MULTIPLE OPTIMAL POINTS : If two corner points of the feasible region are
optimal solutions of the same type i.e both produce the same maximum or
minimum, then any point on the line segment joining these two points is also an
optimal solution of the same type.

### Illustration:

A company produces two types of belts A and B. Profits on these belts are Rs. 2 and Rs. 1.50 per belt respectively. A belt of type A requires twice as much time as belt of type B. The company can produce atmost 1000 belts of type B per day. Material for 800 belts per day is available. Atmost 400 buckles for belts of type A and 700 for type B are available per day. How much belts of each type should the company produce so as to maximize the profit?

**Solution:** Let the company produces *x* no. of belts of type A and *y* no. of belts of type B to maximize the profit.



[Class XII : Maths]

ι	Jsing Corner point method.		
	Corner Points	Obj. fn. z = 2x + 1.5y	
	O (0, 0)	0	
	A (400, 0)	800	
	B (400, 200)	1100	
	C (200, 600)	1300	max <i>z</i> .
	D (100, 700)	1250	
	E (0, 700)	1050	

 $\therefore$  Optimal solution is given by C(200, 600)

i.e. company should produce 200 belts of type A and 600 belts of type B so as to maximize the profit of Rs. 1300.

[Class XII : Maths]

### **ONE MARK QUESTIONS**

- 1. The solution set of the inequation 3x + 4y < 7 is:
  - (a) Whole xy plane except the points lying on the line 3x + 5y = 7
  - (b) Whole xy plane alogn with the points lying on the line 3x + 5y = 7
  - (c) Open half plane containing the origin except the point of line 3x + 5y = 7
  - (d) Open half plane not containing the origin except the point of line 3x + 5y = 7
- 2. Which of the following points solisfies both the inequations  $2x + y \le 10$  and  $x + 2y \ge 8$ ?

- (c) (-5, 6) (d) (4, 2)
- 3. The objective function Z = ax + by of LPP has maximum value 42 at (4, 6) and minimum value 19 at (3, 2). Which of the following is true?

(a) a = 9, b = 1	(b) a = 5, b = 2
(c) a = 3, b = 5	(d) a = 5, b = 3

4. The corner points of the feasible region of a LPP are (0, 4), (7, 0) and  $\left(\frac{20}{2}, \frac{4}{2}\right)$ . If

z = 30x + 24y is the objective functions, then (maximum value of z-minimum value of z) is equal to

- (a) 40 (b) 96
- (c) 120 (d) 136
- 5. The minimum value of z = 3x + 8y subject to the constraints x  $\leq$  20, y  $\geq$  10 and x  $\geq$  0, y  $\geq$  0 is

(a) 80	(b) 140
--------	---------

- (c) 0 (d) 60
- 6. The number of corner points of the feasible region determined by the constraints

$x-y\geq 0,\ 2y\leq x+2,\ x\geq 0,\ y\geq 0 \ is$	
(a) 2	(b) 3

- (c) 4 (d) 5
- 7. The no. of feasible solutions of the L.P.P. given as maximise *z* = 15*x* + 30*y* subject the constraints:

 $3x + y \le 12, x + 2y \le 10, x \ge 0, y \ge 0$  is

- (a) 1 (b) 2
- (c) 3 (d) infinite

8. The feasible region of a linear programming problem is shown in the figure below:



Which of the following are the possible constraints?

- (a)  $x + 2y \ge 4, x + y \le 3, x \ge 0, y \ge 0$
- (b)  $x + 2y \le 4, x + y \le 3, x \ge 0, y \ge 0$
- (c)  $x + 2y \ge 4$ ,  $x + y \ge 3$ ,  $x \ge 0$ ,  $y \ge 0$
- (d)  $x + 2y \ge 4$ ,  $x + y \le 3$ ,  $x \le 0$ ,  $y \le 0$
- 9. L.P.P. is a process of finding
  - (a) Maximum value of the objective function
  - (b) Minimum value of the objective function
  - (c) Optimum value of the objective function
  - (d) None of these
- 10. Which of the following statements is correct?
  - (a) Every L.P.P. admits an optimal solution
  - (b) A L.P.P. admits a unique optimal solution
  - (c) If a L.P.P. admits two optimal solutions, it has an infinite number of optimal solutions
  - (d) The set of all feasible solution of a L.P.P. is not a convex set
- 11. Region represented by  $x \ge 0$ ,  $y \ge 0$  is
  - (a) First quadrant (b) Second quadrant
  - (c) Third quadrant (d) Fourth quadrant

[Class XII : Maths]

12. The feasible region for L.P.P. is shown shaded in the figure. Let f = 3x - 4y be the objective function, then maximum value of *f* is



(c) I, II and III quadrant (d) II, III, IV quadrant

### **ASSERTION-REASON TYPE QUESTIONS**

Directions: Each of these questions contains two statements, Assertion (A) and Reason (R). Choose the correct answer out of the following choices.

(a) Both (A) and (R) are true and (R) is the correct explanation of (A)

- (b) Both (A) and (R) are true but (R) is not the correct explanation of (A)
- (c) (A) is true and (R) is false
- (d) (A) is false but (R) is ture
- 16. Assertion (A) : If a L.P.P. admits two optional solution then it has infinitely many optimal solution.

Reason (R) : If the value of the objective function of a L.P.P. is same at two corners then it is same at every point on the line segment joining the two corner pionts.

17. Assertion (A): The solution region satisfied by the inequalities  $x + y \le 5$ ,  $x \le 4$ ,  $y \le 4$ ,  $x \ge 0$ ,  $y \ge 0$  is bounded.

Reason (R) : A region in x-y plane is said to be bounded if it can be enclosed within a circle.

18. Assertion (A) : Minimize  $z = x^2 + 2xy + y^2$  can be considered as the objective function for the L.P.P.

Reason (R) : Objective function of the L.P.P. is of this type z = ax + by; a and b are real numbers i.e. z is linear function of x and y.

19. Assertion (A) : The region represented by the inequalities  $x \ge 6$ ,  $y \ge 2$ ,  $2x + y \ge 10$ ,  $x \ge 0$ ,  $y \ge 0$  is empty.

Reason (R) : There is no (x, y) that satisfies all the constraints.

Assertion (A) : Corner points of the feasible region for an L.P.P. are (0, 2), (3, 0), (6, 0), (6, 8) and (0, 5). Let F = 4x + 6y be the objective function. The minimum value of F occurs at (0, 2) only.

Reason (R) : Minimum value of F occurs at all the infinite no. points that lie on the line segment joining (0, 2) and (3, 0).

### THREE MARKS QUESTIONS

1. Solve the following linear programming problem graphically:

Maximise z = -3x - 5ysubject to the constraints  $-2x + y \le 4$  $x + y \ge 3$  $x - 2y \le 2$  $x \ge 0, y \ge 0$ 

[Class XII : Maths]

2. Solve the following LPP graphically:

Maximise z = 5x + 3ys.t. the constraints  $3x + 5y \le 15$  $5x + 2y \le 10$ x,  $y \ge 0$ Solve the following LPP graphically Maximise z = x + 2ys.t.  $x + 2y \ge 100$  $2x - y \le 0$ 

3.

$$2x - y \le 0$$
$$2x + y \le 200$$

 $x\geq 0,\ y\geq 0$ 

- 4. The objective function z = 4x + 3y of a LPP under some constraints is to be maximized and minimized. The corner points of the feasible region are A(0, 700), B(100, 700), C(200, 600) and D(400, 200). Find the point at which z is maximum and the point at which z is minimum. Also find the corresponding maximum and minimum values of z.
- 5. Solve graphically

Minimise : z = -3x + 4ys.t.  $3x + 2y \le 12$ 

 $3x + 2y \le 12$ x, y \ge 0

6. Solve the following LPP graphically

 $\begin{array}{ll} \mbox{Minimise:} & Z=60x+80y\\ \mbox{s.t.} & 3x+4y\geq 8\\ & 5x+2y\geq 11\\ & x,\,y\geq 0 \end{array}$ 

7. Solve graphically

Maximise : z = 600x + 400y

s.t. 
$$x + 2y \le 12$$
  
 $2x + y \le 12$   
 $x + 1. 25y \ge 5$   
 $x, y \ge 0$ 

8. Solve graphically

Maximise : P = 100x + 5y

s.t.  $\begin{aligned} x+y &\leq 300\\ 3x+y &\leq 600\\ y &\leq x+200 \end{aligned}$ 

9. Solve the LPP graphically

 $\begin{array}{ll} \mbox{Minimize} & z=5x+10y\\ \mbox{s.t.} & x+2y\leq 120,\,x+y\geq 60,\,x-2y\geq 0\\ & x\geq 0,\,y\geq 0 \end{array}$ 

10. Determine graphically the minimum value of the following objective function:

$$\begin{array}{l} z = 500x + 400y \\ \text{s.t.} \ x + y \ \leq \ 200 \\ x \geq \ 20 \\ y \geq \ 4x \\ y \geq \ 0 \end{array}$$

### **FIVE MARKS QUESTIONS**

Q. 1 Solve the following LPP graphically.

Maximize z = 3x + y subject to the constraints

$$x + 2y \ge 100$$
$$2x - y \le 0$$
$$2x + y \le 200$$
$$x, y \ge 0$$

Q.2 The corner points of the feasible region determined by the system of linear constraints are as shown below.



Answer each of the following :

- (i) Let z = 3x 4y be the objective function. Find the maximum and minimum value of z and also the corresponding points at which the maximum and minimum value occurs.
- (ii) Let z = px + qy where p, q > 0 be the objective function. Find the condition on p and q so that the maximum value of z occurs at B (4, 10) and C (5, 8). Also mention the number of optimal solutions in this case.
- Q. 3 There are two types of fertilisers A and B. A consists of 10% nitrogen and 6% phosphoric acid and B consists of 5% nitrogen and 10% phosphoric acid. After testing the soil conditions, a farmer finds that he needs at least 14 kg of nitrogen and 14 kg of phosphoric acid for his crop. If A costs Rs. 6 per kg and B costs Rs. 5 per kg. determine how much of each type of fertiliser should be used so that nutrient requirements are met at minimum cost. What is the minimum cost?
- Q. 4 A man has Rs. 1500 to purchase two types of of shares of two different companies S1 and S2. Market price of one share of S1 is Rs. 180 and S2 is Rs 120. He wishes to purchase a maximum of ten shares only. If one share of type S1 gives a yield of Rs 11, and of type S2 yields Rs 8 then how much shares of each type must be purchased to get maximum profit? and what will be the maximum profit?
- Q. 5 A company manufactures two types of lamps say A and B. Both lamps go through a cutter and then a finisher. Lamp A requiires 2 hours of the cutter's time and 1 hour of the finisher's time. Lamp B required 1 hr of cutter's, 2 hrs of finisher's time. The cutter has 100 hours and finisher has 80 hours of time available each month. Profit on one lamp A is Rs. 7.00 and on one lamp B is Rs 13.00. Assuming that he can sell all that he produces how many of each type of lamps should be manufactured to obtain maximum profit and what will be the maximum profit?
- Q.6 A dealer wishes to purchase a number of fans and sewing machines. He has only Rs. 5760 to invest and has space for atmost 20 items. A fan and sewing machine cost Rs 360 and Rs. 240 respectively. He can sell a fan at a profit of Rs. 22 and sewing machine at a profit of Rs. 18. Assuming that he can sell whatever he buys, how should he invest money to maximise his profit?
- Q. 7 A producer has 20 and 10 units of labour and capital respectively which he can use to produce two kinds of goods X and Y. To produce one unit of X, 2 units of capital and 1 unit of labour is required. To produce one unit of Y, 3 units of labour and 1 unit of capital is required. If X and Y are priced at Rs. 80 and Rs100 per unit respectively, how should the producer use his resources to maximize revenue?
- Q. 8 A factory owner purchases two types of machines A and B for his factory. The requirements and limitations for the machines are as follows :

Machine	Area Occupied	Labour Force	Daily Output
			(in units)
A	1000 m <sup>2</sup>	12 men	50
В	1200 m <sup>2</sup>	8 men	40

He has maximum area of 7600 m<sup>2</sup> available and 72 skilled labourers who can operate both the machines. How many machines of each type should he buy to maximise the daily output?

Q.9 A manufacturer makes two types of cups A and B. Three machines are required to manufacture the cups and the time in minutes required by each in as given below :

Types of Cup	Machines			
	I	II	111	
A	12	18	6	
В	6	0	9	

Each machine is available for a maximum period of 6 hours per day. If the profit on each cup A is 75 paisa and on B is 50 paisa, find how many cups of each type should be manufactured to maximise the profit per day.

- Q. 10 An aeroplane can carry a maximum of 200 passengers. A profit of Rs. 400 is made on each first class ticket and as profit of Rs. 300 is made on each second class ticket. The airline reserves atleast 20 seats for first class. However at least four times as many passengers prefer to travel by second class than by first class. Determine how many tickets of each type must be sold to maximize profit for the airline.
- Q. 11 A diet for a sick person must contains at least 4000 units of vitamins, 50 units of minerals and 1400 units of calories. Two foods A and B are available at a cost of Rs. 5 and Rs. 4 per unit respectively. One unit of food A contains 200 units of vitamins, 1 unit of minerals and 40 units of calories whereas one unit of food B contains 100 units of vitamins, 2 units of minerals and 40 units of calories. Find what combination of the food A and B should be used to have least cost but it must satisfy the requirements of the sick person.
- Q.12 Anil wants to invest at most Rs. 12000 in bonds A and B. According to the rules, he has to invest at least Rs. 2000 in Bond A and at least Rs. 4000 in bond B. If the rate of interest on bond A and B are 8% and 10% per annum respectively, how should he invest this money for maximum interest? Formute the problem as LPP and solve graphically.

# CASE STUDY QUESTIONS

Q. 1 A man rides his motorcycle at the speed of 50 km/hr. He has to spend Rs 2/km on petrol. But if he rides it at a faster speed of 80 km/hr, the petrol cost increases to Rs 3/km. He has atmost Rs 120 to spend on petrol and one hr's time. he wishes to find the maximum distance that he can travel.



Based on the above information answer the following questions.

- (1) If he travels x km with the speed of 50 km/hr and y km with the speed of 80 km/hr, then write the objective function
- (2) Find the Maximum distance man can travel?
- Q.2 Two tailors A and B earn Rs 150 and Rs 200 per day respectively. A can sticth 6 shirts and 4 pants per day, while B can stitch 10 shirts and 4 pants per day. it is desired to produce atleast 60 shirts and 32 pants at a minimum labour cost.





Tailor B

Based on the above information answer the following.

- (1) If x and y are the number of days A and B work respectively then find the objective function for this LPP
- (2) Find the optimal solution for this LPP and the minimum labour cost?

### SELF ASSESSMENT-1

EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. Objective functin of a L.P.P. is
  - (a) A constraint
  - (b) A function of be optimised
  - (c) A relation between the variables
  - (d) None of these
- 2. The solution set of the inequation 2x + y > 5 is
  - (a) Open half plans that contains the origion
  - (b) Open half plane not containing the origin
  - (c) Whole xy-plane except the points lying on the line 2x + y = 5
  - (d) None of these
- 3. Which of the following statements is correct?
  - (a) Every L.P.P admits an optimal solution
  - (b) A L.P.P. admits unique optimal solution
  - (c) If a LPP admits two optimal solutions, it has an infinite number of optimal solutions
  - (d) None of these
- 4. Solution set of inequation  $x \ge 0$  is
  - (a) Half plane on the left of y-axis
  - (b) Half plane on the right of y-axis excluding the points on y-axis
  - (c) Half plane on the right of y-axis including the points on y-axis
  - (d) None of these
- 5. In a LPP, the constraints on the decision variables x and y are

 $x-3y\geq 0,\ y\geq 0,\ 0\leq x\leq 3.$ 

The feasible region

- (a) is not in the first quadrant
- (b) is bounded in the first quadrant
- (c) is unbounded in the first quadrant
- (d) doesn't exist

### **SELF ASSESSMENT-2**

EAH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

- 1. Solution set of the inequation  $y \le 0$  is
  - (a) Half plane below the x-axis excluding the points on x-axis
  - (b) Half plane below the x-axis including the points on x-axis
  - (c) Half plane above the x-axis
  - (d) None of these
- 2. Regions represented by inequations  $x \ge 0$ ,  $y \ge 0$  is
  - (a) first quadrant
  - (b) second quadrant
  - (c) third quadrant
  - (d) fourth quadrant
- 3. The feasible region for an LPP is always
  - (a) concavo convex polygen
  - (b) concave poloygon
  - (c) convex polygon
  - (d) None of these
- 4. If the constraints in a linear programming problem are changed then
  - (a) the problem is to be reevaluated
  - (b) solution not defined
  - (c) the objective function has to be modified
  - (d) the change in constraints is ignored
- 5. L.P.P. is as follows:
  - Minimize Z = 30x + 50y

Subject to the constraints,

 $3x + 5y \ge 15$ 

 $2x + 3y \le 18$ 

 $x \ge 0, y \ge 0$ 

In the feasible region, the minimum value of Z occurs at

- (a) a unique point
- (b) no point
- (c) infinitely many points
- (d) two points only

### ANSWER

### **One Marks Questions**

1. (c)	2.	(d) (4, 2)	3.	(c) <i>a</i> = 3, <i>b</i> = 5
4. (d) 136	5.	(a) 80	6.	(a) 2
7. (d) infinite	8.	(c)	9.	(c)
10. (c)	11.	(a)	12.	(c) 0
13. (b) unbounded	14.	(c)Atriangle	15.	(c)
16. (a)	17.	(a)	18.	(d)
19. (a)	20.	(d)		

### **Three Marks Questions**

1. Optimal solution 
$$\left(\frac{8}{3}, \frac{1}{3}\right)$$
, maximize =  $\frac{-29}{3}$  feasible region unbounded.  
2. Optimal solution  $\left(\frac{20}{19}, \frac{45}{19}\right)$ , maximize =  $\frac{235}{19} = 12.3$ 

- 3. Optimal solution (0, 200), maximize = 400
- 4. Maximize z = 2600 at C(200, 600) and minimize z is 2100 at A(0, 700)
- 5. Minimize z = 12 at (4, 0)
- 6. Unbounded, minimize z = 160. It occurs at all the points on the line segment joining  $\left(2,\frac{1}{2}\right)$  and  $\left(\frac{8}{3},0\right)$ . So, infinite optimal solutions.
- 7. Maximize z = 4000 at (4, 4)
- 8. Maximize z = 20,000 at (200,0)
- 9. Miximize z = 300 at (60, 0)
- 10. Miximize z =42000 at (20, 80)

### **Five Marks Questions**

- 1. Max z = 250 at x = 50, y = 100
- 2. (i) Max z = 12 at (4,0) and min z = -32 at (0,8)

(ii) P = 2q, infinite solutions lying on the line segment joining the points B and C

- 3. 100 kg of fertilizer A and 80 kg of fertilizer B, minimum cost Rs 1000
- 4. Maximum profit = Rs 95 with 5 shares of each type.
- 5. Lamps of type A = 40, Lamps of type B = 20 Max profit = Rs 540
- 6. Fans: 8, sewing machines: 12, max profit: Rs 392
- 7. X: 2 units, Y: 6 units, max revenue is Rs 760.

[Class XII : Maths]

- 8. Type A : 4, Type B : 3
- 9. Cup A : 15, cup B : 30

10. No of first class ticket = 40, No of second class tickets = 160

11. Food A:5 units, food B: 30 units

12. Maximum interest is Rs 1160 at (2000, 10000)

### CASE STUDIES QUESTIONS

1.	(i) maximize z	= x	+ y	(ii) 54 <mark>2</mark> km		
2.	(i) minimize z = 150x + 200y		)x + 200y	(ii) (5, 3) and Rs	. 1350	
			SELF	ASSESSMENT-1		
1.	(b)	2.	(b) 3	3. (c)	4. (c)	5. (b)
			SELF	ASSESSMENT-2		
1.	(b)	2.	(a) 3	3. (c)	4. (a)	5. (c)
## CHAPTER-13

# PROBABILITY

Probability is the branch of mathematics that deals with assigning a numerical quantity  $(0 \le p \le 1)$  to the happening/non happening of any event.



→ How likely a natural disaster like earthquake, hurricane etc. will strike the country in a given year.



A sports betting company may look at the current record of two teams A and B and determine which team has higher probability of winning and do the sports betting accordingly.

#### TOPICS TO BE COVERED AS PER CBSE LATEST CURRICULUM (2024-25)

- · Conditional probability
- · Multiplication theorem on probability
- · Independent events
- Total probability and Baye's theorem
- · Random variable and its probability distribution
- Mean of random variable

#### **KEY POINTS**

**Conditional Probability :** If *A* and *B* are two events associated with the same sample space of a random experiment, then the conditional probability of the event *A* under the condition that the event *B* has already occurred, written as P(A|B), is given by

$$P(A|\mathsf{B}) = \frac{(P \cap B)}{P(B)}, \quad P(B) \neq 0.$$

**Properties :** 

- (1) P(S|F) = P(F|F) = 1 where S denotes sample space
- (2)  $P((A \cup B)|F) = P(A|F) + P(B|F) P((A \cap B)|F)$
- (3) P(E'|F) = 1 P(E|F)

**Multiplication Rule :** Let *E* and *F* be two events associated with a sample place of an experiment. Then

$$P(E \cap F) = P(E) P(F|E) \text{ provided } P(E) \neq 0$$
$$= P(F) P(E|F) \text{ provided } P(F) \neq 0.$$

If E, F, G are three events associated with a sample space, then

 $P(E \cap F \cap G) = P(E) P(F|E) P(G|(E \cap F))$ 

**Independent Events :** Let *E* and *F* be two events, then if probability of one of them is not affected by the occurrence of the other, then *E* and *F* are said to be independent, i.e.,

(a)  $P(F|E) = P(F), \quad P(E) \neq 0$ 

or (b)  $P(E|F) = P(E), \quad P(F) \neq 0$ 

or (c)  $P(E \cap F) = P(E) P(F)$ 

Three events A, B, C are mutually independent if

$$P(A \cap B \cap C) = P(A) P(B) P(C)$$
$$P(A \cap B) = P(A) P(B)$$
$$P(B \cap C) = P(B) P(C)$$
$$P(A \cap C) = P(A) P(C)$$

**Partition of a Sample Space :** A set of events  $E_1, E_2, ..., E_n$  is said to represent a partition of a sample space S if

(a)  $E_i \cap E_i = \phi; i \neq j; i, j = 1, 2, 3, ..., n$ 

$$(b)E_1 \cup E_2 \cup E_3 \dots \cup E_n = S$$
 and

(c) Each 
$$E_i \neq \phi$$
 i.e.  $P(E_i) > 0 \forall i = 1, 2, ..., n$ 

[Class XII : Maths]

and

**Theorem of Total Probability :** Let  $\{E_1, E_2, ..., E_n\}$  be a partition of the sample space *S*. Let *A* be the any event associated with *S*, then

$$P(A) = \sum_{j=1}^{n} P(E_j) P(A|E_j)$$

**Baye's Theorem :** If  $E_1$ ,  $E_2$ , ...,  $E_n$  are mutually exclusive and exhaustive events associated with a sample space S, and A is any event associated with  $E_i$ 's having non-zero probability, then

$$P(E_{i}|A) = \frac{P(A | E_{i})P(E_{i})}{\sum_{i=1}^{n} P(A | E_{i})P(E_{i})}$$

**Random Variable :** A (r.v.) is a real variable which is associated with the outcome of a random experiment.

Probability Distribution of a r.v. X is the system of numbers given by

X:
 
$$x_1$$
 $x_2$ 
 ....
  $x_n$ 

 P(X = x):
  $p_1$ 
 $p_2$ 
 ....
  $p_n$ 

where

$$p_i > 0, \quad i = 1, 2, ..., n, \sum_{i=1}^n p_i = 1.$$

Mean of a r.v. X :

$$\mu = E(X) = \sum_{i=1}^{n} p_i x_i$$

## Illustration:

Evaluate P(A 
$$\cup$$
 B) if 2P(A) = P(B) =  $\frac{5}{13}$  and P(A|B) =  $\frac{2}{5}$ 

**Solution:** 
$$2P(A) = P(B) = \frac{5}{13}$$
  
 $\Rightarrow P(A) = \frac{5}{26}, P(B) = \frac{5}{13}$ 

As 
$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$
  

$$\Rightarrow \frac{2}{5} = \frac{P(A \cap B)}{(5/13)} \Rightarrow \frac{2}{5} \times \frac{5}{13} = P(A \cap B)$$

$$\Rightarrow \frac{2}{13} = P(A \cap B)$$
Now,  $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ 

$$= \frac{5}{26} + \frac{5}{13} - \frac{2}{13} = \frac{11}{16}$$

## Illustration:

Prove that if E and F are independent events, then the events E and F' are also independent.

**Solution:**  $P(E \cap F) = P(E) P(F)$  (given)

Consider,

$$P(E \cap F') = P(E) - P(E \cap F)$$
$$= P(E) - P(E) P(F)$$
$$= P(E) (1 - P(F))$$
$$P(E \cap F') = P(E) - P(F')$$

So, E and F' are also independent.

## Illustration:

A card from a pack of 52 cards is lost. From the remaining cards of the pack, two cards are drawn. What is the probability that they both are diamonds.

**Solution:** Let  $E_1 = lost card is diamond$ 

 $E_2 = lost card is non-diamond$ 

A = 2 diamonds cards are drawn from the remaining cards

Using Theorem of total probability

$$P(A) = P(A|E_1) P(E_1) + P(A|E_2) P(E_2)$$
$$= \frac{12}{51} \times \frac{11}{50} \times \frac{13}{52} \times \frac{12}{50} \times \frac{13}{51} \times \frac{39}{52^4}$$

$$= \frac{132}{10200} + \frac{468}{10200} = \frac{.600^{-}}{.102000^{-17}} \frac{1}{17}$$

## Illustration:

Three cards are drawn at random (without replacement) from a well shuffled pack of 52 playing cards. Find the probability distribution of number of red cards. Hence, find the mean of the distribution.

Solution: Let X denotes the number of red cards

$$\therefore P(X = 0) = \frac{{}^{26}C_3}{{}^{52}C_3} = \frac{26 \times 25 \times 24}{52 \times 51 \times 50} = \frac{2}{17} = \frac{4}{34}$$

$$P(X = 1) = \frac{{}^{26}C_1 \times {}^{26}C_2}{{}^{52}C_3} = 26 \times \frac{26 \times 25}{2} \times \frac{3 \times 2 \times 1}{52 \times 51 \times 50} = \frac{13}{34}$$

$$P(X = 2) = \frac{{}^{26}C_2 \times {}^{26}C_1}{{}^{52}C_3} = \frac{26 \times 25 \times 26 \times 3 \times 2 \times 1}{2 \times 52 \times 51 \times 50} = \frac{13}{34}$$

$$P(X = 3) = \frac{{}^{26}C_3}{{}^{52}} = \frac{4}{54}$$

$$(X = 3) = \frac{C_3}{5^2 C_2} = \frac{4}{34}$$

## : Probability Distribution

	Х	P(X = x)	X.P(x)		
	0	$\frac{4}{34}$	0		
	1	$\frac{13}{34}$	<u>13</u> 34		
	2	<u>13</u> 34	$\frac{26}{34}$		
	3	$\frac{4}{34}$	<u>12</u> 34		
		$\sum p_i = 1$	$\overline{\mathbf{x}} = \sum \mathbf{p}_i \mathbf{x}_i$		
•	$\overline{x} = \sum p_i x_i = \frac{13}{34} + \frac{26}{34} + \frac{12}{34} = \frac{51}{34} = \frac{3}{2}$				

## **ONE MARK QUESTIONS**

- 1. The events E and F are independent. If P(E) = 0.3 and P(EUF) = 0.5, then P(E/F) P(F/E) equals:
- (b)  $\frac{2}{7}$ (a)  $\frac{1}{7}$ 3 (d)  $\frac{1}{70}$ (c)  $\frac{1}{35}$ 2. For two events A and B, if P(A) = 0.4, P(B) = 0.8, P(B/A) = 0.6, then P(AUB) is: (a) 0.24 (b) 0.3 (d) 0.96 (c) 0.48 If A and B are two events such that  $P(A|B) = 2 \times P(B|A)$  and  $P(A) + P(B) = \frac{2}{3}$ , then 3. P(B) is equal to (a)  $\frac{2}{9}$ (b)  $\frac{7}{9}$ (c)  $\frac{7}{9}$ (d)  $\frac{5}{9}$ 4. Two events A and B will be independent, if; (a) A and B are mutually exclusive (b) P(A) = P(B)(c) P(A'B') = [1 - P(A)] [1 - P(B)](d) P(A) + P(B) = 1If for any two events A and B, P(A) =  $\frac{4}{5}$  and P(A  $\cap$  B) =  $\frac{7}{10}$ , then P(B/A) is equal to 5. (a)  $\frac{1}{10}$ (b)  $\frac{1}{8}$ (c)  $\frac{7}{8}$ (d)  $\frac{17}{20}$ 6. Five fair coins are tossed simultaneously. The probability of the events that atleast one head comes up is (b)  $\frac{5}{32}$ 27 (a) 32 31 (d)  $\frac{1}{32}$ (C) 32 7. Ashima can hit a target 2 out of 3 times. She tried to hit the target twice. The probability that she missed the target exactly one is S 4

(a)	$\frac{2}{3}$	(b) $\frac{1}{3}$
(c)	4 9	(d) $\frac{1}{9}$

8.	If sum of numbers obtained on throwing number obtained on one of the dice is 4	g a pair of dice is 9, then the probability that is
	(a) $\frac{1}{9}$	(b) $\frac{4}{9}$
	(c) $\frac{1}{18}$	(d) $\frac{1}{2}$
9.	X&Y are independent events such that F	$P(X \cap \overline{y}) = \frac{2}{5}$ and $P(X) = \frac{3}{5}$ . Then $P(Y)$ is equal
	to	
	(a) $\frac{2}{3}$	(b) $\frac{2}{5}$
	(c) $\frac{1}{3}$	(d) $\frac{1}{5}$
10.	If for two events A and B, $P(A - B) = \frac{1}{5}$	and P(A) = $\frac{3}{5}$ , then $P\left(\frac{B}{A}\right)$ is equal to
	(a) $\frac{1}{2}$	(b) $\frac{3}{5}$
	(c) $\frac{2}{5}$	(d) $\frac{2}{3}$
11.	If A and B are two events such that P(A)	) > 0 and P(B) $\neq$ 1, then $P(\overline{A}   \overline{B})$ =
	(a) 1- <i>P</i> ( <i>A</i> / <i>B</i> )	(b) $1-P(A/\overline{B})$
	(c) $\frac{1-P(A\cup B)}{P(B)}$	(d) $\frac{P(\overline{A})}{P(B)}$
12.	A and B are events such that $P(A/B) = F$	P(B/A) then
	(a) <i>A</i> ⊂ <i>B</i>	(b) B = A
	(c) $A \cap B = \phi$	(d) $P(A) = P(B)$
13.	Two aeroplanes I and II bomb a target in s a hit correctly are 0.3 and 0.2 respective misses the target. The probability that th	succession. The probabilities of I and II scoring ly. The second plane will bomb only if the first he target is hit by the II plane is
	(a) 0.2	(b) 0.7
	(c) 0.06	(d) 0.14
14.	$P(E \cap F)$ is equal to	
	(a) P(E) P(F/E)	(b) P(F).P(E/F)
	(c) Both (a) & (b)	(d) None of these

15. Two dice are thrown. If it is known that the sum of the numbers on the dice is less than 6, the probability of getting a sum 3 is

(a) 
$$\frac{1}{8}$$
 (b)  $\frac{2}{5}$   
(c)  $\frac{1}{5}$  (d)  $\frac{5}{18}$ 

In following questions Q16 to Q20, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both (A) and (R) are true and (R) is the correct explanation of (A)
- (b) Both (A) and (R) are true but (R) is not the correct explanation of (A)
- (c) (A) is true and (R) is false
- (d) (A) is false but (R) is true
- Assertion (A) : The mean of a random variable X is also called the expectation of x, denoted by E(x).

Reason (R): The mean or expectation of a random variable X is the sum of the producuts of all possible values of x by their respective probabilities.

17. Assertion (A) : Let A and B be two independent events. The  $P(A \cap B) = P(A) + P(B)$ 

Reason (R): Three events A, B and C are said to be independent if

 $P(A \cap B \cap C) = P(A). P(B). P(C)$ 

18. Assertion (A) : Two coins are tossed simultaneously. The probability ofgetting two

heads, if it is known that atleast one head comes up is  $\frac{1}{3}$ .

Reason (R): Let E and F be two events with a random experiment, then

$$P(F / E) = \frac{P(F \cap E)}{P(E)}$$

19. Assertion (A): The mean of the numbers obtained on throwing a die having written 1 on three faces, 2 on two faces and 5 on one face is 2

Reason (R): E(X) = mean of  $x = \sum_{i=1}^{n} p_i x_i$ 

20. Assertion (A) : Bag P contains 6 Red and 4 Blue balls and Bag Q contains 5 red and 6 Blue Balls. A ball is transferred from Bag P to bag Q and then a ball is drawn from Bag

Q. The probability that the ball drawn from bag Q is blue is  $\frac{8}{15}$ .

Reason (R): According to the law of total probability

 $P(A) = P(E_1)P(A / E_1) + P(E_2)P(A / E_2)$  where  $E_1$  and  $E_2$  partitions the sample space S and A is any event connected with  $E_1$  and  $E_2$ .

## **TWO MARKS QUESTIONS**

- 1. A and B are two events such that  $P(A) \neq 0$ , then find P(B|A) if (i) A is a subset of B (ii) A  $\cap$  B =  $\phi$ .
- 2. A random variable *X* has the following probability distribution, find *k*.

X	0	1	2	3	4	5
<i>P</i> ( <i>X</i> )	<u>1</u> 15	к	<u>15K – 2</u> 15	к	<u>15K – 1</u> 15	$\frac{1}{15}$

- 3. Out of 30 consecutive integers two are chosen at random. Find the probability so that their sum is odd.
- 4. Assume that in a family, each child is equally likely to be a boy or a girl. A family with three children is chosen at random. Find the probability that the eldest child is a girl given that the family has atleast one girl.

5. If *A* and *B* are such that  $P(A \cup B) = \frac{5}{9}$  and  $P(\overline{A} \cup \overline{B}) = \frac{2}{3}$ , then find  $P(\overline{A}) + P(\overline{B})$ .

- 6. Prove that if *A* and *B* are independent events, then *A* and *B'* are also independent events.
- 7. If A and B are two independent events such that P(A) = 0.3,  $P(A \cup B) = 0.5$ , then find P(A|B) P(B|A)
- 8. Three faces of an ordinary dice are yellow, two faces are red and one face is blue. The dice is rolled 3 times. Find the probability that yellow, red and blue face appear in the first, second and third throw respectively.
- 9. Find the probability that a leap year will have 53 Fridays or 53 Saturdays.
- 10. A person writes 4 letters and addresses on 4 envelopes. If the letters are placed in the envelopes at random, then what is the probability that all the letters are not placed in the right envelopes.

11. Find the mean of the distribution

X = x	0	1	2	3	4	5
P(X = x)	$\frac{1}{6}$	<u>5</u> 18	2 9	$\frac{1}{6}$	<u>1</u> 9	$\frac{1}{18}$

12. In a class XII of a school, 40% of students study Mathematics, 30% of the students study Biology and 10% of the class study both Mathematics and Biology. If a student is selected at random from the class, then find the probability that he will be studying Mathematics or Biology.

## THREE MARKS QUESTIONS

Q.1. A problem in mathematics is given to three students whose chances of solving it are

 $\frac{1}{2}$ ,  $\frac{1}{3}$  and  $\frac{1}{4}$ . What is the probability that the problem is solved ?

- Q.2. If *A* and *B* are two independent events such that  $P(\overline{A} \cap B) = \frac{2}{15}$  and  $P(A \cap \overline{B}) = \frac{1}{6}$  then find P(A) and P(B).
- Q.3. From a lot of 20 bulbs which include 5 defectives, a sample of 2 bulbs is drawn at random, one by one with replacement. Find the probability distribution of the number of defective bulbs. Also, find the mean of the distribution.
- Q.4. Amit and Nisha appear for an interview for two vacancies in a company. The probability of Amit's selection is 1/5 and that of Nisha's selections is 1/6. What is the probability that
  - (i) both of them are selected?
  - (ii) only one of them is selected?
  - (iii) none of them is selected?
- Q.5. In a game, a man wins a rupee for a six and looses a rupee for any other number when a fair die is thrown. The man decided to throw a die thrice but to quit as and when he gets a six. Find the expected value of the amount he wins/looses.
- Q.6. Suppose that 10% of men and 5% of women have grey hair. A grey haired person is selected at random. What is the probability that the selected person is male assuming that there are 60% males and 40% females ?
- Q.7. Two dice are thrown once. Find the probability of getting an even number on the first die or a total of 8.

- Q.8. Two aeroplanes X and Y bomb a target in succession. There probabilities to hit correctly are 0.3 and 0.2 respectively. The second plane will bomb only if first miss the target. Find the probability that target is hit by Y plane.
- Q.9. The random variable X can take only the values 0, 1, 2. Given that P(X=0) = P(X=1) = p and that  $E(X^2) = E(X)$ , find the value of p.
- Q.10. An urn contains 4 white and 3 red balls. Let X be the number of red balls in a random draw of 3 balls. Find the mean of X.
- Q11. A box contains 10 tickets, 2 of which carry a prize of Rupees 8 each, 5 of which carry a prize of Rupees 4 each and remaining 3 carry a prize of Rupees 2 each. If one ticket is drawn at random, find the mean value of the prize. Using the concept of probability distribution.
- Q.12. The probability distribution of a random variable X is given below:

Х	1	2	3
P(X)	K/2	K/3	K/6

- (i) Find the value of K
- (ii) Find  $P(1 \le X < 3)$
- (iii) Find E(X), the mean of X.
- Q.13. A and B are independent events such that  $P(A \cap \overline{B}) = \frac{1}{4}$  and  $P(\overline{A} \cap B) = \frac{1}{6}$ . Find P(A) and P(B).
- Q.14. A pair of dice is thrown simultaneously. If X denotes the absolute difference of numbers obtained on the pair of dice, then find the probability distribution of x?
- Q.15. There are two coins. One of them is a biased coin such that

P(Head) : P(tail) is 1 : 3 and the other is a fair coin. A coin is selected at random and tossed once. If the coin showed head, then find the probability that it is a biased coin.

- Q.16. Two numbers are selected from first six even natural numbers at random without replacement. If X denotes the greater of two numbers selected, find the probability distribution of X.
- Q.17. A fair coin and an unbiased die are tossed. Let A be the event "Head appears on the coin" and B' be the event, "3 comes on the die". Find whether A and B are independent or not.

## FIVE MARKS QUESTIONS

- Q.1. By examining the chest X-ray, the probability that TB is detected when a person is actually suffering is 0.99. The probability of a healthy person diagnosed to have TB is 0.001. In a certain city, 1 in 1000 people suffers from TB. A person is selected at random and is diagnosed to have TB. What is the probability that he actually has TB ?
- Q.2. Three persons *A*, *B* and *C* apply for a job of Manager in a private company. Chances of their selection (*A*, *B* and *C*) are in the ratio 1 : 2 : 4. The probabilities that *A*, *B* and *C* can introduce charges to improve profits of the company are 0.8, 0.5 and 0.3 respectively. If the change doesn't take place, find the probability that it is due to the appointment of *C*.
- Q.3. A letter is known to have come either from TATANAGAR or from CALCUTTA. On the envelope, just two consecutive letters TA are visible. What is the probability that the letter came from TATANAGAR.
- Q.4. The probability distribution of a random variable X is given as under :

$$P(X = x) = \begin{cases} kx^2 & \text{for } x = 1, 2, 3\\ 2kx & \text{for } x = 4, 5, 6\\ 0 & \text{Otherwise} \end{cases}$$

where k is a constant. Calculate

(*i*) 
$$E(X)$$
 (*ii*)  $E(3X^2)$  (*iii*)  $P(X \ge 4)$ 

- Q.5. Three critics review a book. Odds in favour of the book are 5 : 2, 4 : 3 and 3 : 4 respectively for the three critics. Find the probability that the majority are in favour of the book.
- Q.6. Two numbers are selected at random (without replacement) from positive integers 2, 3, 4, 5, 6, 7. Let X denotes the larger of the two numbers obtained. Find the mean of the probability distribution of X.
- Q.7. An urn contains five balls. Two balls are drawn and are found to be white. What is the probability that all the balls are white?
- Q.8. Two cards are drawn from a well shuffled pack of 52 cards. Find the mean and variance for the number of face cards obtained.
- Q.9. A card from a pack of 52 cards is lost. From the remaining cards of the pack, two cards are drawn at random and are found to be both clubs. Find the possibility of the lost card being of club.
- Q.10. Bag I contains 3 red and 4 black balls and Bag II contains 4 red and 5 black balls. One ball is transferred from Bag I to Bag II and then a ball is drawn from Bag II at random. The ball so drawn is found to be red in colour. Find the probability that the transferred ball is black.

## CASE STUDY QUESTIONS

Q.1. There are different types of Yoga which involve the usage of different poses of Yoga Asanas, Meditation and Pranayam as shown in the figure below:



The Venn Diagram below represents the probabilities of three different types of Yoga, A, B and C performed by the people of a society. Further it is given that probability of a member performing type C Yoga is 0.44.



On the basis of the above information, answer the following questions:

- (i) Find the value of x.
- (ii) Find the value of y.
- (iii) (a) Find  $P\left(\frac{C}{B}\right)$

OR

- (iii) (b) Find the probability that a randomly selected person of the society does Yoga of type A or B but not C.
- Q.2. Recent studies suggest that roughly 12% of the world population is left handed.



Depending upon the parents, the chances of havig a left handed child are as follows:

A : When both father and mother are left handed

Chances of left handed child is 24%

B : When father is right handed and mother is left handed:

Chances of left handed child is 22%

 ${\sf C}$  : When father is left handed and mother is right handed:

Chances of left handed child is 17%

D: When both father and mother are right handed:

Chances of left handed child is 9%

Assuming that  $P(A) = P(B) = P(C) = P(D) = \frac{1}{4}$  and L denotes the event that child is left

handed.

Based on the above information, answer the following questions:

(i) Find P(L/C)

- (ii) Find  $P(\overline{L}/A)$
- (iii) (a) Find P(A/L)

OR

(b) Find the probability that a randomly selected child is left handed given that exactly one of the parents is left handed.

Q.3 An octagonal prism is a three-dimensional polyhedron bounded by two octagonal bases and eight rectangular side faces. It has 24 edges and 16 vertices.



The prism is rolled along the rectangular faces and number on the bottom face (touching the ground) is noted. Let X denote the number obtianed on the bottom face and the following table give the probability distribution of X.

X :	1	2	3	4	5	6	7	8
P(X):	Р	2р	2р	р	2р	p <sup>2</sup>	2p <sup>2</sup>	7p² + p

Based on the above information, answer the following questions:

- (i) Find the value of p
- (ii) FInd P(X > 6)
- (iii) (a) Find P(x = 3m), where m is a natural number

OR

- (iii) (b) Find the mean E(X)
- Q.4. In a birthday party, a magician was being invited by a parent and he had 3 bags that contain number of red and white balls as follows:

Bag 1 contains : 3 red balls, Bag 2 contains : 2 white balls and 1 Red ball

Bag 3 contains : 3 white balls

The probability that the bag i will be chosen by the magician and a ball is selected from

it is 
$$\frac{i}{6}$$
,  $i = 1, 2, 3$ .

Based on the above information, answer the following questions.

- (a) What is the probability that a red ball is selected by the magician?
- (b) What is the probability that a white ball is selected by the magician?
- (c) Given that the magician selects the white ball, what is the probability that the ball was from Bag 2.

Q.5. In an office three employees Vinay, Sonia and Iqbal process incoming copies of a certain form. Vinay process 50% of the forms. Sonia processes 20% and Iqbal the remaining 30% of the forms. Vinay has an error rate of 0.06, Sonia has an error rate of 0.04 and Iqbal has an error rate of 0.03.



Based on the above information answer the following :

- (*i*) Find the conditional probability that an error is committed in processing given that Sonia processed the form?
- (ii) What is probability that Sonia processed the form and committed an error?
- (iii) What is total probability of committing an error in processing the form?

#### SELF ASSESSMENT-1

# EACH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

If A and B are independent events such that P(A) = 0.4, P(B) = x and P(A ∪ B) = 0.5, then x = ?



- 2. If P(A) = 0.8, P(B) = 0.5 and P(B/A) = 0.4, then P(A/B) is
  - (a) 0.32 (b) 0.64
  - (c) 0.16 (d) 0.25
- 3. A couple has two children. What is the probability that both are boys if it is known that one of them is a boy?

(a)	$\frac{1}{3}$	(b)	$\frac{2}{3}$
(c)	$\frac{3}{4}$	(d)	$\frac{1}{4}$

4. The random variable X has a probability distribution P(X) of the following form, where 'k' is some number.

$$P(X = x) = \begin{cases} k, & \text{if } x = 0\\ 2k, & \text{if } x = 1\\ 3k, & \text{if } x = 2\\ 0 & \text{otherwise} \end{cases}$$

Determine the value of k.

- 5. If two events are independent, then
  - (a) they must be mutually exclusive
  - (b) the sum of their probabilities must be equal to 1
  - (c) (a) and (b) both are correct
  - (d) none of the above is correct

## SELF ASSESSMENT-2

# EAH OF THE FOLLOWING MCQ HAS ONE OPTION CORRECT, CHOOSE THE CORRECT ALTERNATIVE.

1. Two cards are drawn from a well shuffled deck of 52 playing cards with replacement. The probability that both cards are queens is

(a) 
$$\frac{1}{13} \times \frac{1}{13}$$
 (b)  $\frac{1}{13} + \frac{1}{13}$   
(c)  $\frac{1}{13} \times \frac{1}{17}$  (d)  $\frac{1}{13} \times \frac{4}{51}$ 

2. The probability distribution of a discrete random variable X is given below:

Х	2	3	4	5
P(X = x)	$\frac{5}{k}$	$\frac{7}{k}$	$\frac{9}{k}$	$\frac{11}{k}$

The values of k is

(a)	8	(b)	16
(c)	32	(d)	48

3. Three persons A, B and C fire at a target in turn, starting with A. Their probability of hitting the target are 0.4, 0.3 and 0.2 respectively. The probability of two hits is

(a)	0.024	(b)	0.188
• •			

- (c) 0.336 (d) 0.452
- 4. If  $4P(A) = 6P(B) = 10P(A \cap B) = 1$ , then P(B/A) = ?

(a)	$\frac{2}{5}$	(b)	$\frac{3}{5}$
(c)	7 10	(d)	<u>19</u> 60

5. A letter is known to have come either from LONDON or CLIFTON; on the postmark only the two consecutive letters ON are legible. The probability that it came from LONDON is

(a)	<u>5</u> 17	(b)	12 17
(c)	<u>17</u> 30	(d)	$\frac{3}{5}$

#### ANSWER

## **One Mark Questions**

1. (d)	<u>1</u> 70	2. (d)0.9	6	3.	(a) $\frac{2}{9}$	4.	(C)
5. (c)	$\frac{7}{8}$	6. (c) <del>31</del> 32	-	7.	(c) $\frac{4}{9}$	8.	(d) $\frac{1}{2}$
9. (c)	$\frac{1}{3}$	10. (d) $\frac{2}{3}$		11.	(b) $1-P(A/\overline{B})$	12.	(d) P(A) = P(B)
13. (d)	0.14	14. (c) Bo	th (a) & (b)	15.	(c) $\frac{1}{5}$	16.	(a)
17. (d)		18. (a)		19.	(a)	20.	(a)

### **Two Marks Questions**

1. (i) 1 (ii)	0 2.	<u>4</u> 15	3.	<u>15</u> 29	4	$\frac{4}{7}$
5. $\frac{10}{9}$	7.	1 70	8.	$\frac{1}{36}$	g	$\frac{3}{7}$
10. $\frac{23}{24}$	11.	<u>35</u> 18	12.	0.6		
		Three M	larks Ques	tions		
1. $\frac{3}{4}$	2.	$P(A) = \frac{1}{5} a$	ind <i>P</i> ( <i>B</i> ) =	1 6 or <i>P</i> (A) =	$=\frac{5}{6}$ and P	$(B)=\frac{4}{5}$
3. $\frac{1}{2} \frac{x}{P(x)} \frac{0}{1}$	$\frac{1}{9} \frac{1}{6} \frac{1}{16} \frac{1}{16}$ 4.	(i) $\frac{1}{30}$ (ii) $\frac{1}{1}$	$\frac{3}{0}$ (iii) $\frac{2}{3}$		5	$5 \frac{91}{54}$
$6.  \frac{3}{4}$	7.	5 9	8.	$\frac{7}{50}$	ç	$\frac{1}{2}$
10. <mark>9</mark> 7	11.	4.2₹	12.	(i) k = 1 (ii	$\frac{5}{6}$ (iii) $\frac{5}{3}$	
13. $\frac{1}{3}$ and $\frac{1}{4}$ ,	$\frac{3}{4}$ and $\frac{2}{3}$					
14.	< 0	1	2	3	4	5
P(	X) $\frac{6}{36}$	$\frac{10}{36}$	$\frac{8}{36}$	$\frac{6}{36}$	$\frac{4}{36}$	$\frac{2}{36}$
15. <sup>1</sup> / <sub>3</sub>						
16.	۲ 4	6	8	10	12	1
P()	X) $\frac{1}{15}$	2 15	3 15	4 15	5 15	
17. Yes,	A and B are inde	pendent.				

Five Marks Questions1.  $\frac{110}{221}$ 2.  $\frac{7}{10}$ 3.  $\frac{7}{11}$ 4. (i) 4.31, (ii) 61.9, (iii)  $\frac{15}{22}$ 

[Class XII : Maths]

251

5. 
$$\frac{209}{343}$$
  
6.  $\overline{x} = \frac{17}{3}$   
7.  $\frac{1}{2}$   
8.  $\overline{x} = \frac{6}{13}$ ,  $\sigma^2 = \frac{60}{169}$   
9.  $\frac{11}{50}$   
10.  $\frac{16}{31}$ 

## CASE STUDY QUESTIONS

1. (i) x = 0.23 (ii) y = 0.04 (iii) (a) 
$$\frac{23}{36}$$
 or (b) 0.46

2. (i) 
$$P(L/C) = 0.17$$
, (ii)  $P(\overline{L}/A) = 0.76$  (iii) (a)  $P(A/L) = \frac{1}{3}$  or (b) 0.39

3. (i)  $P = \frac{1}{10}$  (ii)  $P(x > 6) = \frac{19}{100}$ 

(iii) (a) 
$$\frac{21}{100}$$
 or (b)  $E(x) = 4.06$ 

- 4. (a)  $\frac{5}{18}$  (b)  $\frac{13}{18}$  (c)  $\frac{4}{13}$
- 5. (*i*) 0.04 (*ii*) 0.008 (*iii*) 0.047

## SELF ASSESSMENT-1

1. (c) 2. (b) 3. (a) 4. 
$$k = \frac{1}{6}$$
 5. (d)  
**SELF ASSESSMENT-2**  
1. (a) 2. (c) 3. (b) 4. (a) 5. (b)

## PRACTICE PAPER - I (CBSE - 2023 SAMPLE PAPER)

Session 2023-24

Mathematics (Code-041)

#### Time: 3 hours

Maximum marks: 80

#### **General Instructions:**

- 1. This Question paper contains five sections A, B, C, D and E. Each section is compulsory. However, there are internal choices in some questions.
- 2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
- 3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
- 4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
- 5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.

6. Section E has 3 source based/case based/passage based/integrated units of assessment of 4 marks each with sub-parts.

## Section -A

(Multiple Choice Questions)

Each question carries 1 mark

Q1. If 
$$A = \begin{bmatrix} a_{ij} \end{bmatrix}$$
 is a square matrix of order 2 such that  $a_{ij} = \begin{cases} 1, \text{ when } i \neq j \\ 0, \text{ when } i = j \end{cases}$ , then  $A^2$  is  
(a)  $\begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix}_{2\times 2}$  (b)  $\begin{bmatrix} 1 & 1 \\ 0 & 0 \end{bmatrix}_{2\times 2}$  (c)  $\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}_{2\times 2}$  (d)  $\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}_{2\times 2}$ 

**Q2.** If A and B are invertible square matrices of the same order, then which of the following is not correct?

(a) $\left  \mathbf{AB}^{-1} \right  = \frac{ \mathbf{A} }{ \mathbf{B} }$	(b) $\left  \left( AB \right)^{-1} \right  = \frac{1}{ A   B }$
(c) $(AB)^{-1} = B^{-1}A^{-1}$	(d) $(A+B)^{-1} = B^{-1} + A^{-1}$

Q3. If the area of the triangle with vertices (-3, 0), (3, 0) and (0, k) is 9 sq units, then the value/s of k will

(a) 9 (b) 
$$\pm 3$$
 (c) -9 (d) 6  
Q4. If  $f(x) = \begin{cases} \frac{kx}{|x|}, & \text{if } x < 0 \\ 3, & \text{if } x \ge 0 \end{cases}$  is continuous at  $x = 0$ , then the value of  $k$  is  
(a) -3 (b) 0 (c) 3 (d) any real number

[Class XII : Maths]

be

Q5. The lines  $\vec{r} = \hat{i} + \hat{j} - \hat{k} + \lambda \left(2\hat{i} + 3\hat{j} - 6\hat{k}\right)$  and  $\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu \left(6\hat{i} + 9\hat{j} - 18\hat{k}\right)$ ; (where  $\lambda \& \mu$  are scalars) are

(a) coincident (b) skew (c) intersecting (d) parallel Q6. The degree of the differential equation  $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^3 = \left(\frac{d^2y}{dx^2}\right)^2 is$ (a) 4 (b)  $\frac{3}{2}$  (c) 2 (d) Not defined

Q7. The corner points of the bounded feasible region determined by a system of linear constraints are (0,3),(1,1) and (3,0). Let Z = px + qy, where p, q > 0. The condition on p and q so that the minimum of Z occurs at (3,0) and (1,1) is

(a) 
$$p = 2q$$
 (b)  $p = \frac{q}{2}$  (c)  $p = 3q$  (d)  $p = q$ 

**Q8.** ABCD is a rhombus whose diagonals intersect at **E**. Then  $\overrightarrow{EA} + \overrightarrow{EB} + \overrightarrow{EC} + \overrightarrow{ED}$  equals to

(a) 
$$\vec{0}$$
 (b)  $\overrightarrow{AD}$  (c)  $2\overrightarrow{BD}$  (d)  $2\overrightarrow{AD}$ 

**Q9.** For any integer *n*, the value of  $\int_{-\pi}^{\pi} e^{\cos^2 x} \sin^3(2n+1) x \, dx$  is

Q10. The value of |A|, if  $A = \begin{bmatrix} 0 & 2x - 1 & \sqrt{x} \\ 1 - 2x & 0 & 2\sqrt{x} \\ -\sqrt{x} & -2\sqrt{x} & 0 \end{bmatrix}$ , where  $x \in \mathbb{R}^+$ , is

(a) 
$$(2x+1)^2$$
 (b) 0 (c)  $(2x+1)^3$  (d)  $(2x-1)^2$ 

Q11. The feasible region corresponding to the linear constraints of a Linear Programming Problem is given below.



Which of the following is not a constraint to the given Linear Programming Problem?

(a)  $x + y \ge 2$  (b)  $x + 2y \le 10$  (c)  $x - y \ge 1$  (d)  $x - y \le 1$ 

Q12. If  $\vec{a} = 4\hat{i} + 6\hat{j}$  and  $\vec{b} = 3\hat{j} + 4\hat{k}$ , then the vector form of the component of  $\vec{a}$  along  $\vec{b}$  is

(a) 
$$\frac{18}{5} (3\hat{i} + 4\hat{k})$$
 (b)  $\frac{18}{25} (3\hat{j} + 4\hat{k})$  (c)  $\frac{18}{5} (3\hat{i} + 4\hat{k})$  (d)  $\frac{18}{25} (4\hat{i} + 6\hat{j})$ 

Q13. Given that A is a square matrix of order 3 and |A| = -2, then |adj(2A)| is equal to

(a) 
$$-2^6$$
 (b)  $+4$  (c)  $-2^8$  (d)  $2^8$ 

Q14. A problem in Mathematics is given to three students whose chances of solving it are  $\frac{1}{2}, \frac{1}{3}, \frac{1}{4}$ 

respectively. If the events of their solving the problem are independent then the probability that the problem will be solved, is

(a) 
$$\frac{1}{4}$$
 (b)  $\frac{1}{3}$  (c)  $\frac{1}{2}$  (d)  $\frac{3}{4}$ 

Q15. The general solution of the differential equation ydx - xdy = 0; (Given x, y > 0), is of the form

(a) 
$$xy = c$$
 (b)  $x = cy^2$  (c)  $y = cx$  (d)  $y = cx^2$ ;

(Where 'c' is an arbitrary positive constant of integration)

Q16. The value of  $\lambda$  for which two vectors  $2\hat{i} - \hat{j} + 2\hat{k}$  and  $3\hat{i} + \lambda\hat{j} + \hat{k}$  are perpendicular is (a) 2 (b) 4 (c) 6 (d) 8

Q17. The set of all points where the function f(x) = x + |x| is differentiable, is

(a) 
$$(0,\infty)$$
 (b)  $(-\infty,0)$  (c)  $(-\infty,0) \cup (0,\infty)$  (d)  $(-\infty,\infty)$ 

**Q18.** If the direction cosines of a line are  $<\frac{1}{c}, \frac{1}{c}, \frac{1}{c} >$ , then

(a) 0 < c < 1 (b) c > 2 (c)  $c = \pm \sqrt{2}$  (d)  $c = \pm \sqrt{3}$ 

#### ASSERTION-REASON BASED QUESTIONS

In the following questions, a statement of Assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.

- (a) Both (A) and (R) are true and (R) is the correct explanation of (A).
- (b) Both (A) and (R) are true but (R) is not the correct explanation of (A).
- (c) (A) is true but (R) is false.
- (d) (A) is false but (R) is true.

Q19. Let f(x) be a polynomial function of degree 6 such that  $\frac{d}{dx}(f(x)) = (x-1)^3(x-3)^2$ , then

**ASSERTION (A):** f(x) has a minimum at x = 1.

**REASON (R):** When  $\frac{d}{dx}(f(x)) < 0$ ,  $\forall x \in (a - h, a)$  and  $\frac{d}{dx}(f(x)) > 0$ ,  $\forall x \in (a, a + h)$ ; where 'h' is an infinitesimally small positive quantity, then f(x) has a minimum at x = a, provided f(x) is continuous at x = a. Q20. ASSERTION (A): The relation  $f:\{1,2,3,4\} \rightarrow \{x,y,z,p\}$  defined by  $f=\{(1,x),(2,y),(3,z)\}$  is a bijective function.

**REASON (R)**: The function  $f: \{1,2,3\} \rightarrow \{x,y,z,p\}$  such that  $f = \{(1,x),(2,y),(3,z)\}$  is one-one.

#### Section -B

[This section comprises of very short answer type questions (VSA) of 2 marks each]

Q21. Find the value of 
$$\sin^{-1}\left(\cos\left(\frac{33\pi}{5}\right)\right)$$
.

Find the domain of  $\sin^{-1}(x^2-4)$ .

Q22. Find the interval/s in which the function  $f: \mathbb{R} \to \mathbb{R}$  defined by  $f(x) = xe^x$ , is increasing.

Q23. If 
$$f(x) = \frac{1}{4x^2 + 2x + 1}$$
;  $x \in \mathbb{R}$ , then find the maximum value of  $f(x)$ .

OR

Find the maximum profit that a company can make, if the profit function is given by

 $P(x) = 72 + 42x - x^2$ , where x is the number of units and P is the profit in rupees.

- **Q24.** Evaluate :  $\int_{-1}^{1} \log \left(\frac{2-x}{2+x}\right) dx$ .
- Q25. Check whether the function  $f : \mathbb{R} \to \mathbb{R}$  defined by  $f(x) = x^3 + x$ , has any critical point/s or not? If yes, then find the point/s.

#### <u>Section – C</u> [This section comprises of short answer type questions (SA) of 3 marks each]

Q26. Find: 
$$\int \frac{2x^2+3}{x^2(x^2+9)} dx; x \neq 0.$$

Q27. The random variable X has a probability distribution P(X) of the following form, where 'k' is some real number:

$$P(X) = \begin{cases} k, & \text{if } x=0\\ 2k, & \text{if } x=1\\ 3k, & \text{if } x=2\\ 0, & \text{otherwise} \end{cases}$$

(i) Determine the value of k.

(ii) Find P(X < 2).

(iii) Find 
$$P(X > 2)$$
.  
Q28. Find :  $\int \sqrt{\frac{x}{1-x^3}} dx$ ;  $x \in (0,1)$ .  
OR  
Evaluate:  $\int_0^{\frac{\pi}{4}} \log(1 + \tan x) dx$ .  
Q29. Solve the differential equation:  $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy$ ,  $(y \neq 0)$ .  
OR

Solve the differential equation:  $(\cos^2 x)\frac{dy}{dx} + y = \tan x; \quad \left(0 \le x < \frac{\pi}{2}\right).$ 

Q30. Solve the following Linear Programming Problem graphically:

Minimize: z = x + 2y,

subject to the constraints:  $x + 2y \ge 100$ ,  $2x - y \le 0$ ,  $2x + y \le 200$ ,  $x, y \ge 0$ .

OR

Solve the following Linear Programming Problem graphically: Maximize: z = -x + 2y,

subject to the constraints:  $x \ge 3, x + y \ge 5, x + 2y \ge 6, y \ge 0$ .

Q31. If 
$$(a+bx)e^{\frac{y}{x}} = x$$
 then prove that  $x\frac{d^2y}{dx^2} = \left(\frac{a}{a+bx}\right)^2$ .

#### Section -D

#### [This section comprises of long answer type questions (LA) of 5 marks each]

Q32. Make a rough sketch of the region  $\{(x, y): 0 \le y \le x^2 + 1, 0 \le y \le x + 1, 0 \le x \le 2\}$  and find the

area of the region, using the method of integration.

Q33. Let  $\mathbb{N}$  be the set of all natural numbers and R be a relation on  $\mathbb{N} \times \mathbb{N}$  defined by

 $(a,b) R(c,d) \Leftrightarrow ad = bc$  for all  $(a,b), (c,d) \in \mathbb{N} \times \mathbb{N}$ . Show that **R** is an equivalence relation on

 $\mathbb{N} \times \mathbb{N}$ . Also, find the equivalence class of (2,6), i.e., [(2,6)].

#### OR

Show that the function  $f : \mathbb{R} \to \{x \in \mathbb{R} : -1 < x < 1\}$  defined by  $f(x) = \frac{x}{1+|x|}, x \in \mathbb{R}$  is one-one and onto function.

Q34. Using the matrix method, solve the following system of linear equations :

$$\frac{2}{x} + \frac{3}{y} + \frac{10}{z} = 4, \ \frac{4}{x} - \frac{6}{y} + \frac{5}{z} = 1, \ \frac{6}{x} + \frac{9}{y} - \frac{20}{z} = 2.$$

Q35. Find the coordinates of the image of the point (1, 6, 3) with respect to the line

 $\vec{r} = (\hat{j} + 2\hat{k}) + \lambda(\hat{i} + 2\hat{j} + 3\hat{k});$  where ' $\lambda$ ' is a scalar. Also, find the distance of the image from the y - axis.

OR

An aeroplane is flying along the line  $\vec{r} = \lambda (\hat{i} - \hat{j} + \hat{k})$ ; where ' $\lambda$ ' is a scalar and another aeroplane is flying along the line  $\vec{r} = \hat{i} - \hat{j} + \mu (-2\hat{j} + \hat{k})$ ; where ' $\mu$ ' is a scalar. At what points on the lines should they reach, so that the distance between them is the shortest? Find the shortest possible distance between them.

#### Section -E

[This section comprises of 3 case- study/passage based questions of 4 marks each with sub parts. The first two case study questions have three sub parts (i), (ii), (iii) of marks 1,1,2 respectively. The third case study question has two sub parts of 2 marks each.)

Q36. Read the following passage and answer the questions given below:

In an Office three employees Jayant, Sonia and Oliver process incoming copies of a certain form. Jayant processes 50% of the forms, Sonia processes 20% and Oliver the remaining 30% of the forms. Jayant has an error rate of 0.06, Sonia has an error rate of 0.04 and Oliver has an error rate of 0.03.

Based on the above information, answer the following questions.



(i) Find the probability that Sonia processed the form and committed an error.

(ii) Find the total probability of committing an error in processing the form.

(iii) The manager of the Company wants to do a quality check. During inspection, he selects a form at random from the days output of processed form. If the form selected at random has an error, find the probability that the form is **not** processed by Jayant.

OR

- (iii) Let E be the event of committing an error in processing the form and let  $E_1, E_2$  and  $E_3$  be the events that Jayant, Sonia and Oliver processed the form. Find the value of  $\sum_{i=1}^{3} P(E_i | E)$ .
- Q37. Read the following passage and answer the questions given below:

Teams A, B, C went for playing a tug of war game. Teams A, B, C have attached a rope to a metal ring and is trying to pull the ring into their own area.

Team A pulls with force  $F_1 = 6\hat{i} + 0\hat{j} kN$ ,

Team **B** pulls with force  $F_2 = -4\hat{i} + 4\hat{j} kN$ ,

Team C pulls with force  $F_3 = -3\hat{i} - 3\hat{j} kN$ ,



- (i) What is the magnitude of the force of Team A?
- (ii) Which team will win the game?
- (iii) Find the magnitude of the resultant force exerted by the teams.

## OR

(iii) In what direction is the ring getting pulled?

Q38. Read the following passage and answer the questions given below:

The relation between the height of the plant ('y' in cm) with respect to its exposure to the sunlight

is governed by the following equation  $y = 4x - \frac{1}{2}x^2$ , where 'x' is the number of days exposed to the





- (ii) Does the rate of growth of the plant increase or decrease in the first three days? What will be the height of the plant after 2 days?
- Find the rate of growth of the plant with respect to the number of days exposed to the sunlight.

## PRACTICE PAPER - I (CBSE - 2023 SAMPLE PAPER) CLASS XII

## **MATHEMATICS (CODE-041)**

## SECTION: A (Solution of MCQs of 1 Mark each)

Q no.	ANS	HINTS/SOLUTION
1	(d)	$A = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}, A^2 = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}.$
2	(d)	$(A+B)^{-1}=B^{-1}+A^{-1}.$
3	(b)	Area = $\begin{vmatrix} 1 \\ -3 & 0 & 1 \\ 3 & 0 & 1 \\ 0 & k & 1 \end{vmatrix}$ , given that the area = 9 squnit.
		$\Rightarrow \pm 9 = \frac{1}{2} \begin{vmatrix} -3 & 0 & 1 \\ 3 & 0 & 1 \\ 0 & k & 1 \end{vmatrix}; \text{ expanding along } C_2, \text{ we get } \Rightarrow k = \pm 3.$
4	(a)	Since, $f$ is continuous at $x = 0$ ,
		therefore, $L.H.L = R.H.L = f(0) = a$ finite quantity.
		$\lim_{x \to 0^{-}} f(x) = \lim_{x \to 0^{+}} f(x) = f(0)$
		$\Rightarrow \lim_{x\to 0^-} \frac{-kx}{x} = \lim_{x\to 0^+} 3 = 3 \Rightarrow k = -3.$
5	(d)	Vectors $2\hat{i}+3\hat{j}-6\hat{k} & 6\hat{i}+9\hat{j}-18\hat{k}$ are parallel and the fixed point $\hat{i}+\hat{j}-\hat{k}$ on the
		line $\vec{r} = \hat{i} + \hat{j} - \hat{k} + \lambda \left(2\hat{i} + 3\hat{j} - 6\hat{k}\right)$ does not satisfy the other line
		$\vec{r} = 2\hat{i} - \hat{j} - \hat{k} + \mu (\hat{6i} + \hat{j} - 1\hat{k});$ where $\lambda \& \mu$ are scalars.
6	(c)	The degree of the differential equation $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^3 = \left(\frac{d^2y}{dx^2}\right)^2$ is 2
7	(b)	Z = px + qy(i)
		At $(3,0)$ , $Z = 3p(ii)$ and at $(1,1)$ , $Z = p + q(iii)$
		From (ii) & (iii), $3p = p + q \Rightarrow 2p = q$ .

8	(a)	Given, <i>ABCD</i> is a rhombus whose diagonals bisect each other. $\left  \overrightarrow{EA} \right  = \left  \overrightarrow{EC} \right $ and
		$\left \overline{EB}\right  = \left \overline{ED}\right $ but since they are opposite to each other so they are of opposite signs
		$\Rightarrow \overrightarrow{EA} = -\overrightarrow{EC} \text{ and } \overrightarrow{EB} = -\overrightarrow{ED}.$
		$B \xrightarrow{E} D$ $C$ $\Rightarrow \overline{EA} + \overline{EC} = \vec{O} \dots (i) \text{ and } \overline{EB} + \overline{ED} = \vec{O} \dots (ii)$ Adding (i) and (ii), we get $\overline{EA} + \overline{EB} + \overline{EC} + \overline{ED} = \vec{O}$ .
9	(b)	$f(x) = e^{\cos^2 x} \sin^3(2n+1)x$
		$f(-x) = e^{\cos^2(-x)} \sin^3(2n+1)(-x)$
		$f(-x) = -e^{\cos^2 x} \sin^3(2n+1)x$
		f(-x) = -f(x)
		$So, \int_{-\pi}^{\pi} e^{\cos^2 x} \sin^3(2n+1)x  dx = 0$
10	(b)	Matrix A is a skew symmetric matrix of odd order. $\therefore  A  = 0$ .
11	(c)	We observe, $(0,0)$ does not satisfy the inequality $x-y \ge 1$
		So, the half plane represented by the above inequality will not contain origin
		therefore, it will not contain the shaded feasible region.
12	(b)	Vector component of $\vec{a}$ along $\vec{b} = \left(\frac{\vec{a} \cdot \vec{b}}{\left \vec{b}\right ^2}\right) \vec{b} = \frac{18}{25} \left(3\hat{j} + 4\hat{k}\right).$
13	(d)	$ adj(2A)  =  (2A) ^2 = (2^3 A )^2 = 2^6 A ^2 = 2^6 \times (-2)^2 = 2^8.$
14	(d)	Method 1:
		Let A, B, C be the respective events of solving the problem. Then, $P(A) = \frac{1}{2}$ , $P(B) = \frac{1}{3}$
		and $P(C) = \frac{1}{4}$ . Here, $A, B, C$ are independent events.
		Problem is solved if at least one of them solves the problem.
		Required probability is = $P(A \cup B \cup C) = 1 - P(\overline{A})P(\overline{B})P(\overline{C})$

		$=1-\left(1-\frac{1}{2}\right)\left(1-\frac{1}{3}\right)\left(1-\frac{1}{4}\right)=1-\frac{1}{4}=\frac{3}{4}.$
		Method 2:
		The problem will be solved if one or more of them can solve the problem. The probability is
		$P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC}) + P(\overline{ABC})$
		$=\frac{1}{2}\cdot\frac{2}{3}\cdot\frac{3}{4}+\frac{1}{2}\cdot\frac{1}{3}\cdot\frac{3}{4}+\frac{1}{2}\cdot\frac{2}{3}\cdot\frac{1}{4}+\frac{1}{2}\cdot\frac{1}{3}\cdot\frac{3}{4}+\frac{1}{2}\cdot\frac{2}{3}\cdot\frac{1}{4}+\frac{1}{2}\cdot\frac{1}{3}\cdot\frac{1}{4}+\frac{1}{2}\cdot\frac{1}{3}\cdot\frac{1}{4}+\frac{1}{2}\cdot\frac{1}{3}\cdot\frac{1}{4}=\frac{3}{4}.$
		Method 3:
		Let us think quantitively. Let us assume that there are 100 questions given to A. A
		solves $\frac{1}{2} \times 100 = 50$ questions then remaining 50 questions is given to <i>B</i> and <i>B</i> solves
		$50 \times \frac{1}{3} = 16.67$ questions. Remaining $50 \times \frac{2}{3}$ questions is given to C and C solves
		$50 \times \frac{2}{3} \times \frac{1}{4} = 8.33$ questions.
		Therefore, number of questions solved is $50 + 16.67 + 8.33 = 75$ .
		So, required probability is $\frac{75}{100} = \frac{3}{4}$ .
15	(c)	Method 1:
		$ydx - xdy = 0 \Rightarrow \frac{ydx - xdy}{y^2} = 0 \Rightarrow d\left(\frac{x}{y}\right) = 0 \Rightarrow x = \frac{1}{c}y \Rightarrow y = cx.$
		Method 2:
		$ydx - xdy = 0 \Rightarrow ydx = xdy \Rightarrow \frac{dy}{y} = \frac{dx}{x}$ ; on integrating $\int \frac{dy}{y} = \int \frac{dx}{x}$
		$\log  v  = \log  x  + \log  c $
		since x, v, c > 0, we write $\log v = \log x + \log c \implies v = cx$ .
16	(d)	Dot product of two mutually perpendicular vectors is zero.
1.1	(-)	$\Rightarrow 2 \times 3 + (-1)\lambda + 2 \times 1 = 0 \Rightarrow \lambda = 8.$
17	(c)	Method 1:
		$f(x) = x +  x  = \begin{cases} 2x, x \ge 0 & Y \\ 0, x < 0 & A \end{cases}$
		*
		$y = 2x, x \ge 0$
		$X' \leftarrow \frac{y=0, x<0}{O} \longrightarrow X$
		v'
		There is a sharp corner at $y = 0$ so $f(y)$ is not differentiable at $y = 0$
		There is a sharp corner at $x = 0$ , so $f(x)$ is not differentiable at $x = 0$ .
		Method 2:

		Lf'(0) = 0 & Rf'(0) = 2;  so, the function is not differentiable at  x = 0 For $x \ge 0, f(x) = 2x$ (linear function) & when $x < 0, f(x) = 0$ (constant function)
		Hence $f(x)$ is differentiable when $x \in (-\infty, 0) \cup (0, \infty)$ .
18	(d)	We know, $l^2 + m^2 + n^2 = 1 \Rightarrow \left(\frac{1}{c}\right)^2 + \left(\frac{1}{c}\right)^2 + \left(\frac{1}{c}\right)^2 = 1 \Rightarrow 3\left(\frac{1}{c}\right)^2 = 1 \Rightarrow c = \pm\sqrt{3}.$
19	(a)	$\frac{d}{dx}(f(x)) = (x-1)^3 (x-3)^2$ Assertion : $f(x)$ has a minimum at $x = 1$ is true as $\frac{d}{dx}(f(x)) < 0, \forall x \in (1-h,1) \text{ and } \frac{d}{dx}(f(x)) > 0, \forall x \in (1,1+h); \text{ where,}$ 'h' is an infinitesimally small positive quantity, which is in accordance with the Reason statement.
20	(d)	Assertion is false. As element 4 has no image under $f$ , so relation $f$ is not a function. Reason is true. The given function $f : \{1,2,3\} \rightarrow \{x, y, z, p\}$ is one – one, as for each $a \in \{1,2,3\}$ , there is different image in $\{x, y, z, p\}$ under $f$ .

## Section -B

[This section comprises of solution of very short answer type questions (VSA) of 2 marks each]

21	$\sin^{-1}\left(\cos\left(\frac{33\pi}{5}\right)\right) = \sin^{-1}\cos\left(6\pi + \frac{3\pi}{5}\right) = \sin^{-1}\cos\left(\frac{3\pi}{5}\right) = \sin^{-1}\sin\left(\frac{\pi}{2} - \frac{3\pi}{5}\right)$	1
	$=\frac{\pi}{2}-\frac{3\pi}{5}=-\frac{\pi}{10}.$	1
21 OR	$-1 \le \left(x^2 - 4\right) \le 1 \Longrightarrow 3 \le x^2 \le 5 \Longrightarrow \sqrt{3} \le  x  \le \sqrt{5}$	1
	$\Rightarrow x \in \left[-\sqrt{5}, -\sqrt{3}\right] \cup \left[\sqrt{3}, \sqrt{5}\right].$ So, required domain is $\left[-\sqrt{5}, -\sqrt{3}\right] \cup \left[\sqrt{3}, \sqrt{5}\right].$	1
22	$f(x) = xe^x \Rightarrow f'(x) = e^x(x+1)$	1
	When $x \in [-1,\infty), (x+1) \ge 0$ & $e^x > 0 \Rightarrow f'(x) \ge 0 \therefore f(x)$ increases in this interval.	1
	or, we can write $f(x) = xe^x \Rightarrow f'(x) = e^x(x+1)$	$\frac{1}{2}$
	For $f(x)$ to be increasing, we have $f'(x) = e^x(x+1) \ge 0 \Rightarrow x \ge -1$ as $e^x > 0, \forall x \in \mathbb{R}$	1
	Hence, the required interval where $f(x)$ increases is $[-1,\infty)$ .	$\frac{1}{2}$
23	Method 1: $f(x) = \frac{1}{4x^2 + 2x + 1}$ ,	

Let 
$$g(x) = 4x^2 + 2x + 1 = 4\left(x^2 + 2x\frac{1}{4} + \frac{1}{16}\right) + \frac{3}{4} = 4\left(x + \frac{1}{4}\right)^2 + \frac{3}{4} \ge \frac{3}{4}$$
  
 $\therefore$  maximum value of  $f(x) = \frac{4}{3}$ .  
Method 2:  $f(x) = \frac{1}{4x^2 + 2x + 1}$ , let  $g(x) = 4x^2 + 2x + 1$   
 $\Rightarrow \frac{d}{dx}(g(x)) = g'(x) = 8x + 2$  and  $g'(x) = 0$  at  $x = -\frac{1}{4}$  also  $\frac{d^2}{dx^2}(g(x)) = g''(x) = 8 > 0$   
 $\Rightarrow g(x)$  is minimum when  $x = -\frac{1}{4}$  so,  $f(x)$  is maximum at  $x = -\frac{1}{4}$   
 $\therefore$  maximum value of  $f(x) = f\left(-\frac{1}{4}\right) = \frac{1}{4\left(-\frac{1}{4}\right)^2 + 2\left(-\frac{1}{4}\right) + 1} = \frac{4}{3}$ .  
Method 3:  $f(x) = \frac{1}{4x^2 + 2x + 1}$   
On differentiating w.r.t x , we get  $f'(x) = \frac{-(8x + 2)}{(4x^2 + 2x + 1)^2}$  ....(*i*)  
For maxima or minima, we put  $f'(x) = 0 \Rightarrow 8x + 2 = 0 \Rightarrow x = -\frac{1}{4}$ .  
Again, differentiating equation (i) w.r.t x, we get  
 $f''(x) = -\left\{ \frac{(4x^2 + 2x + 1)^2(8) - (8x + 2)2 \times (4x^2 + 2x + 1)(8x + 2)}{(4x^2 + 2x + 1)^4} \right\}$   
At  $x = -\frac{1}{4}$ ,  $f''(-\frac{1}{4}) < 0$   
 $f(x)$  is maximum at  $x = -\frac{1}{4}$ .  
 $\therefore$  maximum value of  $f(x)$  is  $f\left(-\frac{1}{4}\right) = \frac{-(8x + 2)}{(4x^2 + 2x + 1)^4}$  ....(*i*)  
For maxima or minima, we put  $f'(x) = 0 \Rightarrow 8x + 2 = 0 \Rightarrow x = -\frac{1}{4}$ .  
 $\frac{1}{2}$   
Method 4:  $f(x) = \frac{1}{4x^2 + 2x + 1}$   
On differentiating w.r.t x, we get  $f'(x) = \frac{-(8x + 2)}{(4x^2 + 2x + 1)^4}$  ....(*i*)  
For maxima or minima, we put  $f'(x) = 0 \Rightarrow 8x + 2 = 0 \Rightarrow x = -\frac{1}{4}$ .  
 $\frac{1}{2}$   
Method 4:  $f(x) = \frac{1}{4x^2 + 2x + 1}$   
On differentiating w.r.t x, we get  $f'(x) = \frac{-(8x + 2)}{(4x^2 + 2x + 1)^2}$  ....(*i*)  
For maxima or minima, we put  $f'(x) = 0 \Rightarrow 8x + 2 = 0 \Rightarrow x = -\frac{1}{4}$ .  
 $\frac{1}{2}$   
When  $x \in \left(-h - \frac{1}{4}, -\frac{1}{4}\right)$ , where 'h' is infinitesimally small positive quantity.  
 $4x < -1 \Rightarrow 8x < -2 \Rightarrow 8x + 2 < 0 \Rightarrow -(8x + 2) > 0$  and  $(4x^2 + 2x + 1)^2 > 0 \Rightarrow f'(x) > 0$ 

	and when $x \in \left(-\frac{1}{4}, -\frac{1}{4}+h\right), 4x > -1 \Rightarrow 8x > -2 \Rightarrow 8x + 2 > 0 \Rightarrow -\left(8x + 2\right) < 0$	1
	and $(4x^2 + 2x + 1)^2 > 0 \Rightarrow f'(x) < 0$ . This shows that $x = -\frac{1}{4}$ is the point of local maxima.	2
	$\therefore \text{ maximum value of } f(x) \text{ is } f\left(-\frac{1}{4}\right) = \frac{1}{4\left(-\frac{1}{4}\right)^2 + 2\left(-\frac{1}{4}\right) + 1} = \frac{4}{3}.$	$\frac{1}{2}$
23 OR	For maxima and minima, $P'(x) = 0 \Rightarrow 42 - 2x = 0$	1
	$\Rightarrow x = 21$ and $P''(x) = -2 < 0$	2
	So, $P(x)$ is maximum at $x = 21$ .	$\frac{1}{2}$
	The maximum value of $P(x) = 72 + (42 \times 21) - (21)^2 = 513$ i.e., the maximum profit is ₹ 513.	1
24	Let $f(x) = \log\left(\frac{2-x}{2+x}\right)$	-
	We have, $f(-x) = \log\left(\frac{2+x}{2-x}\right) = -\log\left(\frac{2-x}{2+x}\right) = -f(x)$	1
	So, $f(x)$ is an odd function. $\therefore \int_{-1}^{1} \log \left(\frac{2-x}{2+x}\right) dx = 0.$	1
25	$f(x) = x^3 + x$ , for all $x \in \mathbb{R}$ .	
	$\frac{d}{dx}(f(x)) = f'(x) = 3x^2 + 1; \text{ for all } x \in \mathbb{R}, \ x^2 \ge 0 \Rightarrow f'(x) > 0$	$1\frac{1}{2}$
	Hence, no critical point exists.	$\frac{1}{2}$
	Section –C	9
	[This section comprises of solution short answer type questions (SA) of 3 marks each]	
26	We have, $\frac{2x^2+3}{x^2(x^2+9)}$ . Now, let $x^2 = t$	$\frac{1}{2}$
	So, $\frac{2t+3}{t(t+9)} = \frac{A}{t} + \frac{B}{t+9}$ , we get $A = \frac{1}{3} \& B = \frac{5}{3}$	1
	$\int \frac{2x^2+3}{x^2(x^2+9)} dx = \frac{1}{3} \int \frac{dx}{x^2} + \frac{5}{3} \int \frac{dx}{x^2+9}$	$\frac{1}{2}$
	$= -\frac{1}{3x} + \frac{5}{9} \tan^{-1} \left( \frac{x}{3} \right) + c, \text{ where 'c' is an arbitrary constant of integration.}$	1
27	We have, (i) $\sum P(X_i) = 1 \Rightarrow k + 2k + 3k = 1 \Rightarrow k = \frac{1}{6}$ .	1
		1

	(ii) $P(X < 2) = P(X = 0) + P(X = 1) = k + 2k = 3k = 3 \times \frac{1}{6} = \frac{1}{2}$ .	1
	(iii) $P(X > 2) = 0.$	
28	Let $x^{\frac{3}{2}} = t \Rightarrow dt = \frac{3}{2}x^{\frac{1}{2}}dx$	$\frac{1}{2}$
	$\int \sqrt{\frac{x}{1-x^{3}}}  dx = \frac{2}{3} \int \frac{dt}{\sqrt{1-t^{2}}}$	$\frac{1}{2}$
	$=\frac{2}{3}\sin^{-1}(t)+c$	1
	$=\frac{2}{3}\sin^{-1}\left(x^{\frac{3}{2}}\right)+c, \text{ where 'c' is an arbitrary constant of integration.}$	1
28 OR	Let $I = \int_{0}^{\frac{\pi}{4}} \log_{e} (1 + \tan x) dx$ (i)	
2	$=\int_{0}^{\frac{\pi}{4}}\log_{e}\left(1+\tan\left(\frac{\pi}{4}-x\right)\right)dx,  \text{Using}, \int_{0}^{a}f(x)dx=\int_{0}^{a}f(a-x)dx$	1
	$\Rightarrow I = \int_0^{\frac{\pi}{4}} \log_e \left( 1 + \frac{1 - \tan x}{1 + \tan x} \right) dx = \int_0^{\frac{\pi}{4}} \log_e \left( \frac{2}{1 + \tan x} \right) dx = \int_0^{\frac{\pi}{4}} \log_e 2  dx - I  (\text{ Using(i)})$	1
		1
	$I = \frac{\pi}{4} \log_e 2 \Longrightarrow I = \frac{\pi}{8} \log_e 2.$	
29	$I = \frac{\pi}{4} \log_e 2 \Rightarrow I = \frac{\pi}{8} \log_e 2.$ Method 1: $y e^{\frac{x}{y}} dx = \left(x e^{\frac{x}{y}} + y^2\right) dy \Rightarrow e^{\frac{x}{y}} \left(y dx - x dy\right) = y^2 dy \Rightarrow e^{\frac{x}{y}} \left(\frac{y dx - x dy}{y^2}\right) = dy$	1
29	$I = \frac{\pi}{4} \log_e 2 \Rightarrow I = \frac{\pi}{8} \log_e 2.$ Method 1: $y e^{\frac{x}{y}} dx = \left(x e^{\frac{x}{y}} + y^2\right) dy \Rightarrow e^{\frac{x}{y}} (y dx - x dy) = y^2 dy \Rightarrow e^{\frac{x}{y}} \left(\frac{y dx - x dy}{y^2}\right) = dy$ $\Rightarrow e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = dy$	1
29	$I = \frac{\pi}{4} \log_e 2 \Rightarrow I = \frac{\pi}{8} \log_e 2.$ Method 1: $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy \Rightarrow e^{\frac{x}{y}} (ydx - xdy) = y^2 dy \Rightarrow e^{\frac{x}{y}} \left(\frac{ydx - xdy}{y^2}\right) = dy$ $\Rightarrow e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = dy$ $\Rightarrow \int e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = \int dy \Rightarrow e^{\frac{x}{y}} = y + c, \text{ where 'c' is an arbitrary constant of integration.}$	1 1 1
29	$V = \frac{\pi}{4} \log_e 2 \Rightarrow I = \frac{\pi}{8} \log_e 2.$ Method 1: $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy \Rightarrow e^{\frac{x}{y}} (ydx - xdy) = y^2 dy \Rightarrow e^{\frac{x}{y}} \left(\frac{ydx - xdy}{y^2}\right) = dy$ $\Rightarrow e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = dy$ $\Rightarrow \int e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = \int dy \Rightarrow e^{\frac{x}{y}} = y + c, \text{ where 'c' is an arbitrary constant of integration.}$ Method 2: We have, $\frac{dx}{dy} = \frac{xe^{\frac{x}{y}} + y^2}{y.e^{\frac{x}{y}}}$	1
29	$V = \frac{\pi}{4} \log_e 2 \Rightarrow I = \frac{\pi}{8} \log_e 2.$ Method 1: $ye^{\frac{x}{y}} dx = \left(xe^{\frac{x}{y}} + y^2\right) dy \Rightarrow e^{\frac{x}{y}} (ydx - xdy) = y^2 dy \Rightarrow e^{\frac{x}{y}} \left(\frac{ydx - xdy}{y^2}\right) = dy$ $\Rightarrow e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = dy$ $\Rightarrow \int e^{\frac{x}{y}} d\left(\frac{x}{y}\right) = \int dy \Rightarrow e^{\frac{x}{y}} = y + c, \text{ where 'c' is an arbitrary constant of integration.}$ Method 2: We have, $\frac{dx}{dy} = \frac{xe^{\frac{x}{y}} + y^2}{y.e^{\frac{x}{y}}}$ $\Rightarrow \frac{dx}{dy} = \frac{x}{y} + \frac{y}{e^{\frac{x}{y}}} \dots \dots$	$1$ 1 1 1 $\frac{1}{2}$

	So equation (i) becomes $v + y \frac{dv}{dy} = v + \frac{y}{e^v}$	$\frac{1}{2}$
	$\Rightarrow y \frac{dv}{dy} = \frac{y}{e^v}$	$\frac{1}{2}$
	$\Rightarrow e^{v} dv = dy$	$\frac{1}{2}$
	On integrating we get, $\int e^{y} dv = \int dy \Rightarrow e^{y} = y + c \Rightarrow e^{x/y} = y + c$	1
	where 'c' is an arbitrary constant of integration.	2
29 OR	The given Differential equation is	
	$(\cos^2 x)\frac{dy}{dx} + y = \tan x$	
	Dividing both the sides by $\cos^2 x$ , we get	
	$\frac{dy}{dx} + \frac{y}{\cos^2 x} = \frac{\tan x}{\cos^2 x}$	
	$\frac{dy}{dx} + y(\sec^2 x) = \tan x(\sec^2 x)\dots(i)$	$\frac{1}{2}$
	Comparing with $\frac{dy}{dx} + Py = Q$	
	$P = \sec^2 x, \ Q = \tan x \cdot \sec^2 x$	
	The Integrating factor is, $IF = e^{\int P dx} = e^{\int \sec^2 x dx} = e^{\tan x}$	$\frac{1}{2}$
	On multiplying the equation (i) by $e^{\tan x}$ , we get	
	$\frac{d}{dx}(y.e^{\tan x}) = e^{\tan x} \tan x \left(\sec^2 x\right) \Longrightarrow d\left(y.e^{\tan x}\right) = e^{\tan x} \tan x \left(\sec^2 x\right) dx$	1
	On integrating we get, $y \cdot e^{\tan x} = \int t \cdot e^t dt + c_1$ ; where, $t = \tan x$ so that $dt = \sec^2 x  dx$	
	$= te^t - e^t + c = (\tan x)e^{\tan x} - e^{\tan x} + c$	
	$\therefore y = \tan x - 1 + c.(e^{-\tan x}), where \ 'c_1' \& 'c' \ are \ arbitrary \ constants \ of \ integration.$	1
30	The feasible region determined by the	
	constraints, $x + 2y \ge 100$ , $2x - y \le 0$ , $2x + y \le 200$ , $x, y \ge 0$ , is given below.	




[Class XII : Maths]



#### Section -D

[This section comprises of solution of long answer type questions (LA) of 5 marks each]



	Then, $(a,b)R(c,d) \Rightarrow ad = bc \Rightarrow bc = ad;$ (changing LHS and RHS)				
	$\Rightarrow cb = da;$ (As $a, b, c, d \in \mathbb{N}$ and multiplication is commutative on $\mathbb{N}$ )				
	$\Rightarrow$ $(c,d) R(a,b)$ ; according to the definition of the relation $R$ on $\mathbb{N} \times \mathbb{N}$				
	Thus $(a,b)R(c,d) \Rightarrow (c,d)R(a,b)$				
	So, <b>R</b> is symmetric relation on $\mathbb{N} \times \mathbb{N}$ .				
	Let $(a,b),(c,d),(e,f)$ be arbitrary elements of $\mathbb{N} \times \mathbb{N}$ such that				
	(a,b)R(c,d)and(c,d)R(e,f).				
	Then ${(a,b) R(c,d) \Rightarrow ad = bc} {(c,d) R(e,f) \Rightarrow cf = de} \Rightarrow (ad)(cf) = (bc)(de) \Rightarrow af = be$				
	$\Rightarrow (a,b)R(e,f);  (according to the definition of the relation R on \mathbb{N} \times \mathbb{N})$				
	Thus $(a,b)R(c,d)$ and $(c,d)R(e,f) \Rightarrow (a,b)R(e,f)$				
	So, <b>R</b> is transitive relation on $\mathbb{N} \times \mathbb{N}$ .				
	As the relation $R$ is reflexive, symmetric and transitive so, it is equivalence relation on $\mathbb{N} \times \mathbb{N}$ .	1			
	$[(2,6)] = \{(x,y) \in \mathbb{N} \times \mathbb{N} : (x,y) R(2,6)\}$	2			
	$=\left\{ \left( x,y\right) \in \mathbb{N} \times \mathbb{N} : 3x = y \right\}$	$\frac{1}{2}$			
	$= \{(x,3x): x \in \mathbb{N}\} = \{(1,3), (2,6), (3,9), \dots \}$				
		1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0\\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$ Injectivity: let $x, y \in \mathbb{R}^+ \cup \{0\}$ such that $f(x) = f(y)$ , then	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$ Injectivity: let $x, y \in \mathbb{R}^+ \cup \{0\}$ such that $f(x) = f(y)$ , then $\Rightarrow \frac{x}{1+x} = \frac{y}{1+y} \Rightarrow x + xy = y + xy \Rightarrow x = y$	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$ Injectivity: let $x, y \in \mathbb{R}^+ \cup \{0\}$ such that $f(x) = f(y)$ , then $\Rightarrow \frac{x}{1+x} = \frac{y}{1+y} \Rightarrow x + xy = y + xy \Rightarrow x = y$ So, $f$ is injective function.	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$ Injectivity: let $x, y \in \mathbb{R}^+ \cup \{0\}$ such that $f(x) = f(y)$ , then $\Rightarrow \frac{x}{1+x} = \frac{y}{1+y} \Rightarrow x + xy = y + xy \Rightarrow x = y$ So, $f$ is injective function. Surjectivity: when $x \ge 0$ , we have $f(x) = \frac{x}{1+x} \ge 0$ and $f(x) = 1 - \frac{1}{1+x} < 1$ , as $x \ge 0$	1			
33 OR	We have, $f(x) = \begin{cases} \frac{x}{1+x}, & \text{if } x \ge 0 \\ \frac{x}{1-x}, & \text{if } x < 0 \end{cases}$ Now, we consider the following cases <b>Case 1:</b> when $x \ge 0$ , we have $f(x) = \frac{x}{1+x}$ Injectivity: let $x, y \in \mathbb{R}^+ \cup \{0\}$ such that $f(x) = f(y)$ , then $\Rightarrow \frac{x}{1+x} = \frac{y}{1+y} \Rightarrow x + xy = y + xy \Rightarrow x = y$ So, $f$ is injective function. Surjectivity: when $x \ge 0$ , we have $f(x) = \frac{x}{1+x} \ge 0$ and $f(x) = 1 - \frac{1}{1+x} < 1$ , as $x \ge 0$ Let $y \in [0,1)$ , thus for each $y \in [0,1)$ there exists $x = \frac{y}{1-y} \ge 0$ such that $f(x) = \frac{\frac{y}{1-y}}{1+\frac{y}{1-y}} = y$ .	1			

	So, $f$ is onto function on $[0,\infty)$ to $[0,1)$ .		
	Case 2: when $x < 0$ , we have $f(x) = \frac{x}{1-x}$		
	Injectivity: Let $x, y \in \mathbb{R}^-$ i.e., $x, y < 0$ , such that $f(x) = f(y)$ , then		
	$\Rightarrow \frac{x}{1-x} = \frac{y}{1-y} \Rightarrow x - xy = y - xy \Rightarrow x = y$		
	So, $f$ is injective function.	2	
	Surjectivity: $x < 0$ , we have $f(x) = \frac{x}{1-x} < 0$ also, $f(x) = \frac{x}{1-x} = -1 + \frac{1}{1-x} > -1$	1	
	-1 < f(x) < 0.		
	Let $y \in (-1,0)$ be an arbitrary real number and there exists $x = \frac{y}{1+y} < 0$ such that,		
	$f(x) = f\left(\frac{y}{1+y}\right) = \frac{\frac{y}{1+y}}{1-\frac{y}{1+y}} = y.$		
	So, for $y \in (-1,0)$ , there exists $x = \frac{y}{1+y} < 0$ such that $f(x) = y$ .	1	
	Hence, $f$ is onto function on $(-\infty, 0)$ to $(-1, 0)$ .		
	Case 3:		
	(Injectivity): Let $x > 0$ & $y < 0$ such that $f(x) = f(y) \Rightarrow \frac{x}{1+x} = \frac{y}{1-y}$		
	$\Rightarrow x - xy = y + xy \Rightarrow x - y = 2xy$ , here <i>LHS</i> > 0 but <i>RHS</i> < 0, which is inadmissible.		
	Hence, $f(x) \neq f(y)$ when $x \neq y$ .	1	
	Hence $f$ is one-one and onto function.		
34	The given system of equations can be written in the form $AX = B$ ,		
	Where, $A = \begin{bmatrix} 2 & 3 & 10 \\ 4 & -6 & 5 \\ 6 & 9 & -20 \end{bmatrix}, X = \begin{bmatrix} 1/x \\ 1/y \\ 1/z \end{bmatrix} \text{ and } B = \begin{bmatrix} 4 \\ 1 \\ 2 \end{bmatrix}$		
	Now, $ A  = \begin{vmatrix} 2 & 3 & 10 \\ 4 & -6 & 5 \\ 6 & 9 & -20 \end{vmatrix} = 2(120 - 45) - 3(-80 - 30) + 10(36 + 36)$	$\frac{1}{2}$	
	$= 2(75) - 3(-110) + 10(72) = 150 + 330 + 720 = 1200 \neq 0  \therefore A^{-1} \text{ exists.}$	$\frac{1}{2}$	
	$\therefore adj A = \begin{bmatrix} 75 & 110 & 72 \\ 150 & -100 & 0 \\ 75 & 30 & -24 \end{bmatrix}^{T} = \begin{bmatrix} 75 & 150 & 75 \\ 110 & -100 & 30 \\ 72 & 0 & -24 \end{bmatrix}$	$1\frac{1}{2}$	





On solving the above equations , we get  $\lambda = \frac{2}{3}$  and  $\mu = 0$ 1 So, the position vector of the points, at which they should be so that the distance between them is the shortest, are  $\frac{2}{3}(\hat{i}-\hat{j}+\hat{k})$  and  $\hat{i}-\hat{j}$ . 1  $\overrightarrow{PQ} = \overrightarrow{OQ} - \overrightarrow{OP} = \frac{1}{3}\hat{i} - \frac{1}{3}\hat{j} - \frac{2}{3}\hat{k} \text{ and } \left|\overrightarrow{PQ}\right| = \sqrt{\left(\frac{1}{3}\right)^2 + \left(-\frac{1}{3}\right)^2 + \left(-\frac{2}{3}\right)^2} = \sqrt{\frac{2}{3}}$ 1 The shortest distance =  $\sqrt{\frac{2}{3}}$  units.  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{1}$   $\frac{p(\lambda, -\lambda, \lambda)}{p(\lambda, -\lambda, \lambda)}$ Method 2:  $Q(1,-1-2\mu,\mu)$  $\frac{x-1}{0} = \frac{y+1}{-2} = \frac{z}{1}$ The equation of two given straight lines in the Cartesian form are  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{1}$ .....(*i*) and  $\frac{x-1}{0} = \frac{y+1}{-2} = \frac{z}{1}$ .....(*ii*) The lines are not parallel as direction ratios are not proportional. Let P be a point on straight line 1 2 (i) and Q be a point on straight line (ii) such that line PQ is perpendicular to both of the lines. Let the coordinates of **P** be  $(\lambda, -\lambda, \lambda)$  and that of **Q** be  $(1, -2\mu - 1, \mu)$ ; where ' $\lambda$ ' and ' $\mu$ ' are 1 scalars. 2 Then the direction ratios of the line **PQ** are  $(\lambda - 1, -\lambda + 2\mu + 1, \lambda - \mu)$ Since PQ is perpendicular to straight line (i), we have, 1 2  $(\lambda - 1).1 + (-\lambda + 2\mu + 1).(-1) + (\lambda - \mu).1 = 0$  $\Rightarrow 3\lambda - 3\mu = 2.....(iii)$ 1 Since, PQ is perpendicular to straight line (*ii*), we have 2  $0.(\lambda - 1) + (-\lambda + 2\mu + 1).(-2) + (\lambda - \mu).1 = 0 \implies 3\lambda - 5\mu = 2.....(i\nu)$ 1 Solving (*iii*) and (*iv*), we get  $\mu = 0, \lambda = \frac{2}{3}$ 1 Therfore, the Coordinates of P are  $\left(\frac{2}{3}, -\frac{2}{3}, \frac{2}{3}\right)$  and that of Q are (1, -1, 0)1

So, the required shortest distance is 
$$\sqrt{\left(1-\frac{2}{3}\right)^2 + \left(-1+\frac{2}{3}\right)^2 + \left(0-\frac{2}{3}\right)^2} = \sqrt{\frac{2}{3}}$$
 units.

### Section -E

[This section comprises solution of 3 case- study/passage based questions of 4 marks each with two sub parts. Solution of the first two case study questions have three sub parts (i),(ii),(iii) of marks 1,1,2 respectively. Solution of the third case study question has two sub parts of 2 marks each.)

36 Let 
$$E_1, E_2, E_3$$
 be the events that Jayant, Sonia and Oliver processed the form, which are clearly  
pairwise mutually exclusive and exhaustive set of events.  
Then  $P(E_1) = \frac{50}{100} = \frac{5}{10}$ ,  $P(E_2) = \frac{20}{100} = \frac{1}{5}$  and  $P(E_3) = \frac{30}{100} = \frac{3}{10}$ .  
Also, let  $E$  be the event of committing an error.  
We have,  $P(E | E_1) = 0.06$ ,  $P(E | E_2) = 0.04$ ,  $P(E | E_3) = 0.03$ .  
(i) The probability that Sonia processed the form and committed an error is given by  
 $P(E \cap E_2) = P(E_2) \cdot P(E | E_2) = \frac{1}{5} \times 0.04 = 0.008$ .  
(ii) The total probability of committing an error in processing the form is given by  
 $P(E) = P(E_1) \cdot P(E | E_1) + P(E_2) \cdot P(E | E_2) + P(E_3) \cdot P(E | E_3)$   
 $P(E) = \frac{50}{100} \times 0.06 + \frac{20}{100} \times 0.04 + \frac{30}{100} \times 0.03 = 0.047$ .  
(iii) The probability that the form is processed by Jayant given that form has an error is given by  
 $P(E_1 | E) = \frac{P(E | E_1) \cdot P(E_1 | E_2) \cdot P(E_1 | E_2) \cdot P(E_2) + P(E | E_3) \cdot P(E_3)$   
 $= \frac{0.066 \times \frac{50}{100}}{0.06 \times \frac{20}{100} + 0.03 \times \frac{30}{100}} = \frac{30}{47}$   
Therefore, the required probability that the form is not processed by Jayant given that form has an  
error  $= P(\overline{E_1} | E) = 1 - P(E_1 | E) + P(E_2 | E) + P(E_3 | E) = 1$   
(iii) OR  $\sum_{i=1}^{3} P(E_i | E) = P(E_1 | E) + P(E_2 | E) + P(E_3 | E) = 1$   
Since, sum of the posterior probabilities is 1.

$$(We have, \sum_{i=1}^{3} P(E_i|E) = P(E_i|E) + P(E_i|E) + P(E_i|E)$$

$$= \frac{P(E \cap E_i) + P(E \cap E_i) + P(E \cap E_i)}{P(E)}$$

$$= \frac{P((E \cap E_i) \cup (E \cap E_i) \cup (E \cap E_i))}{P(E)} as E_i \& E_j; i \neq j, are mutually exclusive events$$

$$= \frac{P(E \cap (E_i \cup E_i \cup E_i))}{P(E)} = \frac{P(E \cap S)}{P(E)} = \frac{P(E)}{P(E)} = 1; 'S' being the sample space.)$$
37 We have,
$$[\overline{F_i}] = \sqrt{6^2 + \theta^2} = 6kN, |\overline{F_i}] = \sqrt{(-4)^2 + 4^2} = \sqrt{32} = 4\sqrt{2}kN, |\overline{F_i}] = \sqrt{(-3)^2 + (-3)^2} = \sqrt{18} = 3\sqrt{2}kN.$$
(i) Magnitude of force of Team  $A = 6kN.$ 
(ii) Since  $\overline{a} + \overline{c} = 3(\widehat{i} - \widehat{j})and \overline{b} = -4(\widehat{i} - \widehat{j})$ 
So,  $\overline{b}$  and  $\overline{a} + \overline{c}$  are unlike vectors having same initial point
and  $|\overline{b}| = 4\sqrt{2} \& |\overline{a} + \overline{c}| = 3\sqrt{2}$ 
Thus  $|\overline{F_i}| > |\overline{F_i} + \overline{F_i}| also \overline{F_i}$  and  $\overline{F_i} + \overline{F_i}$  are unlike
Hence B will win the game
(iii)  $\overline{F} = \overline{F_i} + \overline{F_2} + \overline{F_3} = 6\widehat{i} + 0\widehat{j} - 4\widehat{i} + 4\widehat{j} - 3\widehat{i} - 3\widehat{j} = -\widehat{i} + \widehat{j}$ 
 $\therefore |\overline{F}| = \sqrt{(-1)^2 + (1)^2} = \sqrt{2}kN.$ 
I
N
$$\overline{F} = -\widehat{i} + \widehat{j}$$
 $\therefore \theta = \pi - \tan^{-1}(\frac{1}{1}) = \pi - \frac{\pi}{4} = \frac{3\pi}{4};$  where '\theta' is the angle made by the resultant force with the
+ve direction of the x-axis.
38
$$y = 4x - \frac{1}{2}x^2$$
(i) The rate of growth of the plant with respect to the number of days exposed to sunlight
is given by  $\frac{dy}{dx} = 4 - x.$ 
(ii) Let rate of growth be represented by the function  $g(x) = \frac{dy}{dx}$ .

Now, 
$$g'(x) = \frac{d}{dx} \left( \frac{dy}{dx} \right) = -1 < 0$$
  
 $\Rightarrow g(x)$  decreases.  
So the rate of growth of the plant decreases for the first three days.  
Height of the plant after 2 days is  $y = 4 \times 2 - \frac{1}{2}(2)^2 = 6 cm$ .

# PRACTICE PAPER – 2 <u>CLASS XII</u> MATHEMATICS (CODE: 041)

## Time Allowed: 3 HOURS

Maximum Marks: 80

### **General Instructions:**

- 1. This question paper contains **FIVE sections A, B, C, D & E**. Each part is compulsory. However, there are internal choices in some questions.
- 2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
- 3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
- 4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
- 5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
- 6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

SECTION – A (Multiple Choice Questions) Each question carries 1 mark Each MCQ has four options with only one correct option, choose the correct option.					
1.	A function $f: R \to R$ defined as $f(x) = x^2 - 4x + 5$ is (a) injective but not surjective (b) surjective but not injective (c) both injective and surjective (d) neither injective nor surjective	1			
	SOLUTION : $f(x) = (x-2)^2 + 1$ As, $f(1) = f(3) \Rightarrow 1 \neq 3$ (Not injective) $f(x) \ge 1 \Rightarrow$ codomain $\neq$ Range (Not surjective) $OPTION(d)$				
2.	If $A = \begin{pmatrix} a & c & -1 \\ b & 0 & 5 \\ 1 & -5 & 0 \end{pmatrix}$ is a skew-symmetric matrix, then the value of $2a - (b + c)$ is	1			
	$(a) 0  (b) 1  (c) -10  (d) 10$ $SOLUTION : As, A \text{ is a skew-symmetric matrix} \Rightarrow a = 0, b = -c$ $Thus, 2a - (b + c) = 0 - 0 = 0  OPTION (a)$				

3.
 If A is a square matrix of order 3 such that the value of 
$$|adjA|=8$$
, then the value of  $|A^T|$  is

 (a)  $\sqrt{2}$ 
 (b)  $-\sqrt{2}$ 
 (c)8
 (d)  $2\sqrt{2}$ 
 (e)  $-\sqrt{2}$ 
 (c)8
 (d)  $2\sqrt{2}$ 
 (e)  $2\sqrt{2}$ 
 (f)  $2\sqrt{2}$ 
 (g)  $2\sqrt{2}$ 
2\sqrt{2}
 (g)  $2\sqrt{2}$ 
 (g

7.	If $xe^y = 1$ , then the value of $\frac{dy}{dx}$ at $x = 1$ is				1
	( <i>a</i> ) -1	(b)1	(c)-e	$(d)\frac{-1}{e}$	
	SOLUTION: $xe^{y} = 1 \Rightarrow \frac{dy}{dx} = \frac{-1}{x} \Rightarrow \frac{dy}{dx}\Big _{x=1} = -1$ OPTION (a)				
8.	Derivative of $e^{\sin^2 x}$ with respect to $\cos x$ is				1
	(a) $\sin x \cdot e^{\sin^2 x}$		$(b)\cos x$	$e^{\sin^2 x}$	
	$(c) - 2\cos x \cdot e^{\sin^2}$	x	$(d)-2 \operatorname{si}$	$n^2 x \cos x \cdot e^{\sin^2 x}$	
	SOLUTION : Le	$et y = e^{1 - \cos^2 x} \Longrightarrow \frac{d}{d(e^{1 - \cos^2 x})}$	$\frac{dy}{\cos x} = e^{1 - \cos^2 x} (-1)$	$2\cos x$ )	
	$\frac{dy}{d(\cos x)} = -2\cos x$	$os x.e^{\sin^2 x}$ OPTIC	DN (c)		
9.	The function $f($	$x = \frac{x}{2} + \frac{2}{x}$ has a lo	ocal minima at $x$	equal to:	1
	( <i>a</i> ) 2	(b)1	(c)0 (	d)-2	
	SOLUTION : A	$ls, f'(x) = \frac{1}{2} - \frac{2}{x^2} =$	$\Rightarrow x = \pm 2$		
	$so, f''(x) = \frac{4}{x^3} >$	> 0 when $x = 2$	OPTION(a)		
10.	Given a curve y	$=7x - x^3$ and x inc	creases at the rate	e of 2 units / sec.	1
	The rate at which the slope of curve changing, when $x = 5$ is				
	(a) - 60 units/se	c	( <i>b</i> ) 60 ι	inits/sec	
	$\frac{(c)-70 \text{ units/se}}{(c)-70 \text{ units/se}}$	c	$(d) - 1^2$	40 units/sec	
	SOLUTION : Gi	ven, $\frac{dx}{dt} = 2$ units /	sec		
	As, $y = 7x - x^3 \Rightarrow m = \frac{dy}{dx} = 7 - 3x^2$				
	Now, $\frac{dm}{dt} = -6x$	$c\frac{dx}{dt} = -30 \times 2 = -60$	) units/sec OPT	ION(a)	

11. 
$$\int \frac{1}{x(\log x)^2} dx \text{ is equal to:} \qquad 1$$
(a)  $2\log(\log x) + c$ 
(b)  $\frac{-1}{\log x} + c$ 
(c)  $\frac{(\log x)^3}{3} + c$ 
(d)  $\frac{3}{(\log x)^3} + c$ 
SOLUTION :  $Let, I = \int \frac{1}{x} (\log x)^2 dx = \frac{(\log x)^{-2+1}}{-2+1} + c = \frac{-1}{\log x} + c$ 
(OPTION (b)
  
12. The value of  $\int_{-1}^{1} x |x| dx$  is:
$$\frac{(a)}{6} \frac{1}{6} (b) \frac{1}{3} (c) \frac{-1}{6} (d) 0$$
SOLUTION : As, For  $f(x) = x |x| \Rightarrow f(-x) = -f(x)$ 
Thus,  $\int_{-1}^{1} x |x| dx = 0$  (Option (d)
  
13. Area of the region bounded by the curve  $y^2 = 4x$  and the x-axis between  $x = 0$  &  $x = 1$  is (in units)
$$\frac{(a)}{2} \frac{2}{3} (b) \frac{8}{3} (c) 3 (d) \frac{4}{3}$$
SOLUTION :  $A = 2\int_{0}^{1} 2\sqrt{x} dx = 4\left(\frac{x^{\frac{3}{2}}}{\frac{3}{2}}\right)_{0}^{1} = \frac{8}{3}$  (OPTION (b)

14. The order of the differential equation 
$$\frac{d^4y}{dx^4} - \sin\left(\frac{d^2y}{dx^2}\right) = 5$$
 is:  
(a) 4 (b) 3 (c) 2 (d) not defined  
SOLUTION : order of the differential equation = 4  
[OPTION (a)]  
15. The position vectors of points P & Q are  $\vec{p}$  and  $\vec{q}$  respectively. The  
point R divides line segment PQ in the ratio 3:1 & S is the mid-point  
of the line segment PR. The position vector of S is  
(a)  $\frac{\vec{p}+3\vec{q}}{4}$  (b)  $\frac{\vec{p}+3\vec{q}}{8}$  (c)  $\frac{5\vec{p}+3\vec{q}}{4}$  (d)  $\frac{5\vec{p}+3\vec{q}}{8}$   
SOLUTION :  $\overset{P}{<} - \cdots \overset{S}{-\cdots} \overset{R}{-\cdots} \overset{Q}{-}$   
As, S divides line segment PQ in the ratio  $\frac{3}{2}:(\frac{3}{2}+1)=3:5$   
so,  $S = \frac{3Q+5P}{3+5} = \frac{5\vec{p}+3\vec{q}}{8}$  [OPTION (d)]  
16. The angle which the line  $\frac{x}{1} = \frac{y}{-1} = \frac{z}{0}$  makes with the positive  
direction of Y-axis is  
(a)  $\frac{5\pi}{6}$  (b)  $\frac{3\pi}{4}$  (c)  $\frac{5\pi}{4}$  (d)  $\frac{7\pi}{4}$   
SOLUTION : As,  $\cos \beta = \frac{-1}{\sqrt{2}} = \cos \frac{3\pi}{4} \Rightarrow \beta = \frac{3\pi}{4}$   
[OPTION (b)]

17.	The cartesian equation of the line passing through the point		
	$(1, -3, 2)$ and parallel to the line $\vec{r} = (2 + \lambda)\hat{i} + \lambda\hat{j} + (2\lambda - 1)\hat{k}$		
	is		
	(a) $\frac{x-1}{2} = \frac{y+3}{0} = \frac{z-2}{-1}$ (b) $\frac{x+1}{1} = \frac{y-3}{1} = \frac{z+2}{2}$		
	$\underbrace{(c) \ \frac{x+1}{2} = \frac{y-3}{0} = \frac{z+2}{-1}}_{(d) \ \frac{x+1}{1} = \frac{y+3}{1} = \frac{z-2}{2}}$		
	SOLUTION: $\vec{r} = (2+\lambda)\hat{i} + \lambda\hat{j} + (2\lambda-1)\hat{k} = (2\hat{i} - \hat{k}) + \lambda(\hat{i} + \hat{j} + 2\hat{k})$		
	D.r. of the Parallel line = $<1,1,2>$ OPTION (d)		
18.	If A & B are events such that $P(A   B) = P(B   A) \neq 0$ , then		
	(a) $A \subset B$ but $A \neq B$ (b) $A = B$	1	
	(c) $A \cap B = \phi$ (d) $P(A) = P(B)$		
	SOLUTION: As, $P(\frac{A}{B}) = P(\frac{B}{A}) \Rightarrow P(A) = P(B)$		
	OPTION (d)		
	ASSERTION-REASON BASED QUESTIONS (Q.19 & Q.20)		
	In the following questions, a statement of assertion (A) is followed by a statement of Reason (R). Choose the correct answer out of the following choices.		
	(a) Both A and R are true and R is the correct explanation of A.		
	(b) Both A and R are true but R is not the correct explanation of A. (c) A is true but R is false.		
	(d) A is false but R is true.		
		1	

19.ASSERTION(A): Domain of 
$$y = \cos^{-1}x$$
 is  $[-1,1]$ .  
REASONING(R): The range of the principal value branch of  
 $y = \cos^{-1}x$  is  $[0,\pi] - \{\frac{\pi}{2}\}$ .  
SOLUTION: As, **A is true but R is false**.  
 $OPTION(c)$ 20.ASSERTION(A): The vectors  $\vec{a} = 6\hat{i} + 2\hat{j} - 8\hat{k}$ ,  $\vec{b} = 10\hat{i} - 2\hat{j} - 6\hat{k}$   
and  $\vec{c} = 4\hat{i} - 4\hat{j} + 2\hat{k}$  represents the sides of a right  
angled triangle.  
REASONING(R): Three non-zero vectors of which none of the two are  
collinear forms a triangle if their resultant is zero  
vector or sum of any two vectors is equal to the third.  
SOLUTION :  $As, \vec{a} + \vec{b} + \vec{c} = \vec{0} \Rightarrow$  Forms a triangle but for right angled  
triangle we have to further analyse.  
Since,  $|\vec{a}| = \sqrt{104}$ ,  $|\vec{b}| = \sqrt{140}$ ,  $|\vec{c}| = \sqrt{36} \Rightarrow |\vec{b}|^2 = |\vec{a}|^2 + |\vec{c}|^2$   
 $OR$  We can observe  $\vec{a}.\vec{c} = 0$   
So, The vectors  $\vec{a} = 6\hat{i} + 2\hat{j} - 8\hat{k}$ ,  $\vec{b} = 10\hat{i} - 2\hat{j} - 6\hat{k}$  and  $\vec{c} = 4\hat{i} - 4\hat{j} + 2\hat{k}$   
represents the sides of a right angled triangle.  
Both A and R are true but R is not the correct explanation of A.  
 $OPTION(b)$ 21.Find the value of k if  $\sin^{-1}\left[k \tan\left(2\cos^{-1}\frac{\sqrt{3}}{2}\right]\right] = \frac{\pi}{3}$ .  
 $SOLUTION : \left[k \tan\left(2\cos^{-1}\frac{\sqrt{3}}{2}\right)\right] = \sin\frac{\pi}{3} = \frac{\sqrt{3}}{2} \Rightarrow \left[k \tan\left(\frac{\pi}{3}\right)\right] = \frac{\sqrt{3}}{2}$   
 $\Rightarrow k\sqrt{3} = \frac{\sqrt{3}}{2} \Rightarrow \left[k = \frac{1}{2} = 0.5\right]$ 

22. (a) Verify whether the function f defined by  

$$f(x) = \begin{cases} x. \sin\left(\frac{1}{x}\right), x \neq 0 \\ 0, x = 0 \end{cases}$$
is continuous at  $x = 0$  or not.  
SOLUTION : We know that  $f(x)$  is continuous at  $x = 0$  when  
LHL=RHL =  $f(0)$ . So,  
LHL(Left Hand Limit)  $x = a - h = 0 - h$   
 $\lim_{h \to 0} f(0 - h) = \lim_{h \to 0} (-h) \sin\left(\frac{1}{-h}\right) = \lim_{h \to 0} h. \sin\left(\frac{1}{h}\right)$   
 $= 0 \times (\text{Any finite value between - 1 to 1})$   
 $= 0$   
NOTE :  $\sin(-\theta) = -\sin\theta \& -1 \le \sin\theta \le 1$   
RHL(Right Hand Limit)  $x = a + h = 0 + h$   
 $\lim_{h \to 0} f(0 - h) = \lim_{h \to 0} h. \sin\left(\frac{1}{h}\right) = 0 \times (\text{Any finite value between - 1 to 1})$   
 $= 0$   
Now,  $f(x = 0) = 0$  (Given)  
Since, LHL = RHL =  $f(x = 0)$   
So,  $f(x)$  is Continuous at  $x = 0$ 

OR  
Check the differentiability of the function f defined by  

$$f(x) = |x-5|$$
 at the point  $x = 5$ .  
SOLUTION : First of all we have to redefine  $f(x)$ , so  
 $f(x) = |x-5| = \begin{cases} x-5, \text{ when } x > 5\\ 5-x, \text{ when } x \le 5 \end{cases}$   
(L.H.D.):  $Lf'(5) = \lim_{h \to 0} \frac{f(5-h)-f(5)}{-h} = \lim_{h \to 0} \frac{h-0}{-h} = -1$   
(R.H.D.):  $Rf'(5) = \lim_{h \to 0} \frac{f(5+h)-f(5)}{h} = \lim_{h \to 0} \frac{h-0}{h} = 1$   
Since,  $LHD \neq RHD$   
so function  $f(x)$  is not differentiable at  $x = 5$ .  
<sup>23.</sup> The area of the circle is increasing at a uniform rate of 2 cm<sup>2</sup>/sec.  
How fast is the circumference of the circle increasing when the  
radius  $r = 5cm$ ?  
SOLUTION :  $As, \frac{dA}{dt} = 2 cm^2/sec$ . so,  $A = \pi r^2$   
 $\Rightarrow \frac{dA}{dt} = 2\pi r \frac{dr}{dt} = 10\pi \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = \frac{1}{5\pi} cm/sec}$   
Now, Circumference  $C = 2\pi r$   
 $\Rightarrow \frac{dC}{dt} = 2\pi \frac{dr}{dt} = \frac{2}{5} cm/sec}$   
Alternative:  $A = \pi r^2 \Rightarrow \frac{dA}{dt} = r\left(2\pi \frac{dr}{dt}\right) = 5\frac{dC}{dt}$   
 $\Rightarrow \frac{dC}{dt} = \frac{2}{5} cm/sec}$ 

<sup>24.</sup> (a) Find: 
$$\int \cos^3 x \cdot e^{\log \sin x} dx$$
  
SOLUTION:  $I = \int \cos^3 x \cdot e^{\log \sin x} dx = \int \cos^3 x \cdot \sin x dx$   
 $I = -\int \cos^3 x d(\cos x) = \frac{-\cos^4 x}{4} + c$   
OR  
(b) Find:  $\int \frac{1}{5+4x-x^2} dx$   
SOLUTION:  $I = \int \frac{-1}{x^2-4x-5} dx = -\int \frac{1}{(x-2)^2-3^2} dx$   
 $I = \frac{-1}{6} \ln \left| \frac{x-5}{x+1} \right| + c OR I = \frac{1}{6} \ln \left| \frac{x+1}{x-5} \right| + c$   
<sup>25.</sup> Find the vector equation of the line passing through the point  
(2,3,-5) and making equal angles with the coordinates axes.  
and  
 $\vec{r} = (7\hat{i} - 6\hat{k}) + \mu(\hat{i} + 2\hat{j} + 2\hat{k})$   
SOLUTION: Here,  $\alpha = \beta = \gamma \Rightarrow 3\cos^2 \alpha = 1$   
d.r. of Line  $= \langle \frac{\pm 1}{\sqrt{3}}, \frac{\pm 1}{\sqrt{3}}, \frac{\pm 1}{\sqrt{3}} \rangle$   
Now, Equation of the line  
 $\vec{r} = (2\hat{i} + 3\hat{j} - 5\hat{k}) + \lambda \cdot (\frac{\pm 1}{\sqrt{3}}\hat{i} + \frac{\pm 1}{\sqrt{3}}\hat{j} + \frac{\pm 1}{\sqrt{3}}\hat{k})$   
 $OR[\vec{r} = (2\hat{i} + 3\hat{j} - 5\hat{k}) + \lambda(\hat{i} + \hat{j} + \hat{k})]$ 

SECTION C  
(This section comprises of short answer type questions (SA) of 3 marks each)  
26. (a) Find 
$$\frac{dy}{dx}$$
, if  $(\cos x)^y = (\cos y)^x$ .  
SOLUTION: As,  $(\cos x)^y = (\cos y)^x$   
 $\Rightarrow y \ln(\cos x) = x \ln(\cos y)$   
on differentiating both sides with respect to x, we get  
 $y(-\tan x) + \ln(\cos x)\frac{dy}{dx} = x(-\tan y)\frac{dy}{dx} + \ln(\cos y)$   
 $\therefore \left[\frac{dy}{dx} = \left(\frac{\ln(\cos y) + y \tan x}{\ln(\cos x) + x \tan y}\right)\right]$   
OR  
(b) If  $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$ , then prove that  $\frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$   
SOLUTION : so, Let  $x = \sin A, y = \sin B$   
Thus,  $\sqrt{1-\sin^2 A} + \sqrt{1-\sin^2 B} = a(\sin A - \sin B)$   
 $\Rightarrow \cos A + \cos B = a(\sin A - \sin B)$   
 $\Rightarrow 2\cos(\frac{A+B}{2}) \cos(\frac{A-B}{2}) = a(2\cos(\frac{A+B}{2})) \sin(\frac{A-B}{2}))$   
 $\Rightarrow \cot(\frac{A-B}{2}) = a \Rightarrow A - B = 2\cot^{-1} a$   
 $\Rightarrow \frac{\sin^{-1}x - \sin^{-1}y = 2\cot^{-1} a}{\sqrt{1-y^2} \frac{dy}{dx}} = 0 \Rightarrow \frac{1}{\sqrt{1-y^2}}\frac{dy}{dx} = \frac{1}{\sqrt{1-x^2}}$   
 $\therefore \frac{dy}{dx} = \frac{\sqrt{1-y^2}}{\sqrt{1-x^2}} = \sqrt{\frac{1-y^2}{1-x^2}}$ 

<sup>27.</sup> If 
$$x = a \sin^3 \theta$$
,  $y = b \cos^3 \theta$ , then find  $\frac{d^2 y}{dx^2}$  at  $\theta = \frac{\pi}{4}$ .  
SOLUTION :  $As, x = a \sin^3 \theta \Rightarrow \frac{dx}{d\theta} = 3a \sin^2 \theta \cdot \cos \theta$   
 $y = b \cos^3 \theta \Rightarrow \frac{dy}{d\theta} = -3b \cos^2 \theta \cdot \sin \theta$   
 $\frac{dy}{dx} = \frac{-b}{a} \cot \theta$   
Now, Differentiate again with respect to x both the sides,  
 $\frac{d^2 y}{dx^2} = \frac{b}{a} \csc^2 \theta \frac{d\theta}{dx} = \frac{b}{a} \csc^2 \theta \frac{1}{3a \sin^2 \theta \cdot \cos \theta}$   
 $\frac{d^2 y}{dx^2} = \frac{b}{3a^2} \csc^2 \theta \cdot \sec \theta$   
Now,  $\frac{d^2 y}{dx^2} = \frac{b}{3a^2} \csc^2 \theta \cdot \sec \theta$ 

28. (a) Evaluate : 
$$\int_{0}^{\pi} \frac{e^{\cos x}}{e^{\cos x} + e^{-\cos x}} dx$$
SOLUTION : Let,  $I = \int_{0}^{\pi} \frac{e^{\cos x}}{e^{\cos x} + e^{-\cos x}} dx$ ...(1)
On applying 
$$\int_{0}^{a} f(x) dx = \int_{0}^{a} f(a-x) dx$$
,
we get  $I = \int_{0}^{\pi} \frac{e^{-\cos x}}{e^{-\cos x} + e^{\cos x}} dx$ ...(2)
On adding Eq. (1) and (2), we get
$$2I = \int_{0}^{\pi} \frac{e^{-\cos x} + e^{\cos x}}{e^{-\cos x} + e^{-\cos x}} dx = \int_{0}^{\pi} 1 dx = (x)_{0}^{\pi} = \pi - 0 = \pi$$

$$\therefore \boxed{I = \int_{0}^{\pi} \frac{e^{\cos x} + e^{-\cos x}}{e^{\cos x} + e^{-\cos x}} dx = \frac{\pi}{2}}$$
OR
(b) Find:  $\int \frac{2x+1}{(x+1)^{2}(x-1)} dx$ 
Solution :  $I = \int \frac{x+1+x}{(x+1)^{2}(x-1)} dx = \int \frac{1}{(x+1)(x-1)} dx + \frac{1}{2} \int \frac{(x+1)+(x-1)}{(x+1)^{2}(x-1)} dx$ 

$$I = \frac{3}{2} \int \frac{1}{(x+1)(x-1)} dx + \frac{1}{2} \int \frac{1}{(x+1)^{2}} dx$$

$$I = \frac{3}{4} \int \left(\frac{1}{x-1} - \frac{1}{x+1}\right) dx + \frac{1}{2} \int \frac{1}{(x+1)^{2}} dx$$

$$\therefore \boxed{I = \frac{3}{4} \ln \left| \frac{x-1}{x+1} \right| - \frac{1}{2} \left( \frac{1}{x+1} \right) + c}$$
(Student may try using Partial Fraction also)

29. (a) Find the particular solution of the differential equation  

$$\frac{dy}{dx} - 2xy = 3x^2 \cdot e^{x^2}; y(0) = 5$$
Solution: On Comparing  $\frac{dy}{dx} - 2xy = 3x^2 \cdot e^{x^2}$  with  

$$\frac{dy}{dx} + P(x) \cdot y = Q(x), \text{ we get } P(x) = -2x \quad \& \quad Q(x) = 3x^2 \cdot e^{x^2}$$
Now, Integrating Factor  $= e^{\int P(x)dx} = e^{\int -2xdx} = e^{-x^2}$   
Thus, solution of given Differential equation as  
 $y \cdot I \cdot F \cdot = \int Q(x) \cdot I \cdot F \cdot dx$   
 $\Rightarrow y \cdot e^{-x^2} = \int e^{-x^2} \cdot 3x^2 \cdot e^{x^2} dx$   
 $\Rightarrow y \cdot e^{-x^2} = \int (3x^2) dx = x^3 + c$   
 $\therefore \boxed{y \cdot e^{-x^2} = x^3 + c}$   
Now, For  $y(0) = 5 \Rightarrow 5 = 0 + c \Rightarrow \boxed{c = 5}$   
 $\therefore \boxed{y \cdot e^{-x^2} = x^3 + 5}$   
Required Solution of the given Differential Equation

OR  
(b) Solve the following differential equation:  

$$x^2 dy + y(x+y)dx = 0$$
  
Solution: Here,  $\frac{dy}{dx} = \frac{-(yx+y^2)}{x^2} = -\frac{y}{x} - \left(\frac{y}{x}\right)^2 = f(\frac{y}{x})...(1)$   
so, its a homogeneous differential equation  
Let  $y = vx \Rightarrow \frac{dy}{dx} = v + x\frac{dv}{dx}$   
Now, Equation (1) becomes  $v + x\frac{dv}{dx} = -v - v^2$   
 $\Rightarrow x\frac{dv}{dx} = -(2v+v^2)$   
 $\Rightarrow \frac{dv}{(v+1)^2 - 1} = -\frac{1}{x}dx$   
 $\Rightarrow \int \frac{dv}{(v+1)^2 - 1} = \int -\frac{1}{x}dx$   
 $\Rightarrow \frac{1}{2}\ln\left|\frac{v}{v+2}\right| + \ln x = \ln c' \Rightarrow \frac{v}{v+2} = \frac{c}{x^2}$   
 $\therefore \boxed{x^2y = c(y+2x)}$   
Required Solution of the given Differential Equation

30. Find a vector of magnitude 4 units perpendicular to each of  
vectors 
$$2\hat{i} - \hat{j} + \hat{k}$$
 and  $\hat{i} + \hat{j} - \hat{k}$  and hence verify your answer.  
Solution : Say  $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ ,  $\vec{b} = \hat{i} + \hat{j} - \hat{k}$   
 $So, \vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = 0\hat{i} + 3\hat{j} + 3\hat{k}$   
Now, vector of magnitude 4 units perpendicular to  $\vec{a} \ll \vec{b}$  is  
 $4\left(\frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|}\right) = \pm 2\sqrt{2}(0\hat{i} + \hat{j} + \hat{k})$   
 $\vec{r} = (0\hat{i} + 2\sqrt{2}\hat{j} + 2\sqrt{2}\hat{k})$  or  $(0\hat{i} - 2\sqrt{2}\hat{j} - 2\sqrt{2}\hat{k})$   
Any one of the answer is acceptable.  
Verification:  $|\vec{r}| = \sqrt{0 + 8 + 8} = 4$   
 $\vec{r}.\vec{a} = (0\hat{i} + 2\sqrt{2}\hat{j} + 2\sqrt{2}\hat{k}).(\hat{i} + \hat{j} - \hat{k}) = 0 + 2\sqrt{2} - 2\sqrt{2} = 0$   
 $\vec{r}.\vec{b} = (0\hat{i} + 2\sqrt{2}\hat{j} + 2\sqrt{2}\hat{k}).(\hat{i} + \hat{j} - \hat{k}) = 0 + 2\sqrt{2} - 2\sqrt{2} = 0$   
Hence,  $\vec{r}$  vector has magnitude 4 units and perpendicular to  
each of vectors  $2\hat{i} - \hat{j} + \hat{k}$  and  $\hat{i} + \hat{j} - \hat{k}$ .



32. (a) If  $A = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 0 & -3 \\ 1 & 2 & 0 \end{pmatrix}$ , then find  $A^{-1}$  and hence solve the following 5 system of equations: x + 2y - 3z = 1, 2x - 3z = 2, x + 2y = 3SOLUTION :  $As, A = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 0 & -3 \\ 1 & 2 & 0 \end{pmatrix}$  $adjA = \begin{pmatrix} 6 & -6 & -6 \\ -3 & 3 & -3 \\ 4 & 0 & -4 \end{pmatrix}, |A| = 6 - 6 - 12 = -12$ Thus,  $A^{-1} = \frac{-1}{12} \begin{pmatrix} 6 & -6 & -6 \\ -3 & 3 & -3 \\ 4 & 0 & -4 \end{pmatrix}$ so, Given equation can be written into a matrix equation as  $\begin{pmatrix} 1 & 2 & -3 \\ 2 & 0 & -3 \\ 1 & 2 & 0 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} \Longrightarrow X = \mathbf{A}^{-1}.B$  $A \quad X = B$  $\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{-1}{12} \begin{pmatrix} 6 & -6 & -6 \\ -3 & 3 & -3 \\ 4 & 0 & -4 \end{pmatrix} \begin{pmatrix} 1 \\ 2 \\ 3 \end{pmatrix} = \frac{-1}{12} \begin{pmatrix} -24 \\ -6 \\ -8 \end{pmatrix} = \begin{pmatrix} 2 \\ 1/2 \\ 2/3 \end{pmatrix}$  $\therefore x = 2, y = \frac{1}{2}, z = \frac{2}{3}$ 

$$OR$$
(b) Find the product of the matrices  $\begin{pmatrix} 1 & 2 & -3 \\ 2 & 3 & 2 \\ 3 & -3 & -4 \end{pmatrix} \begin{pmatrix} -6 & 17 & 13 \\ 14 & 5 & -8 \\ -15 & 9 & -1 \end{pmatrix}$ 
and hence solve the system of linear equations:  
 $x + 2y - 3z = -4, 2x + 3y + 2z = 2, 3x - 3y - 4z = 11$ 

$$SOLUTION : Let, AB = \begin{pmatrix} 1 & 2 & -3 \\ 2 & 3 & 2 \\ 3 & -3 & -4 \end{pmatrix} \begin{pmatrix} -6 & 17 & 13 \\ 14 & 5 & -8 \\ -15 & 9 & -1 \end{pmatrix}$$

$$So, AB = \begin{pmatrix} 67 & 0 & 0 \\ 0 & 67 & 0 \\ 0 & 0 & 67 \end{pmatrix} = 67I$$
Thus,  $A^{-1} = \frac{1}{67}B = \frac{1}{67} \begin{pmatrix} -6 & 17 & 13 \\ 14 & 5 & -8 \\ -15 & 9 & -1 \end{pmatrix}$ 

$$so, Given equation can be written into a matrix equation as$$

$$\begin{pmatrix} 1 & 2 & -3 \\ 2 & 3 & 2 \\ 3 & -3 & -4 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} -4 \\ 2 \\ 11 \end{pmatrix} \Rightarrow X = A^{-1}.C$$

$$A \qquad X = C$$

$$\begin{pmatrix} x \\ y \\ z \end{pmatrix} = \frac{1}{67} \begin{pmatrix} -6 & 17 & 13 \\ 14 & 5 & -8 \\ -15 & 9 & -1 \end{pmatrix} \begin{pmatrix} -4 \\ 2 \\ 11 \end{pmatrix} = \frac{1}{67} \begin{pmatrix} 201 \\ -134 \\ 67 \end{pmatrix} = \begin{pmatrix} 3 \\ -2 \\ 1 \end{pmatrix}$$

$$\therefore \boxed{x = 3, y = -2, z = 1}$$

<sup>33.</sup> Find the area of the region bounded by the curve 
$$4x^2 + y^2 = 36$$
  
using integration.  
SOLUTION:  $4x^2 + y^2 = 36$   
 $\Rightarrow \frac{x^2}{3^2} + \frac{y^2}{6^2} = 1$ . Thus,  $y = 2\sqrt{3^2 - x^2}$   
This is the equation of an Ellipse.  
Area of ellipse = 4(Area of AoB)  
 $A = 4\int_{0}^{3} 2\sqrt{3^2 - x^2} dx = 8\int_{0}^{3} \sqrt{3^2 - x^2} dx$   
 $A = 8\left[\frac{x\sqrt{3^2 - x^2}}{2} + \frac{9}{2}\sin^{-1}\left(\frac{x}{3}\right)\right]_{0}^{3} = 8\left[\frac{9}{2}\sin^{-1}(1)\right]$   
 $A = 36\frac{\pi}{2} = 18\pi$  unit<sup>2</sup>  
 $\therefore$  Area bounded by the curve,  $A = 18\pi$  unit<sup>2</sup>



OR (b) Find the shortest distance between the lines  $L_1 \& L_2$ Given below:  $L_1$ : The line passing through (2, -1, 1) and parallel to  $\frac{x}{1} = \frac{y}{1} = \frac{z}{3}$  $L_{2}: \vec{r} = \hat{i} + (2\mu + 1)\hat{j} - (\mu + 2)\hat{k}$ SOLUTION: Equations of Line  $L_1 \& L_2$  are  $L_1: \vec{r} = (2\hat{i} - \hat{j} + \hat{k}) + \lambda(\hat{i} + \hat{j} + 3\hat{k})$ L<sub>2</sub>:  $\vec{r} = (\hat{i} + \hat{j} - 2\hat{k}) + \mu(\hat{0i} + 2\hat{j} - \hat{k})$ Now,  $\vec{a_2} - \vec{a_1} = (\hat{i} + \hat{j} - 2\hat{k}) - (2\hat{i} - \hat{j} + \hat{k}) = (-\hat{i} + 2\hat{j} - 3\hat{k})$  $\begin{vmatrix} \vec{b}_1 \times \vec{b}_2 \\ \vec{b}_1 \times \vec{b}_2 \end{vmatrix} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 1 & 1 & 3 \\ 0 & 2 & -1 \end{vmatrix} = -7\hat{i} + \hat{j} + 2\hat{k}$  $|\vec{b_1} \times \vec{b_2}| = \sqrt{49 + 1 + 4} = \sqrt{54} = 3\sqrt{6}$ Thus, Shortest Distance between the lines is S.D. =  $\left| \frac{(\vec{a_2} - \vec{a_1}) \cdot (\vec{b_1} \times \vec{b_2})}{|\vec{b_1} \times \vec{b_1}|} \right| = \left| \frac{7 + 2 - 6}{3\sqrt{6}} \right| = \frac{1}{\sqrt{6}} = \frac{\sqrt{6}}{6}$  units



This section comprises of 3 case-study/passage-based questions of 4 marks each with two sub-parts.

First two case study questions have three sub parts (A), (B) & (C) of marks 1, 1, 2 respectively. The third case study question has two sub-parts of 2 marks each.






38.	A departmental store sends bills to charge its customers once a month.	4		
	Past Experience show that 70% of its customers pay their first month			
	bill in time. The store also found that the customer who pays the bill			
	in time has the probability of 0.8 of paying in the next month and the			
	customer who doesn't pay in time has the probability of 0.4 of paying			
	in time the next month.			
	Based on the above information, answer the following questions:			
	(i) Let $E_1$ and $E_2$ respectively denote the event of customer paying or			
	not paying the first month bill in time. Find $P(E_1)$ and $P(E_2)$ .			
	(ii) Let A denotes the event of customer paying second month's bill			
	in time, then find $P(A E_1)$ and $P(A E_2)$ .			
	(iii) Find the probability of customer paying second month's bill in time.			
	OR			
	Find the probability of customer paying first month's bill in time if			
	it is found that customer has paid the second month's bill in time.			
	Solution: (i) $P(E_1) = \frac{70}{100} = \frac{7}{10}$ and $P(E_2) = 1 - \frac{7}{10} = \frac{3}{10}$			
	(ii) $P(A E_1) = 0.8$ and $P(A E_2) = 0.4$			
	(iii) Probability of customer paying second month's bill in time			
	is $P(A) = P(A E_1) \times P(E_1) + P(A E_2) \times P(E_2)$			
	$P(A) = \frac{7 \times 0.8}{10} + \frac{3 \times 0.4}{10} = \frac{5.6 + 1.2}{10} = 0.68$			
	OR			
	(iii) Probability of customer paying first month's bill in time if			
	it is found that customer has paid the second month's bill in time			
	is $P(E_1 A) = \frac{P(A E_1) \times P(E_1)}{P(A E_1) \times P(E_1)}$			
	$P(A E_1) \times P(E_1) + P(A E_2) \times P(E_2)$			
	$\Rightarrow P(E_1 A) = \frac{0.56}{0.68} = \frac{14}{17}$			

# PRACTICE PAPER – 3 <u>CLASS XII</u> MATHEMATICS (CODE: 041)

### Time Allowed: 3 HOURS

#### Maximum Marks: 80

## **General Instructions:**

- 1. This question paper contains **FIVE sections A, B, C, D & E**. Each part is compulsory. However, there are internal choices in some questions.
- 2. Section A has 18 MCQ's and 02 Assertion-Reason based questions of 1 mark each.
- 3. Section B has 5 Very Short Answer (VSA)-type questions of 2 marks each.
- 4. Section C has 6 Short Answer (SA)-type questions of 3 marks each.
- 5. Section D has 4 Long Answer (LA)-type questions of 5 marks each.
- 6. Section E has 3 source based/case based/passage based/integrated units of assessment (4 marks each) with sub parts.

		SEC	TION – A		
		(Multiple Cl	noice Questions)		
Ea	ch MCQ has four	options with only or	n carries 1 mark e correct option, c	hoose the correct opti	on.
1.	A set of value constraints and (a)Unbounded (b) Feasible s	s of decision varial d non–negativity co l solution	bles that satisfies to bles that satisfies to bless that satisfies to bless bless that bless that bless that bless bles	he linear P.P. is called its :	1
	(c) Optimum s (d) None of t	olution hese			
2.	The value of the expression $\cos ec^{-1}(2) + \cos^{-1}(\frac{1}{2}) + \tan^{-1}(-1)$ is			1	
·	(a) $\frac{\pi}{4}$	(b) $\frac{\pi}{3}$	(c) $\frac{\pi}{6}$	(d) $\frac{\pi}{2}$	
3.	If $A = \begin{pmatrix} 1 & 2 \\ 2 & 1 \end{pmatrix}$	$B = \begin{pmatrix} 1\\4\\0 \end{pmatrix} \& C = \begin{pmatrix} 0\\3 \end{pmatrix}$	$\begin{pmatrix} 0 & 1 \\ 1 & 2 \end{pmatrix}$ , then order of	of Matrix P is,	1
	where $P = AC$	CB			
	(a) $2 \times 1$	(b) $2 \times 3$	(c) $2 \times 2$	(d) 3×2	

4.	If A is square matrix of order $3 \times 3$ such that $ adj A  = 16$ , such that	
	$( 2A )^2=2^p$ , then the $p=$	1
	(a) 4  (b) 5  (c) 8  (d) 10	
5.	For an L.P.P. the objective function is $Z = 400x + 300y$ , and the feasible	
	region determined by a set of constraints (linear inequations) is shown in the	1
	graph.Coordinates at which the objective function is maximum is	
	(a) (20, 0) (b) (40, 0) (c) (40, 160) (d) (20, 180)	
6.	If A and B are two independent events with $P(A) = \frac{3}{5} \& P(B) = \frac{4}{9}$ , then $9P(\overline{A} \cap \overline{B}) =$	1
	(a) 1 (b) 2 (c) 3 (d) 4	
7.	If $\vec{a}$ is a unit vector and $(\vec{x} + \vec{a})$ $(\vec{x} - \vec{a}) = 15$ then $ \vec{x}  = 15$	
	$(x + a) \cdot (x + a) = 15$ , since $ x  = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) = 15$ , $(x + a) \cdot (x + a) \cdot (x$	1
8	$ \begin{array}{c} (a) & 1 \\ (b) & 2 \\ (c) & 5 \\ (a) & 4 \\ \end{array} $	
0.	Suppose P, Q and R are different matrices of order $3 \times 5$ , $a \times b$	1
	and $c \times d$ respectively, then value of $ac + bd$ is, if matrix	
	2P + 3Q - 4R is defined	
	(a) 9  (b) 30  (c) 34  (d) 15	
9.	If A is a symmetric matrix then which of the following is not	1
	Symmetric matrix,	1
	$(a)A + A^{T}$ $(b)AA^{T}$ $(c)A - A^{T}$ $(d)A^{T}$	

10.	If $y^{\frac{1}{x}} = a$ , then	$\frac{dy}{dx} =$			1
	(a)y	<i>(b) ay</i>	(c)ax	(d) $y(\log_e a)$	
11.	If a non–singu	lar Matrix A satis	$fy \ 2A^2 + A \ -I = 0$	$O, then A^{-1} =$	1
	(a) 2A - I	(b) 2 $A$ + $I$	(c) 4A + 2I	(d) 2A - 4I	1
12.	$If\left \vec{a}\right  = 13, \left \vec{b}\right  =$	$1 \text{ and } \left  \vec{a} \cdot \vec{b} \right  = 12,$	then $\left  \vec{a} \times \vec{b} \right  =$		1
	(a) 5	( <i>b</i> ) 4	(c) 1	( <i>d</i> ) 3	
13.	<i>The</i> sum of ord	er & degree of th	ne differential equa	ation	1
	$\left \frac{d}{dx}\left(\left(\frac{d^2y}{dx^2}\right)^3\right)\right  =$	= 0 is			-
	(a) 5	<i>(b)</i> 4	<i>(c)</i> 2	( <i>d</i> ) 3	
14.	If the sum of th of their different	e two unit vectors ce is	is a unit vector, the	n the magnitude	1
	( <i>a</i> ) 3	(b) $\sqrt{3}$	(c) 1	(d) $\sqrt{2}$	
15.	If $(2\hat{i} + 6\hat{j} + 9\hat{k})$	$(\hat{k}) \times (\hat{i} + p\hat{j} + q\hat{k}) =$	$=\vec{0}, then \ p+2q =$		1
	( <i>a</i> )10	(b)11	( <i>c</i> )12	( <i>d</i> )13	-
16.	If a line makes $\cos 2\alpha + \cos 2\beta$	$\alpha$ an angle $\alpha$ , $\beta$ , $\beta + \cos 2\gamma =$	$\gamma$ with the axes res	pectively, then	1
	( <i>a</i> ) -1	<i>(b)</i> 0	(c) 1	( <i>d</i> ) 2	
17.	The general so	lution of the dif	ferential equation	$\frac{dx}{dy} = \frac{x}{y}$ is	1
1	(a) x = y + c	(b)x-y=c	$c \qquad (c)xy = c$	$(d)\frac{x}{y} = c$	



21.	Find the principal value of $\sin^{-1}(\sin\frac{3\pi}{5})$ .	2
	OR	
	A relation $R$ in the set of real numbers $R$ is given by	
	$R = \{ (a, b) : a > b, \text{ such that } a, b \in R \}.$	
	Check the Transitivity of Relation R.	
22.	For the curve $y = 5x - 2x^3$ , if x increases at the rate of	2
	2 units / sec, then how fast is the slope of curve changing	
	when $x = 3$ ?	
23.	If $\sin y = x \sin(a + y)$ , then prove that $\frac{dy}{dx} = \frac{\sin^2(a + y)}{\sin a}$ .	2
	OR	
	If $5^x + 5^y = 5^{x+y}$ , then prove that $\frac{dy}{dx} = -5^{y-x}$ .	
24.	If $\vec{a} = \hat{i} + 2\hat{j} - \hat{k}$ and $\vec{b} = \hat{i} - \hat{j} - \hat{k}$ , then Find a unit vector	2
	OPPOSITE to the direction of $(\vec{a} + \vec{b})$ .	
25.	Find the Direction cosines of the line $\frac{x-1}{1} = \frac{2y-6}{4} = \frac{4-z}{1}$ .	2
	SECTION C	
	(This section comprises of short answer type questions (SA) of 3 marks each)	
26.	Solve the following Differential Equation:	2
	$\left(x^2 - y^2\right)dx + 2xy  dy = 0$	
	OR	
	Find the general solution of the differential equation	
	$ydx - \left(x + 2y^2\right)dy = 0$	

27.	Find the intervals in which $f(x) = \sin x + \cos x, x \in [0, 2\pi]$ is	
	(a) strictly Increasing	5
	(b) strictly Decreasing	
28.	If $x^p \cdot y^q = (x+y)^{p+q}$ , Prove that $\frac{dy}{dx} = \frac{y}{x}$ .	3
	OR	
	If $x\sqrt{1+y} + y\sqrt{1+x} = 0$ , $x \neq y$ , then prove that $\frac{dy}{dx} = \frac{-1}{(1+x)^2}$ .	
29.	Evaluate : $I = \int \frac{e^x dx}{e^{2x} - 4e^x + 5}$	3
	OR	
	Evaluate : $I = \int_{-3}^{5}  x - 2  dx$	
20	Solve the following Linear Programming Problem graphically:	2
50.	Minimise $Z = 13x - 15y$ subject to the constraints	5
	$x + y \le 7, \ 2x - 3y + 6 \ge 0, x \ge 0, y \ge 0$	
31.	In a group of 50 scouts in a camp, 30 are well trained in first aid	2
	techniques while the remaining are well trained in hospitality but	
	not in first aid. Two scouts are selected at random from the group.	
	Find the probability distribution of number of selected scouts who	
	are well trained in first aid.	

	OR		
	In a hostel, 60% of the students read Hindi newspaper, 40% read English		
	newspaper and 20% read both Hindi and English newspaper. A student is		
	selected at random.		
	(a) Find the probability that the student reads neither Hindi nor English		
	newspaper.		
	(b) If she reads Hindi newspaper, find the probability that she reads English		
	(c) If she reads English newspaper, find the probability that she reads Hindi		
	newspaper.		
	SECTION D		
	(This section comprises of long answer-type questions (LA) of 5 marks each)		
32.	Show that each of the relation R in the set $A = \{x \in Z : 0 \le x \le 12\}$ ,		
	given by $R = \{(a,b) :   a-b   \text{ is a multiple of 5}\}$		
	Find the set of all elements related to 1 in each case.		
	OR		
	Let $A = R - \{3\}$ and $B = R - \{1\}$ . Consider the function		
	$f: A \to B$ defined by $f(x) = \frac{x-2}{x-3}$ . Is f one – one and onto?		
	Justify your answer.		
33.	[3 4 5]	_	
	Find the inverse of the matrix $2 - 1 = 8$ and hence solve	5	
	5 -2 7		
	the system of equations :		
	3x + 4y + 5z = 18		
	5x - 2y + 7z = 20		
	2x - y + 8z = 13		



(B) If a cm be the width and b cm be the height of the poster, then Expressed the area of the poster in terms of a only.

(C) Find the values of a & b, so that area of the poster is minimized.

37. Read the following passage and answer the questions given below:



A tank with rectangular base and rectangular sides, open at the top is to be constructed so that its depth is 2 m and volume is 8 m<sup>3</sup>. If the building of tank costs Rs 70 per sq. meter for the base and Rs 45 per sq. meter for the sides.

(i) What is the length of the tank for which construction cost is least?(ii) What is the breadth of the tank for which construction cost is least?(iii) What is the least cost of construction?

1 1 2

4



# NOTES

19 <del>1</del>	
	-
2	
22	
-	
÷	
144	
<del>ت</del> ــــــــــــــــــــــــــــــــــــ	12