

केन्द्रीय विद्यालय संगठन, भोपाल संभाग
KENDRIYA VIDYALAYA SANGATHAN, BHOPAL REGION

कक्षा- XII /Class–XII

SESSION/सत्र -2024 -2025

SET-2/सेट-2

Marking Scheme

Mathematics (Code – 041)

Q. No.	Answer	Hints/solution
		Section: A (Multiple Choice Questions- 1 Mark each)
1.	(c)	bijjective function
2.	(a)	$\sin \left[\frac{\pi}{3} + \sin^{-1} \left(\frac{1}{2} \right) \right] = \sin \left(\frac{\pi}{3} + \frac{\pi}{6} \right) = \sin \frac{\pi}{2} = 1$
3.	(a)	<p>For PY to exist $k = 3$ Order of $PY = p \times k$</p> <p>Order of $WY = n \times k$</p> <p>For $PY + WY$ to exist $\text{order}(PY) = \text{order}(WY)$ $\therefore p = n$</p>
4.	(b)	Number of Symmetric matrices of order $3 \times 3 = 2^6 = 64$
5.	(a)	$4 = (3x - 2) - (x + 6)$ $x = 6$
6.	(d)	$ 2A^T = 2^3 A^T = 8 A = 24$
7.	(d)	$\frac{dy}{dt} = 3\cos^2 t.(-\sin t), \quad \frac{dx}{dt} = 3\sin^2 t.(\cos t)$ $\frac{dy}{dx} = \frac{-3\cos^2 t. \sin t}{3\sin^2 t. \cos t} = -\cot t$
8.	(d)	The graph of the function $f: \mathbb{R} \rightarrow \mathbb{Z}$ defined by $f(x) = [x]$; (where $[.]$ denotes G.I.F) is a straight line $\forall x \in (2.5 - h, 2.5 + h)$, 'h' is an infinitesimally small positive quantity. Hence, the function is continuous and differentiable at $x = 2.5$.

9.	(a)	<p>We know, $\int_0^{2a} f(x) dx = 0$, if $f(2a-x) = -f(x)$</p> <p>Let $f(x) = \operatorname{cosec}^7 x$.</p> <p>Now, $f(2\pi - x) = \operatorname{cosec}^7(2\pi - x) = -\operatorname{cosec}^7 x = -f(x)$</p> <p>$\therefore \int_0^{2\pi} \operatorname{cosec}^7 x dx = 0$; Using the property $\int_0^{2a} f(x) dx = 0$, if $f(2a-x) = -f(x)$.</p>
10.	(c)	<p>$\frac{d}{dx} \left(\frac{dy}{dx} \right)^4 = 0 \Rightarrow \frac{d}{dx} (y')^4 = 0$</p> <p>Solving the above, $4y'^3 \times y'' = 0$</p> <p>Order, $m=2$ Degree, $n=1$ and $m+n=3$</p>
11.	(b)	<p>A differential equation of the form $\frac{dy}{dx} = f(x, y)$ is said to be homogeneous, if $f(x, y)$ is a homogeneous function of degree 0.</p> <p>Now, $x^n \frac{dy}{dx} = y \left(\log_e \frac{y}{x} + \log_e e \right) \Rightarrow \frac{dy}{dx} = \frac{y}{x} \left(\log_e e + \left(\frac{y}{x} \right) \right) = f(x, y)$; (Let) $f(x, y)$ will be a homogeneous function of degree 0, if $n=1$.</p>
12.	(c)	<p>$\vec{a} = 3, \vec{b} = 4, \vec{a} + \vec{b} = 5$</p> <p>We have, $\vec{a} + \vec{b} ^2 + \vec{a} - \vec{b} ^2 = 2(\vec{a} ^2 + \vec{b} ^2) = 2(9 + 16) = 50 \Rightarrow \vec{a} - \vec{b} = 5$.</p>
13.	(c)	<p>Any vector in the direction of a vector \vec{a} is given by</p> $\frac{\vec{a}}{ \vec{a} } = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{\sqrt{1^2 + 2^2 + 2^2}} = \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3}$ <p>\therefore Vector in the direction of \vec{a} with magnitude 9 is $9 \cdot \frac{\vec{a}}{ \vec{a} } = 9 \cdot \frac{\hat{i} - 2\hat{j} + 2\hat{k}}{3} = 3(\hat{i} - 2\hat{j} + 2\hat{k})$</p>
14.	(a)	<p>We have, $\vec{a} = 2\hat{i} + 3\hat{j} + 2\hat{k}$ and $\vec{b} = \hat{i} + 2\hat{j} + \hat{k}$</p> <p>$\therefore \vec{a} \cdot \vec{b} = (2\hat{i} + 3\hat{j} + 2\hat{k}) \cdot (\hat{i} + 2\hat{j} + \hat{k}) = 2 + 6 + 2 = 10$</p> <p>and $\vec{b} = \sqrt{1^2 + 2^2 + 1^2} = \sqrt{6}$</p> <p>Hence, projection of \vec{a} on \vec{b} is $\frac{\vec{a} \cdot \vec{b}}{ \vec{b} } = \frac{10}{\sqrt{6}}$.</p>

15.	(b)	We know that $\cos^2\alpha + \cos^2\beta + \cos^2\gamma = 1$ so $\sin^2\alpha + \sin^2\beta + \sin^2\gamma = 2$
16.	(c)	Equation of line $\frac{x-1/3}{1/6} = \frac{y+1/3}{1/3} = \frac{z-1}{1/2}$ so DR of line are 1, 2, 3
17.	(a)	Convex Polygon
18.	(a)	$P\left(\frac{A}{B}\right) = \frac{3}{8}$
19.	(c)	Option (c) is correct, because A is true and R is false
20.	(b)	<p>Answer (b) is correct , because</p> <p>Corner points are (0, 0), (10, 0), (20/3, 20/3) and (0, 10)</p> <p>$Z_{\max} = x + 3y = 0 + 3 \times 10 = 30$</p> <p>both A and R are true but R is not the correct explanation for A</p>


SECTION: B
(VSA Questions of 2 marks each)

21.	$\sin [\cot^{-1}\{\cos(\tan^{-1}1)\}] = \sin [\cot^{-1}\{\cos\frac{\pi}{4}\}]$ $= \sin [\cot^{-1}\frac{1}{\sqrt{2}}]$ $= \sin [\sin^{-1}\frac{\sqrt{2}}{\sqrt{3}}]$ $= \frac{\sqrt{2}}{\sqrt{3}}$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
22.	<p>RHL (at $x = 3$) = $3b + 3$</p> <p>LHL (at $x = 3$) = $3a + 1$</p> <p>$f(3) = 3a + 1$</p> <p>$f(x)$ is continuous at $x = 3$ so $3b + 3 = 3a + 1$</p> <p>Relation between a and b is $3a - 3b = 2$</p>	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

23.(a)	$y = \tan^{-1} x$ and $z = \log_e x$ Then $\frac{dy}{dx} = \frac{1}{1+x^2}$ and $\frac{dz}{dx} = \frac{1}{x}$ So, $\frac{\frac{dy}{dx}}{\frac{dz}{dx}} = \frac{\frac{dy}{dx}}{\frac{1}{x}}$ $= \frac{1}{1+x^2} \cdot \frac{x}{1} = \frac{x}{1+x^2}$ OR Let $y = (\cos x)^x$. log y = x log cos x On differentiating both sides with respect to x , we get $\Rightarrow \frac{dy}{dx} = (\cos x)^x \left\{ \log_e \cos x \frac{d}{dx}(x) + x \frac{d}{dx}(\log_e \cos x) \right\}$ $\Rightarrow \frac{dy}{dx} = (\cos x)^x \left\{ \log_e \cos x + x \cdot \frac{1}{\cos x} (-\sin x) \right\} \Rightarrow \frac{dy}{dx} = (\cos x)^x (\log_e \cos x - x \tan x)$	$\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$
23.(b)	$\Rightarrow \frac{dy}{dx} = (\cos x)^x \left\{ \log_e \cos x \frac{d}{dx}(x) + x \frac{d}{dx}(\log_e \cos x) \right\}$ $\Rightarrow \frac{dy}{dx} = (\cos x)^x \left\{ \log_e \cos x + x \cdot \frac{1}{\cos x} (-\sin x) \right\} \Rightarrow \frac{dy}{dx} = (\cos x)^x (\log_e \cos x - x \tan x)$	$\frac{1}{2}$ $\frac{1}{2}$ 1
24.(a)	We have $\vec{b} + \lambda \vec{c} = (-1 + 3\lambda)\hat{i} + (2 + \lambda)\hat{j} + \hat{k}$ $(\vec{b} + \lambda \vec{c}) \cdot \vec{a} = 0 \Rightarrow 2(-1 + 3\lambda) + 2(2 + \lambda) + 3 = 0$ $\lambda = -\frac{5}{8}$ OR	$\frac{1}{2}$ 1 $\frac{1}{2}$
24.(b)	$\overrightarrow{BA} = \overrightarrow{OA} - \overrightarrow{OB} = (4\hat{i} + 3\hat{k}) - \hat{k} = 4\hat{i} + 2\hat{k}$ Direction cosines are $\frac{2}{\sqrt{5}}, 0, \frac{1}{\sqrt{5}}$	$\frac{1}{2}$ 1.5
25.	Writing the equation of line Putting value Proving perpendicular to z axis	1/2 1/2 1


	SECTION: C (SA Questions of 3 marks each)	
26.(a)	<p>Let $I = \int_{-1}^2 x^3 - x dx$</p> <p>Again let $f(x) = x^3 - x = x(x^2 - 1) = x(x - 1)(x + 1)$ Now break the limit at $x=0, 1$ (because on putting $f(x)=0$ we get $x=0, 1, -1$)</p> <p>It is clear that $x^3 - x \geq 0$ on $[-1, 0]$ $x^3 - x \leq 0$ on $[0, 1]$ $x^3 - x \geq 0$ on $[1, 2]$</p> <p>Hence, the interval of the integral can be subdivided as</p> $\int_{-1}^2 x^3 - x dx = \int_{-1}^0 (x^3 - x) dx - \int_0^1 (x^3 - x) dx + \int_1^2 (x^3 - x) dx$ $= \frac{11}{4}$ <p>OR</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
26.(b)	<p>Ans:</p> <p>Let $I = \int_0^{\pi} \frac{x \sin x}{1 + \cos^2 x} dx \quad \dots (i)$</p> $= \int_0^{\pi} \frac{(\pi - x) \sin(\pi - x)}{1 + [\cos(\pi - x)]^2} dx \quad \left[\because \int_0^a f(x) dx = \int_0^a f(a - x) dx \right] \Rightarrow I = \int_0^{\pi} \frac{(\pi - x) \sin x}{1 + \cos^2 x} dx \quad \dots (ii)$ <p>Adding (i) and (ii), we get $2I = \int_0^{\pi} \frac{\pi \sin x}{1 + \cos^2 x} dx$</p> <p>Put $\cos x = t \Rightarrow -\sin x dx = dt$ Also, $x = 0 \Rightarrow t = 1$ and $x = \pi \Rightarrow t = -1$</p> $\therefore 2I = \int_1^{-1} \frac{-\pi dt}{1 + t^2} \Rightarrow I = \frac{\pi}{2} \int_{-1}^1 \frac{dt}{1 + t^2}$ $\therefore I = \frac{\pi}{2} [\tan^{-1} t]_{-1}^1 = \frac{\pi}{2} [\tan^{-1}(1) - \tan^{-1}(-1)] = \frac{\pi}{2} \left[\frac{\pi}{4} - \left(-\frac{\pi}{4} \right) \right] = \frac{\pi^2}{4}$	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>

27.	<p>Putting $\frac{-x}{2} = t$, we get $x = -2t$ and $dx = -2dt$.</p> $\therefore I = \int \frac{\sqrt{1 - \sin x}}{(1 + \cos x)} e^{-x/2} dx$ $= \int \frac{\sqrt{1 - \sin(-2t)}}{ 1 + \cos(-2t) } e^t (-2dt) = -2 \int \frac{\sqrt{1 + \sin 2t}}{(1 + \cos 2t)} e^t dt$ $= -2 \int \frac{\sqrt{\cos^2 t + \sin^2 t + 2 \sin t \cos t}}{2 \cos^2 t} e^t dt$ $= -2 \int \frac{(\cos t + \sin t)}{2 \cos^2 t} e^t dt = - \int (\sec t + \sec t \tan t) e^t dt$ $= - \int e^t [f(t) + f'(t)] dt, \text{ where } f(t) = \sec t$ $= -e^t f(t) + C = -e^{-x/2} \sec\left(\frac{-x}{2}\right) + C = -e^{-x/2} \sec \frac{x}{2} + C.$	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p>
28.	<p>Putting $\sin x = t$ and $\cos x dx = dt$, we get</p> $I = \frac{\cos x}{(1 - \sin x)(2 - \sin x)} dx = \int \frac{dt}{(1-t)(2-t)}$ <p>Let $\frac{1}{(1-t)(2-t)} = \frac{A}{(1-t)} + \frac{B}{(2-t)}$</p> $\Rightarrow 1 = A(2-t) + B(1-t). \quad \dots (i)$ <p>Putting $t = 1$ in (i), we get $A = 1$.</p> <p>Putting $t = 2$ in (i), we get $B = -1$.</p> $\therefore \frac{1}{(1-t)(2-t)} = \frac{1}{(1-t)} - \frac{1}{(2-t)}$ $\Rightarrow \int \frac{\cos x}{(1 - \sin x)(2 - \sin x)} dx = \int \frac{dt}{(1-t)(2-t)}$ $= \int \left\{ \frac{1}{(1-t)} - \frac{1}{(2-t)} \right\} dt = \int \frac{dt}{(1-t)} - \int \frac{dt}{(2-t)}$ $= -\log 1-t + \log 2-t + C$ $= \log \left \frac{2-t}{1-t} \right + C = \log \left \frac{2 - \sin x}{1 - \sin x} \right + C.$	<p>½</p> <p>½</p> <p>½</p> <p>½</p> <p>1</p>
29.(a)	<p>The given differential equation can be written as</p> $\frac{dy}{dx} = \frac{1+y^2}{(\tan^{-1}y-x)} \text{ or } \frac{dx}{dy} = \frac{(\tan^{-1}y-x)}{1+y^2}$ $\Rightarrow \frac{dx}{dy} + \frac{x}{1+y^2} = \frac{(\tan^{-1}y)}{1+y^2} \text{ linear in } x \text{ where } P = \frac{1}{1+y^2}, Q = \frac{(\tan^{-1}y)}{1+y^2}$ <p>I.F. = $e^{\tan^{-1}y}$</p> <p>GS is $x e^{\tan^{-1}y} = (\tan^{-1}y) e^{\tan^{-1}y} - e^{\tan^{-1}y} + C$ or $x = (\tan^{-1}y) - 1 + C e^{-\tan^{-1}y}$</p>	<p>½+½</p> <p>½</p> <p>1</p>

29.(b)	<p>When $x = 0, y = 0, C = 1$ So PS is $x = (\tan^{-1}y) + e^{-\tan^{-1}y} - 1$</p> <p style="text-align: center;">OR</p> <p>The given differential equation can be written as</p> $\frac{dy}{dx} = \frac{[xy \cos(\frac{y}{x}) + y^2 \sin(\frac{y}{x})]}{[xy \sin(\frac{y}{x}) - x^2 \cos(\frac{y}{x})]}$ <p>On showing this is homogenous equation</p> <p>To solve it, we make the substitution</p> $y = vx$ <p>or</p> $\frac{dy}{dx} = v + x \frac{dv}{dx}$ <p>or</p> $v + x \frac{dv}{dx} = \frac{v \cos v + v^2 \sin v}{v \sin v - \cos v}$ <p>or</p> $x \frac{dv}{dx} = \frac{2v \cos v}{v \sin v - \cos v}$ <p>Solve by variable separation</p> $\log \sec v - \log v = 2 \log x + \log C_1 $ $\frac{\sec(\frac{y}{x})}{(\frac{y}{x})(x^2)} = C \text{ where, } C = \pm C_1$ $\sec(\frac{y}{x}) = C xy$ <p>which is the general solution of the given differential equation.</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
30.	<p>On correct graph of the system of linear inequalities On correct corner points A, B, C and D are (0, 10), (5, 5), (15, 15) and (0, 20)</p> 	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>

	<table><tr><th>Corner Point</th><th colspan="2">Corresponding value of $Z = 3x + 9y$</th></tr><tr><td>A (0, 10)</td><td colspan="2">90</td></tr><tr><td>B (5, 5)</td><td colspan="2">60 (Minimum)</td></tr><tr><td>C (15, 15)</td><td>180 (Maximum)</td><td rowspan="2">(Multiple optimal solutions)</td></tr><tr><td>D (0, 20)</td><td>180 (Maximum)</td></tr></table> <p>Minimum value of Z is 60 at the point B (5, 5) of the feasible region. The maximum value of Z on the feasible region occurs at the two corner points C (15, 15) and D (0, 20) and it is 180 in each case.</p>	Corner Point	Corresponding value of $Z = 3x + 9y$		A (0, 10)	90		B (5, 5)	60 (Minimum)		C (15, 15)	180 (Maximum)	(Multiple optimal solutions)	D (0, 20)	180 (Maximum)	1
Corner Point	Corresponding value of $Z = 3x + 9y$															
A (0, 10)	90															
B (5, 5)	60 (Minimum)															
C (15, 15)	180 (Maximum)	(Multiple optimal solutions)														
D (0, 20)	180 (Maximum)															
31.(a)	<p>Since the event of raining today and not raining today are complementary events so if the probability that it rains today is 0.4 then the probability that it does not rain today is 1- 0.4 = 0.6 $\Rightarrow P_1 = 0.6$</p> <p>If it rains today, the probability that it will rain tomorrow is 0.8 then the probability that it will not rain tomorrow is . 1- 0.8 = 0.2 $\Rightarrow P_2 = 0.2$</p> <p>If it does not rain today, the probability that it will rain tomorrow is 0.7 $\Rightarrow P_3 = 0.7$ then the probability that it will not rain tomorrow is 1 - 0.7 = 0.3 $\Rightarrow P_4 = 0.3$</p> <p>(i) $P_1 \times P_4 - P_2 \times P_3 = 0.6 \times 0.3 - 0.2 \times 0.7 = 0.04$.</p> <p>(ii) Let E_1 and E_2 be the events that it will rain today and it will not rain today respectively.</p> <p>$P(E_1) = 0.4$ & $P(E_2) = 0.6$</p> <p>A be the event that it will rain tomorrow. $P\left(\frac{A}{E_1}\right) = 0.8$ & $P\left(\frac{A}{E_2}\right) = 0.7$</p> <p>We have, $P(A) = P(E_1)P\left(\frac{A}{E_1}\right) + P(E_2)P\left(\frac{A}{E_2}\right) = 0.4 \times 0.8 + 0.6 \times 0.7 = 0.74$.</p> <p>The probability of rain tomorrow is 0.74.</p> <p style="text-align: center;">OR</p> <p>Let S denote the success (getting a '6') and F denote the failure (not getting a '6'). Thus, $P(S) = \frac{1}{6}$ and $P(F) = \frac{5}{6}$ $P(\text{A wins in the first throw}) = P(S) = \frac{1}{6}$ A gets the third throw, when the first throw by A and second throw by B result into failures. Therefore, $P(\text{A wins in the 3rd throw}) = P(\text{FFS}) = P(F)P(F)P(S) = \frac{5}{6} \times \frac{5}{6} \times \frac{1}{6} = \left(\frac{5}{6}\right)^2 \times \frac{1}{6}$ $P(\text{A wins in the 5th throw}) = P(\text{FFFFS}) = \left(\frac{5}{6}\right)^4 \times \frac{1}{6}$ and so on. Hence, $P(\text{A wins}) = \frac{1}{6} + \left(\frac{5}{6}\right)^2 \times \frac{1}{6} + \left(\frac{5}{6}\right)^4 \times \frac{1}{6} + \dots$ $= \frac{\frac{1}{6}}{1 - \frac{25}{36}} = \frac{6}{11}$ $P(\text{B wins}) = 1 - P(\text{A wins}) = \frac{5}{11}$</p>	1 <														

(Long answer type questions (LA) of 5 marks each)

32.	<p> $A = \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix}, B = \begin{bmatrix} 1 & -1 & 1 \\ 1 & -2 & -2 \\ 2 & 1 & 3 \end{bmatrix}$ </p> <p>Then the product AB is</p> $AB = 8 \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} = 8I$ $\Rightarrow B^{-1} = \frac{1}{8}A$ $\Rightarrow B^{-1} = \frac{1}{8} \begin{bmatrix} -4 & 4 & 4 \\ -7 & 1 & 3 \\ 5 & -3 & -1 \end{bmatrix}$ <p>Write equations in matrix form such a way that coeff. Matrix = B and $C = \begin{bmatrix} 4 \\ 9 \\ 1 \end{bmatrix}$</p> <p>Solving the equation $X = B^{-1}C$, we get</p> $x = 3, y = -2, z = -1$	<p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>2</p>
33.(a)	<p>Let r, h and α be as in Fig. Then $\tan \alpha = \frac{r}{h}$</p>  <p>Given $\tan \alpha = 0.5$ so $\frac{r}{h} = 0.5$</p> $\Rightarrow r = \frac{h}{2}$ <p>Let V be the volume of the cone. Then $V = \frac{1}{3}\pi r^2 h = \frac{\pi h^3}{12}$</p> <p>Therefore $\frac{dV}{dt} = \frac{\pi h^2}{4} \frac{dh}{dt}$</p> <p>Given $\frac{dV}{dt} = 5 \text{ m}^3/\text{h}$ and $h = 4$ so $\frac{dh}{dt} = \frac{5}{4\pi}$ or $\frac{35}{88} \text{ m/h}$</p> <p style="text-align: center;">OR</p> <p>$f(x) = \sin x + \cos x$, so $f'(x) = \cos x - \sin x$</p> <p>Now $f'(x) = 0$ gives $\sin x = \cos x$ which gives that $x = \frac{\pi}{4}, \frac{5\pi}{4}$ as $0 \leq x \leq 2\pi$</p> <p>The points $x = \frac{\pi}{4}, \frac{5\pi}{4}$ divide the interval $[0, 2\pi]$ into three disjoint intervals namely $[0, \frac{\pi}{4})$, $(\frac{\pi}{4}, \frac{5\pi}{4})$, $(\frac{5\pi}{4}, 2\pi]$</p> <p>$f'(x) > 0$ if $x \in [0, \frac{\pi}{4}) \cup (\frac{5\pi}{4}, 2\pi]$ so $f(x) = \sin x + \cos x$ is strictly increasing in $[0, \frac{\pi}{4}) \cup (\frac{5\pi}{4}, 2\pi]$</p> <p>$f'(x) < 0$ if $x \in (\frac{\pi}{4}, \frac{5\pi}{4})$ so $f(x) = \sin x + \cos x$ is strictly decreasing in $(\frac{\pi}{4}, \frac{5\pi}{4})$</p>	<p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1+1/2</p> <p>1</p>
33.(b)		

[illegible]

	$\frac{x-1}{32/7} = \frac{y-2}{-34/7} = \frac{z-1}{12/7} \Rightarrow \frac{x-1}{16} = \frac{y-2}{-17} = \frac{z-1}{6},$ <p style="text-align: center;">OR</p>	1
35.(b)	<p>The given equation of line can be written as</p> $\vec{r} = \hat{i} + (-2)\hat{j} + (3)\hat{k} + t(-\hat{i} + \hat{j} + (-2)\hat{k}) \text{ and }$ $\vec{r} = \hat{i} + (-1)\hat{j} - \hat{k} + s(\hat{i} + (2)\hat{j} - (2)\hat{k})$ <p>So $\vec{a}_1 = \hat{i} + (-2)\hat{j} + (3)\hat{k}$, $\vec{a}_2 = \hat{i} + (-1)\hat{j} - \hat{k}$</p> $\vec{b}_1 = -\hat{i} + \hat{j} + (-2)\hat{k}, \quad \vec{b}_2 = \hat{i} + (2)\hat{j} - (2)\hat{k}$ $\vec{b}_1 \times \vec{b}_2 = 2\hat{i} - 4\hat{j} - 3\hat{k}$ $\vec{a}_2 - \vec{a}_1 = \hat{j} - 4\hat{k}$ $SD = \frac{8}{\sqrt{29}}$ <p>So the given lines does not intersect to each other</p>	<p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p>

SECTION E

(Case Studies/Passage based questions of 4 Marks each)

36.	<p>(i) Number of relations is equal to the number of subsets of the set $B \times G = 2^{n(B \times G)}$</p> $= 2^{n(B) \times n(G)} = 2^{3 \times 2} = 2^6$ <p>(Where $n(A)$ denotes the number of the elements in the finite set A)</p> <p>(ii) Smallest Equivalence relation on G is $\{(g_1, g_1), (g_2, g_2)\}$</p> <p>(iii) (a) (A) reflexive but not symmetric =</p> $\{(b_1, b_2), (b_2, b_1), (b_1, b_1), (b_2, b_2), (b_3, b_3), (b_2, b_3)\}.$ <hr/> <p>So the minimum number of elements to be added are</p> $(b_1, b_1), (b_2, b_2), (b_3, b_3), (b_2, b_3)$ <p>{Note : it can be any one of the pair from, (b_3, b_2), (b_1, b_3), (b_3, b_1) in place of (b_2, b_3) also}</p> <p>(B) reflexive and symmetric but not transitive =</p> $\{(b_1, b_2), (b_2, b_1), (b_1, b_1), (b_2, b_2), (b_3, b_3), (b_2, b_3), (b_3, b_2)\}.$ <hr/> <p>So the minimum number of elements to be added are</p> $(b_1, b_1), (b_2, b_2), (b_3, b_3), (b_2, b_3), (b_3, b_2)$ <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>
-----	---	-------------------------------------

	<p>One-one and onto function</p> <p>$x^2 = 4y$. let $y = f(x) = \frac{x^2}{4}$</p> <p>Let $x_1, x_2 \in [0, 20\sqrt{2}]$ such that $f(x_1) = f(x_2) \Rightarrow \frac{x_1^2}{4} = \frac{x_2^2}{4}$</p> <p>$\Rightarrow x_1^2 = x_2^2 \Rightarrow (x_1 - x_2)(x_1 + x_2) = 0 \Rightarrow x_1 = x_2$ as $x_1, x_2 \in [0, 20\sqrt{2}]$</p> <p>$\therefore f$ is one-one function</p> <p>Now, $0 \leq y \leq 200$ hence the value of y is non-negative and $f(2\sqrt{y}) = y$</p> <p>\therefore for any arbitrary $y \in [0, 200]$, the pre-image of y exists in $[0, 20\sqrt{2}]$ hence f is onto function.</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>
37.	<p>(i) Let E denote the event that the person selected is actually having HIV and A the event that the person's HIV test is diagnosed as +ive. We need to find $P(A E)$.</p> <p>$P(A E) = P(\text{Person tested as HIV+ive given that he/she is actually having HIV})$</p> <p>$= 90\% = 90/100 = 0.9$</p> <p>(ii)(a) We need to find $P(E A)$. Also F denotes the event that the person selected is actually not having HIV</p> <p>Then $P(E) = 0.001$, $P(F) = 1 - P(E) = 0.999$,</p> <p>$P(A E) = P(\text{Person tested as HIV+ive given that he/she is actually having HIV})$</p> <p>$= 90\% = 90/100 = 0.9$</p> <p>$P(A F) = P(\text{Person tested as HIV +ive given that he/she is actually not having HIV})$</p> <p>$= 1\% = 0.01$</p> <p>Now, by Bayes' theorem</p> <p>$P(E A) = \frac{P(E)P(A E)}{P(E)P(A E) + P(F)P(A F)}$</p> <p>$= \frac{90}{1089} = 0.083$ approx.</p>	<p>2</p> <p>1</p> <p>1</p>

38.	<p>(i) Let length of the side of square base be x cm and height of the box be y cm \therefore volume of box, $V = x^2y = 1024$</p> <p>(ii) Let C denotes the cost of the box $C = 2x^2 \times 5 + 4xy \times 2.5$</p> $\Rightarrow C(x) = 10x^2 + \frac{10240}{x}$ <p>(iii)(a) On differentiating both sides w.r.t. x we get</p> $C'(x) = 20x - \frac{10240}{x^2}$ $C''(x) = 20 + \frac{20480}{x^3}$ <p>Now, $C'(x) = 0 \Rightarrow x = 8$</p> <p>As, $C''(8) > 0$. So, at $x = 8$ the function has minimum value.</p> <p>For $x=8$ cost is minimum and the corresponding least cost of the box is:</p> $C(8) = 10 \times 64 + \frac{10240}{8} = 1920$ <p>\therefore least cost =Rs 1920</p> <p style="text-align: center;">OR</p> <p>(iii)(b) Dimensions of required box are $8(2^{1/3})$cm, $8(2^{1/3})$cm and $8(2^{1/3})$cm.</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>2</p>
-----	---	--