

केंद्रीय विद्यालय संगठन, अहमदाबाद संभाग  
**KENDRIYA VIDYALAYA SANGATHAN, AHMEDABAD REGION**  
 प्री-बोर्ड परीक्षा 2025-26  
**PRE BOARD-I 2025-26**

**SUBJECT: MATHEMATICS (041)**  
**CLASS: XI**

**M.M.: 80**  
**TIME: 3 HOURS**


**ANSWER KEY**  
**SET-1 (A)**

**SECTION A**

1	(c) $\frac{\pi}{3}$	1
2	(b) 64	1
3	(a) 8	1
4	(d) a scalar matrix	1
5	(b) =1	1
6	(d) $R - \{4\}$	1
7	(c) $-2\sqrt{\pi}$	1
8	(b) 2	1
9	<b>(b)</b> $\tan(xe^x) + C$	1
10	(d) $(2, \infty)$	1
11	(b) 0	1
12	(c) (3,3)	1
13	(d) $q=3p$	1
14	(c) 12	1
15	(c) $\frac{\pi}{3}$	1
16	(b) $\vec{a} \perp \vec{b}$	1
17	(d) (4,0)	1
18	(c) $\frac{2}{3}$	1
19	(d) Assertion(A) is false but reason(R) is true	1
20	(c) Assertion (A) is true but reason(R) is false.	1

	<b>SECTION B</b>	
21	<p>Substitute  <math>x = \tan(\theta)</math>  <math>\sin^{-1}\left(\frac{\tan \theta}{\sec \theta}\right) =</math>  <math>\sin^{-1}(\sin \theta) = \theta</math>  <math>= \tan^{-1} x</math>  <b>OR</b>  <math>-1 \leq -x^2 \leq 1</math>  <math>(i) x^2 \geq -1</math>  This is true for all real numbers, as the square of any real number is non-negative.  <math>(ii) -1 \leq -x^2</math>  <math>1 \geq x^2,</math>  <math>-1 \leq x \leq 1</math></p> <p>The domain of the function is <b><math>[-1, 1]</math></b></p>	<p>1</p> <p>1</p> <p>0.5</p> <p>0.5</p> <p>1</p>
22	<p>Taking log both side  <math>\log x = \frac{y}{x}</math>  Differentiate both side w.r.t. x  <math>\frac{1}{x} = \frac{y - x \frac{dy}{dx}}{y^2}</math>  <math>\frac{dy}{dx} = \frac{xy - y^2}{x^2} = \frac{y(x - y)}{x^2}</math>  Substitute  <math>y = \frac{x}{\log x}</math>  <math>\frac{dy}{dx} = \frac{x - y}{x \log x}</math></p>	<p>1</p> <p>1</p>
23	<p><math>u = 2^{\cos^2 x}, v = \cos^2 x</math>  <math>\frac{du}{dx} = 2^{\cos^2 x} (\log 2) (-2 \cos x \sin x), \frac{dv}{dx} = (-2 \cos x \sin x)</math>  <math>\frac{du}{dv} = \frac{\frac{du}{dx}}{\frac{dv}{dx}} = 2^{\cos^2 x} (\log 2)</math></p>	<p>1</p> <p>1</p>
24	<p>Put <math>\sqrt{x} = t, dx = 2t dt</math>  <math>\int_0^{\frac{\pi}{2}} \sin t dt</math>  <math>= 2[-\cos t]_0^{\frac{\pi}{2}} = 2(0 - 1) = -2</math>  <b>OR</b>  Put <math>1 + 2x = t^2, x = \frac{t^2 - 1}{2}</math>  <math>2dx = 2t dt, dx = t dt</math>  <math>= \int \frac{t^2 - 1}{2} t dt = \frac{1}{2} \int t^4 - t^2 dt</math>  <math>= \frac{1}{2} \left( \frac{t^5}{5} - \frac{t^3}{3} \right) + c</math>  <math>\frac{1}{2} \left( \frac{(1 + 2x)^{5/2}}{5} - \frac{(1 + 2x)^{3/2}}{3} \right) + c</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>



	$\frac{d^2y}{dx^2} = \frac{\sec^3\theta \tan\theta}{a^2}$ $\frac{d^2y}{dx^2} \text{ at } \theta = \frac{\pi}{4} = \frac{2\sqrt{2}}{a^2}$ <p>OR</p> $y = (\tan^{-1}x)^2$ <p>Differentiate w.r.t x</p> $\frac{dy}{dx} = \frac{2\tan^{-1}x}{1+x^2}$ $(1+x^2)\frac{dy}{dx} = 2\tan^{-1}x$ <p>Again differentiate w.r.t x</p> $(x^2+1)y_2 + 2xy_1 = \frac{2}{1+x^2}$ $(x^2+1)^2y_2 + 2x(x^2+1)y_1 = 2$	0.5 0.5  1 0.5  1 0.5
29	 <p>Area = <math>4 \int_0^2 y dx = 4 \int_0^2 \frac{1}{2} \sqrt{4^2 - x^2} dx</math></p> $= 2 \left[ \frac{x}{2} \sqrt{4^2 - x^2} + 8 \sin^{-1} \left( \frac{x}{4} \right) \right]_0^2$ $2 \left[ \sqrt{12} + \frac{8\pi}{6} \right] = 4\sqrt{3} + \frac{8\pi}{3}$ <p style="text-align: center;">Type equation here.</p> <p>OR</p> <p>Required area <math>A = \int_0^4 x(4-x) dx + \left  \int_4^5 x(4-x) dx \right </math></p> $A = \left[ 2x^2 - \frac{x^3}{3} \right]_0^4 + \left  \left[ 2x^2 - \frac{x^3}{3} \right]_4^5 \right $ $= 32/3 + 7/3 = 13 \text{ sq. units.}$	1 1 1  1 1 1
30	<p>That the dr<sup>o</sup> of given lines are not proportional so, they are not parallel lines.</p> $(a_2 - a_1) = \hat{j} - 4\hat{k}$ $(b_1 \times b_2) = 2\hat{i} - 4\hat{j} - 3\hat{k}$ <p>Consider <math>(a_2 - a_1) \cdot (b_1 \times b_2) = 8 \neq 0</math></p> <p>Hence line will not intersect. So the lines are skew.</p> <p>Shortest distance = <math>\left  \frac{(a_2 - a_1) \cdot (b_1 \times b_2)}{ b_1 \times b_2 } \right  = \frac{8}{\sqrt{4+16+9}} = \frac{8}{\sqrt{29}} \text{ units}</math></p> <p>OR</p> <p>Let the wicket keeper divides the line segment in ratio k:1</p> $\vec{W} = \frac{k \cdot \vec{F} + 1 \cdot \vec{B}}{k+1}$ $6\hat{i} + 12\hat{j} = \left( \frac{12k+2}{k+1} \right) \hat{i} + \left( \frac{18k+8}{k+1} \right) \hat{j}$	1/2  1/2 1/2    1/2 1   1 1

	On comparing the components, we get $6 = \left(\frac{12k+2}{k+1}\right), \quad k = \frac{2}{3}$	1
31	<p>E= the first die showed an even number.  F= the sum of the numbers on the dice is 9= {(6,3), (3,6), (5,4), (4,5)}</p> <p><math>E \cap F = \{(6,3), (4,5)\}</math>  <math>n(S) = 36, n(E) = 18, n(F) = 4, n(E \cap F) = 2</math>  <math>P\left(\frac{E}{F}\right) = \frac{P(E \cap F)}{P(F)} = \frac{2/36}{4/36} = \frac{1}{2}</math> Type equation here.</p>	1 1 1
	SECTION D	
32	<p><b>Solution</b> The given differential equation can be written as</p> $\frac{dx}{dy} + \frac{x}{1+y^2} = \frac{\tan^{-1}y}{1+y^2} \quad \dots (1)$ <p>Now (1) is a linear differential equation of the form <math>\frac{dx}{dy} + P_1 x = Q_1</math>,</p> <p>where, <math>P_1 = \frac{1}{1+y^2}</math> and <math>Q_1 = \frac{\tan^{-1}y}{1+y^2}</math>.</p> <p>Therefore, <math>I.F = e^{\int \frac{1}{1+y^2} dy} = e^{\tan^{-1}y}</math></p> <p>Thus, the solution of the given differential equation is</p> $x e^{\tan^{-1}y} = \int \left( \frac{\tan^{-1}y}{1+y^2} \right) e^{\tan^{-1}y} dy + C \quad \dots (2)$ <p>Let <math>I = \int \left( \frac{\tan^{-1}y}{1+y^2} \right) e^{\tan^{-1}y} dy</math></p> <p>Substituting <math>\tan^{-1}y = t</math> so that <math>\left( \frac{1}{1+y^2} \right) dy = dt</math>, we get</p> $I = \int t e^t dt = t e^t - \int 1 \cdot e^t dt = t e^t - e^t = e^t (t - 1)$ <p>or <math>I = e^{\tan^{-1}y} (\tan^{-1}y - 1)</math></p> <p>Substituting the value of I in equation (2), we get</p> $x \cdot e^{\tan^{-1}y} = e^{\tan^{-1}y} (\tan^{-1}y - 1) + C$ <p>or <math>x = (\tan^{-1}y - 1) + C e^{-\tan^{-1}y}</math></p> <p>which is the general solution of the given differential equation.</p>	1   1   1   1   1
33	<p><math>(kA) \left( \frac{1}{k} A^{-1} \right) = k \frac{1}{k} (A A^{-1}) = I</math>  <math>(kA)^{-1} = \frac{1}{k} A^{-1}</math>.</p> <p>Hence calculate <math>(3A)^{-1} = \frac{1}{3} A^{-1}</math>,</p> $A = \begin{bmatrix} 2 & -1 & 1 \\ -1 & 2 & -1 \\ 1 & -1 & 2 \end{bmatrix}$ <p><math>\det(A) = 4 \neq 0, A^{-1}</math> exist</p> $\text{adj}A = \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$	1 1/2  1/2 2

	$A^{-1} = \frac{1}{4} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$ $(3A)^{-1} = \frac{1}{3} A^{-1} = \frac{1}{12} \begin{bmatrix} 3 & 1 & -1 \\ 1 & 3 & 1 \\ -1 & 1 & 3 \end{bmatrix}$	$\frac{1}{2}$  $\frac{1}{2}$
34	<p> <math display="block">I = \int \frac{\cos x}{(4 + \sin^2 x)(5 - 4 \cos^2 x)} dx</math> <math display="block">= \int \frac{\cos x}{(4 + \sin^2 x)(1 + 4 \sin^2 x)} dx</math> Let, <math>\sin x = t \Rightarrow \cos x dx = dt</math> <math display="block">I = \int \frac{dt}{(4 + t^2)(1 + 4t^2)}</math> <math display="block">= -\frac{1}{15} \int \frac{dt}{4 + t^2} + \frac{4}{15} \int \frac{dt}{1 + 4t^2}</math> <math display="block">(\because \text{using partial fraction})</math> For the second term, let <math>u = 2t</math>, so <math>du = 2dt</math>. <math display="block">I = -\frac{1}{15} \cdot \frac{1}{2} \tan^{-1}\left(\frac{t}{2}\right) + \frac{4}{15} \int \frac{1}{1 + u^2} \frac{du}{2}</math> <math display="block">I = -\frac{1}{30} \tan^{-1}\left(\frac{t}{2}\right) + \frac{2}{15} \tan^{-1}(u) + C</math> <math display="block">I = -\frac{1}{30} \tan^{-1}\left(\frac{\sin x}{2}\right) + \frac{2}{15} \tan^{-1}(2 \sin x) + C</math> Substitute back <math>t = \sin x</math>. <math display="block">I = -\frac{1}{30} \tan^{-1}\left(\frac{\sin x}{2}\right) + \frac{2}{15} \tan^{-1}(2 \sin x) + C</math> </p> <p>OR</p> <p> <math display="block">I = \int_0^{\pi} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}</math> <math display="block">= 2 \int_0^{\frac{\pi}{2}} \frac{dx}{a^2 \cos^2 x + b^2 \sin^2 x}</math> <math display="block">= 2 \int_0^{\frac{\pi}{2}} \frac{\sec^2 x}{a^2 + b^2 \tan^2 x} dx</math> <math display="block">\tan x = t</math> <math display="block">\sec^2 x dx = dt</math> <math display="block">x = 0 \Rightarrow t = 0</math> <math display="block">x = \frac{\pi}{2} \Rightarrow t = \infty</math> <math display="block">I = 2 \int_0^{\infty} \frac{dt}{a^2 + b^2 t^2}</math> <math display="block">= \frac{2}{b^2} \cdot \frac{b}{a} \left[ \tan^{-1}\left(\frac{bt}{a}\right) \right]_0^{\infty}</math> <math display="block">= \frac{2}{ab} \left( \frac{\pi}{2} - 0 \right) = \frac{\pi}{ab}</math> </p>	1  1  1  1    1.5    1  1  1+0.5

35	$\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$ <p>Any arbitrary point on line is <math>M(\lambda, 2\lambda+1, 3\lambda+2)</math>          Dir's of AM are <math>\langle \lambda-1, 2\lambda-5, 3\lambda-1 \rangle</math>          AM perpendicular to line 1  <math>\lambda = 1</math>  <math>\therefore M(1, 3, 5)</math> is the foot of perpendicular of the point A to the given line.          Let the image of point A in the line be <math>A'(\alpha, \beta, \gamma)</math>          Since M is the mid point of <math>AA'</math>,  <math>M\left(\frac{1+\alpha}{2}, \frac{6+\beta}{2}, \frac{3+\gamma}{2}\right) = M(1, 3, 5)</math>  <math>A'(1, 0, 7)</math>          Also, Equation of <math>AA'</math> is <math>\frac{x-1}{0} = \frac{y-6}{-6} = \frac{z-3}{4}</math></p> <p style="text-align: center;"><b>OR</b></p> <p>line <math>\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}</math>,          any random point on the line will be given by <math>P(\lambda-5, 4\lambda-3, -9\lambda+6)</math>          Since <math>PQ=7</math>  <math>\sqrt{(\lambda-7)^2 + (4\lambda-7)^2 + (-9\lambda+7)^2} = 7</math>  <math>\lambda = 1</math>  <math>P(-4, 1, -3)</math>          The Equation of line PQ is  <math>\frac{x+4}{6} = \frac{y-1}{3} = \frac{z+3}{2}</math> or <math>\frac{x-2}{6} = \frac{y-4}{3} = \frac{z+1}{2}</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
	<b>CASE STUDY</b>	
36	<p>(i) No. of possible relation from S to J = <math>2^{12} = 4096</math>          ( <math>n(S \times J) = 12</math> )</p> <p>(ii) Condition: <math> S  \leq  J </math> (hear, <math>4 &gt; 3</math>)          It is impossible to assign all speakers to distinct judges.          one-one functions can be there from set S to set J=0</p> <p>(iii) Given function from S to J as  <math>f = \{(S_1, J_1), (S_2, J_2), (S_3, J_2), (S_4, J_3)\}</math>  <math>S_2</math> and <math>S_3</math> are both assigned to <math>J_2 \rightarrow</math> not injective          All judges <math>J = \{J_1, J_2, J_3\}</math> are assigned <math>\rightarrow</math> Surjective          Since f is not bijective</p> <p style="text-align: center;"><b>OR</b></p> <p>relation <math>R_1 = \{(S_1, S_2), (S_2, S_4)\}</math>          Add reflexive relation pairs <math>(S_1, S_1), (S_2, S_2), (S_3, S_3), (S_4, S_4)</math>          Keep <math>(S_1, S_2)</math> but excluded <math>(S_2, S_1)</math>,          Final relation <math>R_1 = \{(S_1, S_1), (S_2, S_2), (S_3, S_3), (S_4, S_4), (S_1, S_2), (S_2, S_4)\}</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>2</p>

37	<p>(i) Capacity=area×depth = <math>x^2h = 250</math>  <math>Cost (C) = 500x^2 + 4000h^2</math>  <math>C = 500\left(\frac{250}{h}\right) + 4000h^2</math></p> <p>(ii) <math>\frac{dC}{dh} = -\frac{125000}{h^2} + 8000h</math>  <math>\frac{dC}{dh} = 0, h = \frac{5}{2}m</math> or 2.5 m</p> <p>(iii) <math>\left(\frac{d^2C}{dh^2}\right)_{h=2.5} &gt; 0</math>  Cost is minimum when h=2.5 m  Minimum cost = <math>C = \frac{125000}{\frac{5}{2}} + 4000\left(\frac{5}{2}\right)^2 = \text{Rs } 75,000</math></p> <p style="text-align: center;"><b>OR</b></p> <p>h=2.5 m when <math>\frac{dC}{dh} = 0</math>  For value of h less than 5/2 and closed o 5/2, <math>\frac{dC}{dh} &lt; 0</math>  For value of h less more than 5/2 and closed o 5/2, <math>\frac{dC}{dh} &gt; 0</math>  By first derivative test , C is minimum at h=5/2  Now <math>x^2 = \frac{250}{h}, x = 10 m, \text{ also, } x = 4h</math></p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
38	<p>Let <math>E_1</math>=customer avails loan on fixed rate  <math>E_2</math>=customer avails loan on floating rate  <math>E_3</math>=customer avails loan on variable rate  A= Person defaults on the loan  <math>P(E_1) = \frac{1}{10}, P(E_2) = \frac{2}{10}, P(E_3) = \frac{7}{10},</math>  <math>P(A/E_1) = \frac{15}{100}, P(A/E_2) = \frac{3}{100}, P(A/E_3) = \frac{1}{100},</math>  (i) <math>P(A) = P(E_1)P(A/E_1) + P(E_2)P(A/E_2) + P(E_3)P(A/E_3) = 9/500</math></p> <p>(ii) <math>P\left(\frac{E_3}{A}\right) = \frac{P(E_3)P(A/E_3)}{P(E_1)P(A/E_1) + P(E_2)P(A/E_2) + P(E_3)P(A/E_3)} = 7/18</math></p>	<p>1+1</p> <p>1+1</p>