

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025
SUBJECT NAME MATHEMATICS (Q.P. CODE – 65/1/1)


General Instructions: -

| | |
|----------|--|
| 1 | You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully. |
| 2 | “Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.” |
| 3 | Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded. |
| 4 | The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly. |
| 5 | The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators. |
| 6 | Evaluators will mark (√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing. |
| 7 | If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly. |
| 8 | If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly. |
| 9 | If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” . |

| | |
|----|---|
| 10 | No marks to be deducted for the cumulative effect of an error. It should be penalized only once. |
| 11 | A full scale of marks _____ (example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it. |
| 12 | Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper. |
| 13 | <p>Ensure that you do not make the following common types of errors committed by the Examiner in the past: -</p> <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks |
| 14 | While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks. |
| 15 | Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously. |
| 16 | The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation. |
| 17 | Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words. |
| 18 | The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme. |

| | | |
|------------|---|---|
| Q4. | Which of the following is <u>not</u> a homogeneous function of x and y ? (A) $y^2 - xy$ (B) $x - 3y$ (C) $\sin^2 \frac{y}{x} + \frac{y}{x}$ (D) $\tan x - \sec y$ | |
| Ans | (D) $\tan x - \sec y$ | 1 |
| Q5. | If $f(x) = x + x - 1 $, then which of the following is correct ? (A) $f(x)$ is both continuous and differentiable, at $x = 0$ and $x = 1$. (B) $f(x)$ is differentiable but not continuous, at $x = 0$ and $x = 1$. (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. (D) $f(x)$ is neither continuous nor differentiable, at $x = 0$ and $x = 1$. | |
| Ans | (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. | 1 |
| Q6. | If A is a square matrix of order 2 such that $\det(A) = 4$, then $\det(4 \operatorname{adj} A)$ is equal to : (A) 16 (B) 64 (C) 256 (D) 512 | |
| Ans | (B) 64 | 1 |
| Q7. | If E and F are two independent events such that $P(E) = \frac{2}{3}$, $P(F) = \frac{3}{7}$, then $P(E/\bar{F})$ is equal to : (A) $\frac{1}{6}$ (B) $\frac{1}{2}$ (C) $\frac{2}{3}$ (D) $\frac{7}{9}$ | |
| Ans | (C) $\frac{2}{3}$ | 1 |
| Q8. | The absolute maximum value of function $f(x) = x^3 - 3x + 2$ in $[0, 2]$ is : (A) 0 (B) 2 (C) 4 (D) 5 | |
| Ans | (C) 4 | 1 |

| | | |
|--------------------|---|----------|
| <p>Q9.</p> | <p>Let $A = \begin{bmatrix} 1 & -2 & -1 \\ 0 & 4 & -1 \\ -3 & 2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} -2 \\ -5 \\ -7 \end{bmatrix}$, $C = [9 \ 8 \ 7]$, which of the following is defined ?</p> <p>(A) Only AB (B) Only AC (C) Only BA (D) All AB, AC and BA</p> | |
| <p>Ans</p> | <p>(A) Only AB</p> | <p>1</p> |
| <p>Q10.</p> | <p>If $\int \frac{2^x}{x^2} dx = k \cdot 2^{\frac{1}{x}} + C$, then k is equal to</p> <p>(A) $\frac{-1}{\log 2}$ (B) $-\log 2$ (C) -1 (D) $\frac{1}{2}$</p> | |
| <p>Ans</p> | <p>(A) $\frac{-1}{\log 2}$</p> | <p>1</p> |
| <p>Q11.</p> | <p>If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, $\vec{a} = \sqrt{37}$, $\vec{b} = 3$ and $\vec{c} = 4$, then angle between \vec{b} and \vec{c} is</p> <p>(A) $\frac{\pi}{6}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$</p> | |
| <p>Ans</p> | <p>(C) $\frac{\pi}{3}$</p> | <p>1</p> |
| <p>Q12.</p> | <p>The integrating factor of differential equation $(x + 2y^3) \frac{dy}{dx} = 2y$ is</p> <p>(A) $e^{\frac{y^2}{2}}$ (B) $\frac{1}{\sqrt{y}}$ (C) $\frac{1}{y^2}$ (D) $e^{-\frac{1}{y^2}}$</p> | |
| <p>Ans</p> | <p>(B) $\frac{1}{\sqrt{y}}$</p> | <p>1</p> |

| | | |
|-------------|--|----------|
| Q13. | <p>If $A = \begin{bmatrix} 7 & 0 & x \\ 0 & 7 & 0 \\ 0 & 0 & y \end{bmatrix}$ is a scalar matrix, then y^x is equal to</p> <p>(A) 0 (B) 1 (C) 7 (D) ± 7</p> | |
| Ans | (B) 1 | 1 |
| Q14. | <p>The corner points of the feasible region in graphical representation of a L.P.P. are (2, 72), (15, 20) and (40, 15). If $Z = 18x + 9y$ be the objective function, then</p> <p>(A) Z is maximum at (2, 72), minimum at (15, 20) (B) Z is maximum at (15, 20) minimum at (40, 15) (C) Z is maximum at (40, 15), minimum at (15, 20) (D) Z is maximum at (40, 15), minimum at (2, 72)</p> | |
| Ans | (C) Z is maximum at (40, 15), minimum at (15, 20) | 1 |
| Q15. | <p>If A and B are invertible matrices, then which of the following is <u>not</u> correct ?</p> <p>(A) $(A + B)^{-1} = B^{-1} + A^{-1}$ (B) $(AB)^{-1} = B^{-1}A^{-1}$ (C) $\text{adj}(A) = A A^{-1}$ (D) $A ^{-1} = A^{-1}$</p> | |
| Ans | (A) $(A + B)^{-1} = B^{-1} + A^{-1}$ | 1 |
| Q16. | <p>If the feasible region of a linear programming problem with objective function $Z = ax + by$, is bounded, then which of the following is correct ?</p> <p>(A) It will only have a maximum value. (B) It will only have a minimum value. (C) It will have both maximum and minimum values. (D) It will have neither maximum nor minimum value.</p> | |
| Ans | (C) It will have both maximum and minimum values. | 1 |
| Q17. | <p>The area of the shaded region bounded by the curves $y^2 = x$, $x = 4$ and the x-axis is given by</p>  <p>(A) $\int_0^4 x \, dx$ (B) $\int_0^2 y^2 \, dy$ (C) $2 \int_0^4 \sqrt{x} \, dx$ (D) $\int_0^4 \sqrt{x} \, dx$</p> | |
| Ans | (D) $\int_0^4 \sqrt{x} \, dx$ | 1 |

| | |
|--------------------|--|
| <p>Q18.</p> | <p>The graph of a trigonometric function is as shown. Which of the following will represent graph of its inverse ?</p> <p>(A) </p> <p>(B) </p> <p>(C) </p> <p>(D) </p> |
|--------------------|--|

| | | |
|-------------------|-------------|----------|
| <p>Ans</p> | <p>(C) </p> | <p>1</p> |
|-------------------|-------------|----------|

Assertion – Reason Based Questions

Direction : Question numbers 19 and 20 are Assertion (A) and Reason (R) based questions carrying 1 mark each. Two statements are given, one labelled Assertion (A) and other labelled Reason (R). Select the correct answer from the options (A), (B), (C) and (D) as given below.

- (A) Both Assertion (A) and Reason (R) are true and the Reason (R) is the correct explanation of the Assertion (A).
- (B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).
- (C) Assertion (A) is true, but Reason (R) is false.
- (D) Assertion (A) is false, but Reason (R) is true.

| | | |
|--------------------|--|-------------------|
| <p>Q19.</p> | <p>Assertion (A) : Let Z be the set of integers. A function $f : Z \rightarrow Z$ defined as $f(x) = 3x - 5, \forall x \in Z$ is a bijective.</p> <p>Reason (R) : A function is a bijective if it is both surjective and injective.</p> | <p>Ans</p> |
| <p>Ans</p> | <p>(D) Assertion (A) is false, but Reason (R) is true.</p> | <p>1</p> |

| | | |
|------|--|---|
| Q20. | <p>Assertion (A) : $f(x) = \begin{cases} 3x - 8, & x \leq 5 \\ 2k, & x > 5 \end{cases}$</p> <p>is continuous at $x = 5$ for $k = \frac{5}{2}$.</p> <p>Reason (R) : For a function f to be continuous at $x = a$,</p> $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = f(a).$ | |
| Ans | (D) Assertion (A) is false, but Reason (R) is true. | 1 |

SECTION B

This section comprises very short answer (VSA) type questions of **2 marks each**.

| | | |
|------|---|--|
| Q21. | <p>(a) Differentiate $2^{\cos^2 x}$ w.r.t $\cos^2 x$.</p> <p style="text-align: center;">OR</p> <p>(b) If $\tan^{-1}(x^2 + y^2) = a^2$, then find $\frac{dy}{dx}$.</p> | |
|------|---|--|

| | | |
|--------|---|--------------------------------|
| Ans(a) | <p>Let $u = 2^{\cos^2 x} \Rightarrow \frac{du}{dx} = 2^{\cos^2 x} (-2 \cos x \sin x) \log 2$</p> <p>Let $v = \cos^2 x \Rightarrow \frac{dv}{dx} = -2 \cos x \sin x$</p> <p>Now $\frac{du}{dv} = \frac{\left(\frac{du}{dx}\right)}{\left(\frac{dv}{dx}\right)} = 2^{\cos^2 x} \log 2$</p> | <p>1</p> <p>1/2</p> <p>1/2</p> |
|--------|---|--------------------------------|

OR

| | | |
|--------|--|--------------------------------|
| Ans(b) | <p>$\tan^{-1}(x^2 + y^2) = a^2 \Rightarrow x^2 + y^2 = \tan a^2$</p> <p>Differentiate both sides wrt x,</p> $2x + 2y \frac{dy}{dx} = 0$ $\Rightarrow \frac{dy}{dx} = -\frac{x}{y}$ | <p>1/2</p> <p>1</p> <p>1/2</p> |
|--------|--|--------------------------------|

| | | |
|------|---|--|
| Q22. | Evaluate : $\tan^{-1} \left[2 \sin \left(2 \cos^{-1} \frac{\sqrt{3}}{2} \right) \right]$ | |
| Ans | $\tan^{-1} \left[2 \sin \left(2 \cos^{-1} \frac{\sqrt{3}}{2} \right) \right]$ $= \tan^{-1} \left[2 \sin \left(2 \times \frac{\pi}{6} \right) \right] = \tan^{-1} \left[2 \sin \frac{\pi}{3} \right]$ $= \tan^{-1} \left[2 \times \frac{\sqrt{3}}{2} \right] = \tan^{-1} \sqrt{3} = \frac{\pi}{3}$ | <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> |
| Q23. | The diagonals of a parallelogram are given by $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + 3\hat{j} - \hat{k}$. Find the area of the parallelogram. | |
| Ans | $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 1 \\ 1 & 3 & -1 \end{vmatrix} = -2\hat{i} + 3\hat{j} + 7\hat{k}$ <p style="text-align: center;">Area of parallelogram = $\frac{1}{2} \vec{a} \times \vec{b}$</p> $= \frac{1}{2} \sqrt{(-2)^2 + 3^2 + 7^2} = \frac{\sqrt{62}}{2}$ | <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> |
| Q24. | Find the intervals in which function $f(x) = 5x^{\frac{3}{2}} - 3x^{\frac{5}{2}}$ is (i) increasing (ii) decreasing. | |
| Ans | $f(x) = 5x^{3/2} - 3x^{5/2} \Rightarrow f'(x) = \frac{15}{2} \sqrt{x} (1-x)$ <p>For increasing / decreasing, put $f'(x) = 0$</p> <p>$\Rightarrow x = 0, 1$</p> <p>(i) When $x \in [0, 1], f'(x) \geq 0$. So, f is increasing when $x \in [0, 1]$ (The intervals $(0, 1), [0, 1)$ or $(0, 1]$ can also be considered.)</p> <p>(ii) When $x \in [1, \infty), f'(x) \leq 0$. So, f is decreasing when $x \in [1, \infty)$ (The interval $(1, \infty)$ can also be considered.)</p> | <p style="text-align: center;">1</p> <p style="text-align: center;">1/2</p> <p style="text-align: center;">1/2</p> |

| | |
|-------------|---|
| Q25. | <p>(a) Two friends while flying kites from different locations, find the strings of their kites crossing each other. The strings can be represented by vectors $\vec{a} = 3\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 2\hat{i} - 2\hat{j} + 4\hat{k}$. Determine the angle formed between the kite strings. Assume there is no slack in the strings.</p> <p style="text-align: center;">OR</p> <p>(b) Find a vector of magnitude 21 units in the direction opposite to that of \vec{AB} where A and B are the points A(2, 1, 3) and B(8, -1, 0) respectively.</p> |
|-------------|---|

| | | |
|---------------|--|--------------------|
| Ans(a) | <p>Let the required angle between the kite strings be θ.</p> <p>Then, $\cos\theta = \frac{\vec{a} \cdot \vec{b}}{ \vec{a} \vec{b} }$</p> $\Rightarrow \cos\theta = \frac{(3\hat{i} + \hat{j} + 2\hat{k})(2\hat{i} - 2\hat{j} + 4\hat{k})}{\sqrt{9+1+4} \sqrt{4+4+16}} = \frac{12}{\sqrt{336}} = \frac{3}{\sqrt{21}}$ $\Rightarrow \theta = \cos^{-1}\left(\frac{12}{\sqrt{336}}\right) \text{ or } \cos^{-1}\left(\frac{3}{\sqrt{21}}\right)$ | <p>1½</p> <p>½</p> |
|---------------|--|--------------------|

OR

| | | |
|---------------|--|----------------------------|
| Ans(b) | <p>$\vec{BA} = -6\hat{i} + 2\hat{j} + 3\hat{k}$</p> <p>Required unit vector of magnitude 21</p> $= 21 \times \left(\frac{-6\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{36+4+9}} \right)$ $= 3(-6\hat{i} + 2\hat{j} + 3\hat{k}) \text{ or } -18\hat{i} + 6\hat{j} + 9\hat{k}$ | <p>1</p> <p>½</p> <p>½</p> |
|---------------|--|----------------------------|

SECTION C

This section comprises short answer (SA) type questions of **3 marks each**.

| | | |
|-------------|--|-----------------------------|
| Q26. | <p>The side of an equilateral triangle is increasing at the rate of 3 cm/s. At what rate its area increasing when the side of the triangle is 15 cm ?</p> | |
| Ans | <p>Let 'a' be the side of the triangle, so $\frac{da}{dt} = 3 \text{ cm/s}$</p> <p>Now area of an equilateral triangle, $A = \frac{\sqrt{3}}{4} a^2$</p> $\Rightarrow \frac{dA}{dt} = \frac{\sqrt{3}a}{2} \times \frac{da}{dt}$ $\therefore \left. \frac{dA}{dt} \right _{a=15 \text{ cm}} = \frac{\sqrt{3} \times 15}{2} \times 3 = \frac{45\sqrt{3}}{2} \text{ cm}^2/\text{s}$ | <p>½</p> <p>1½</p> <p>1</p> |

Q27.

Solve the following linear programming problem graphically :

Maximise $Z = x + 2y$


Subject to the constraints :

$$x - y \geq 0$$

$$x - 2y \geq -2$$

$$x \geq 0, y \geq 0$$

Ans



| Corner Point | Value of $Z = x + 2y$ |
|--------------|-----------------------|
| $O(0,0)$ | 0 |
| $A(2,2)$ | 6 |

Since feasible region is unbounded. Plot $x + 2y > 6$ which has common region with feasible region, thus Z has no maximum value.

For correct graph and shading
1½

For correct table
1

½


Ans

(a) Find : $\int \frac{x + \sin x}{1 + \cos x} dx$


OR

(b) Evaluate : $\int_0^{\frac{\pi}{4}} \frac{dx}{\cos^3 x \sqrt{2 \sin 2x}}$

| | | |
|------------------|--|---|
| <p>Ans(a)</p> | $\int \frac{x + \sin x}{1 + \cos x} dx$ $= \int \frac{x + 2\sin \frac{x}{2} \cos \frac{x}{2}}{2\cos^2 \frac{x}{2}} dx$ $= \int x \left(\frac{1}{2} \sec^2 \frac{x}{2} \right) dx + \int \tan \frac{x}{2} dx$ $= x \tan \frac{x}{2} - \int \tan \frac{x}{2} dx + \int \tan \frac{x}{2} dx$ $= x \tan \frac{x}{2} + C$ | <p>1</p> <p>1/2</p> <p>1</p> <p>1/2</p> |
| <p>OR</p> | | |
| <p>Ans(b)</p> | $\int_0^{\pi/4} \frac{dx}{\cos^3 x \sqrt{2\sin 2x}}$ $= \frac{1}{2} \int_0^{\pi/4} \frac{dx}{\cos^4 x \sqrt{\tan x}}$ $= \frac{1}{2} \int_0^{\pi/4} \frac{(1 + \tan^2 x) \sec^2 x}{\sqrt{\tan x}} dx$ <p>Put $\tan x = t \Rightarrow \sec^2 x dx = dt$</p> $\therefore I = \frac{1}{2} \int_0^1 \frac{1+t^2}{\sqrt{t}} dt$ $= \frac{1}{2} \int_0^1 \left(\frac{1}{\sqrt{t}} + t^{3/2} \right) dt$ $= \frac{1}{2} \left[2\sqrt{t} + \frac{2}{5} t^{5/2} \right]_0^1$ $= \frac{6}{5}$ | <p>1/2</p> <p>1/2</p> <p>1</p> <p>1/2</p> |

| | | |
|----------------------|--|---|
| <p>Q29.</p> | <p>(a) Verify that lines given by $\vec{r} = (1 - \lambda)\hat{i} + (\lambda - 2)\hat{j} + (3 - 2\lambda)\hat{k}$ and $\vec{r} = (\mu + 1)\hat{i} + (2\mu - 1)\hat{j} - (2\mu + 1)\hat{k}$ are skew lines. Hence, find shortest distance between the lines.</p> <p style="text-align: center;">OR</p> <p>(b) During a cricket match, the position of the bowler, the wicket keeper and the leg slip fielder are in a line given by $\vec{B} = 2\hat{i} + 8\hat{j}$, $\vec{W} = 6\hat{i} + 12\hat{j}$ and $\vec{F} = 12\hat{i} + 18\hat{j}$ respectively. Calculate the ratio in which the wicketkeeper divides the line segment joining the bowler and the leg slip fielder.</p> | |
| <p>Ans(a)</p> | <p>Rewriting the lines, we get</p> $\vec{r} = (\hat{i} - 2\hat{j} + 3\hat{k}) + \lambda(-\hat{i} + \hat{j} - 2\hat{k}) \text{ and } \vec{r} = (\hat{i} - \hat{j} - \hat{k}) + \mu(\hat{i} + 2\hat{j} - 2\hat{k})$ <p>Let $\vec{a}_1 = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{a}_2 = \hat{i} - \hat{j} - \hat{k}$, $\vec{b}_1 = -\hat{i} + \hat{j} - 2\hat{k}$, $\vec{b}_2 = \hat{i} + 2\hat{j} - 2\hat{k}$</p> <p>Note that the dr's of given lines are not proportional so, they are not parallel lines. The lines will be skew if they do not intersect each other also.</p> <p>Here $\vec{a}_2 - \vec{a}_1 = \hat{j} - 4\hat{k}$, $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & -2 \\ 1 & 2 & -2 \end{vmatrix} = 2\hat{i} - 4\hat{j} - 3\hat{k}$</p> <p>Consider $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)$</p> $= (\hat{j} - 4\hat{k}) \cdot (2\hat{i} - 4\hat{j} - 3\hat{k}) = 8 \neq 0$ <p>Hence lines will not intersect. So the lines are skew.</p> <p>Shortest Distance = $\frac{ (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) }{ \vec{b}_1 \times \vec{b}_2 }$</p> $= \frac{8}{\sqrt{4 + 16 + 9}} = \frac{8}{\sqrt{29}}$ | <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2 + 1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> |
| OR | | |
| <p>Ans(b)</p> | <p>Let the wicket keeper divides the line segment in ratio $k : 1$</p> $\therefore \vec{W} = \frac{k\vec{F} + 1\vec{B}}{k + 1}$ $\Rightarrow 6\hat{i} + 12\hat{j} = \left(\frac{12k + 2}{k + 1}\right)\hat{i} + \left(\frac{18k + 8}{k + 1}\right)\hat{j}$ $\Rightarrow k = \frac{2}{3}$ <p>Hence, the required ratio is 2 : 3</p> <div style="text-align: center;">  </div> | <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> |

| | | | | | | | | | | | | |
|----------------------|---|---|----|---|---|---|-------------|---|----|----|---|--|
| <p>Q30.</p> | <p>(a) The probability distribution for the number of students being absent in a class on a Saturday is as follows :</p> <table border="1" data-bbox="304 237 783 353"> <tbody> <tr> <td>X</td> <td>0</td> <td>2</td> <td>4</td> <td>5</td> </tr> <tr> <td>P(X)</td> <td>p</td> <td>2p</td> <td>3p</td> <td>p</td> </tr> </tbody> </table> <p>Where X is the number of students absent.</p> <p>(i) Calculate p. 1</p> <p>(ii) Calculate the mean of the number of absent students on Saturday. 2</p> <p style="text-align: center;">OR</p> <p>(b) For the vacancy advertised in the newspaper, 3000 candidates submitted their applications. From the data it was revealed that two third of the total applicants were females and other were males. The selection for the job was done through a written test. The performance of the applicants indicates that the probability of a male getting a distinction in written test is 0.4 and that a female getting a distinction is 0.35. Find the probability that the candidate chosen at random will have a distinction in the written test.</p> | X | 0 | 2 | 4 | 5 | P(X) | p | 2p | 3p | p | |
| X | 0 | 2 | 4 | 5 | | | | | | | | |
| P(X) | p | 2p | 3p | p | | | | | | | | |
| <p>Ans(a)</p> | <p>(i) Since $\sum P(X) = 1 \Rightarrow p + 2p + 3p + p = 1$ $\Rightarrow p = \frac{1}{7}$</p> <p>(ii) Mean = $\sum X.P(X) = 0(p) + 2(2p) + 4(3p) + 5(p)$ $= 21p = 21\left(\frac{1}{7}\right) = 3$</p> | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> | | | | | | | | | | |
| OR | | | | | | | | | | | | |
| <p>Ans(b)</p> | <p>Let E_1 : The applicant is a male E_2 : The applicant is a female A : The candidate chosen will have distinction in the written test.</p> <p>$P(E_1) = \frac{1}{3}, P(E_2) = \frac{2}{3}, P(A E_1) = 0.4, P(A E_2) = 0.35$</p> <p>$\therefore P(A) = P(E_1)P(A E_1) + P(E_2)P(A E_2)$ $= \frac{1}{3} \times 0.4 + \frac{2}{3} \times 0.35$ $= \frac{11}{30}$</p> | <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> | | | | | | | | | | |

| | | | |
|-------------|---|--|---|
| Q31. | Sketch the graph of $y = x + 3 $ and find the area of the region enclosed by the curve, x -axis, between $x = -6$ and $x = 0$, using integration. | | |
| Ans | <p>Required Area</p> $= \int_{-6}^0 y \, dx$ $= 2 \int_{-3}^0 (x + 3) \, dx$ $= 2 \left[\frac{(x + 3)^2}{2} \right]_{-3}^0$ $= 9$ |  | <p>For correct graph: 1 mark</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |

SECTION D


This section comprises long answer (LA) type questions of **5 marks each**.

| | | | |
|-------------|---|--|--|
| Q32. | <p>(a) If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$, then prove that $\frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$.</p> <p align="center">OR</p> <p>(b) If $x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$ and $y = \sin \theta$, then find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{4}$.</p> | | |
|-------------|---|--|--|

| | | |
|---------------|--|--|
| Ans(a) | <p>Let $x = \sin A, y = \sin B \Rightarrow A = \sin^{-1} x, B = \sin^{-1} y$</p> <p>$\therefore \sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$</p> <p>$\Rightarrow \cos A + \cos B = a(\sin A - \sin B)$</p> <p>$\Rightarrow 2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) = 2a \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right)$</p> <p>$\Rightarrow \cot \left(\frac{A-B}{2} \right) = a \Rightarrow A - B = 2 \cot^{-1} a$</p> <p>$\Rightarrow \sin^{-1} x - \sin^{-1} y = 2 \cot^{-1} a$</p> <p>differentiate both sides wrt x,</p> $\frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = 0$ <p>$\Rightarrow \frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1/2</p> <p>1 1/2</p> |
|---------------|--|--|

OR

| | | |
|---------------|--|--|
| <p>Ans(b)</p> | $x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$ $\Rightarrow \frac{dx}{d\theta} = a \left(-\sin \theta + \frac{1}{\tan \frac{\theta}{2}} \times \sec^2 \frac{\theta}{2} \times \frac{1}{2} \right)$ $= a \left(-\sin \theta + \frac{1}{\sin \theta} \right) = a \left(\frac{1 - \sin^2 \theta}{\sin \theta} \right)$ $\frac{dx}{d\theta} = a \cot \theta \cos \theta$ <p>Also, $y = \sin \theta \Rightarrow \frac{dy}{d\theta} = \cos \theta$</p> $\therefore \frac{dy}{dx} = \frac{\tan \theta}{a}$ <p>Differentiating wrt x,</p> $\frac{d^2 y}{dx^2} = \frac{\sec^2 \theta}{a} \times \frac{d\theta}{dx}$ $= \frac{\sec^3 \theta \tan \theta}{a^2}$ $\left. \frac{d^2 y}{dx^2} \right]_{\text{at } \theta = \frac{\pi}{4}} = \frac{2\sqrt{2}}{a^2}$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> |
| <p>Q33.</p> | <p>Find the absolute maximum and absolute minimum of function $f(x) = 2x^3 - 15x^2 + 36x + 1$ on $[1, 5]$.</p> | |
| <p>Ans</p> | $f(x) = 2x^3 - 15x^2 + 36x + 1$ $\Rightarrow f'(x) = 6(x^2 - 5x + 6) = 6(x - 2)(x - 3)$ $f'(x) = 0 \Rightarrow x = 2, 3 \in [1, 5]$ <p>Now $f(1) = 24, f(2) = 29, f(3) = 28, f(5) = 56$</p> <p>Hence, the absolute maximum value is 56 and the absolute minimum value is 24.</p> | <p>1</p> <p>1</p> <p>2</p> <p>1</p> |
| <p>Q34.</p> | <p>(a) Find the image A' of the point A(1, 6, 3) in the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$. Also, find the equation of the line joining A and A'.</p> <p style="text-align: center;">OR</p> <p>(b) Find a point P on the line $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}$ such that its distance from point Q(2, 4, -1) is 7 units. Also, find the equation of line joining P and Q.</p> | |


| | | |
|----------------------|---|--|
| <p>Ans(a)</p> | <p>The equation of given line is $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda$</p> <p>Any arbitrary point on the line is $M(\lambda, 2\lambda+1, 3\lambda+2)$</p> <p>dr's of AM are $\langle \lambda-1, 2\lambda-5, 3\lambda-1 \rangle$</p> <p>Here $1(\lambda-1) + 2(2\lambda-5) + 3(3\lambda-1) = 0$</p> <p>$\Rightarrow \lambda = 1$</p> <p>$\therefore M(1, 3, 5)$ is the foot perpendicular of the point A to the given line.</p> <p>Let image of point A in the line be $A'(\alpha, \beta, \gamma)$</p> <p>Since M is the mid-point of AA', so $M\left(\frac{1+\alpha}{2}, \frac{6+\beta}{2}, \frac{3+\gamma}{2}\right) = M(1, 3, 5)$</p> <p>$\Rightarrow A'(1, 0, 7)$ is the image of A.</p> <p>Also, Equation of AA' is $\frac{x-1}{0} = \frac{y-6}{-3} = \frac{z-3}{2}$</p> |  <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p> |
| <p>OR</p> | | |
| <p>Ans(b)</p> | <p>The given line is $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9} = \lambda$ and $Q(2, 4, -1)$</p> <p>Any random point on the line will be given by $P(\lambda-5, 4\lambda-3, -9\lambda+6)$</p> <p>Since $PQ = 7 \Rightarrow \sqrt{(\lambda-7)^2 + (4\lambda-7)^2 + (-9\lambda+7)^2} = 7$</p> <p>$\Rightarrow 98(\lambda^2 - 2\lambda + 1) = 0 \Rightarrow \lambda = 1$</p> <p>Hence, the required point is $P(-4, 1, -3)$</p> <p>The equation of line PQ is $\frac{x+4}{6} = \frac{y-1}{3} = \frac{z+3}{2}$ or $\frac{x-2}{6} = \frac{y-4}{3} = \frac{z+1}{2}$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
| <p>Q35.</p> | <p>A school wants to allocate students into three clubs : Sports, Music and Drama, under following conditions :</p> <ul style="list-style-type: none"> The number of students in Sports club should be equal to the sum of the number of students in Music and Drama club. The number of students in Music club should be 20 more than half the number of students in Sports club. The total number of students to be allocated in all three clubs are 180. <p>Find the number of students allocated to different clubs, using matrix method.</p> | |

| | | |
|------------|---|--|
| Ans | <p>Let x, y and z be the no. of students allocated to Sports, Music and Drama clubs respectively.</p> <p>Here, $x = y + z, y = \frac{x}{2} + 20, x + y + z = 180$</p> <p>$\Rightarrow x - y - z = 0, x - 2y = -40, x + y + z = 180$</p> <p>Given equations can be written as $AX = B$</p> <p>where, $A = \begin{bmatrix} 1 & -1 & -1 \\ 1 & -2 & 0 \\ 1 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$</p> <p>$A = -4 \neq 0 \Rightarrow A^{-1}$ exists.</p> <p>$adjA = \begin{bmatrix} -2 & 0 & -2 \\ -1 & 2 & -1 \\ 3 & -2 & -1 \end{bmatrix}$</p> <p>$A^{-1} = \frac{1}{ A } \times adjA = \frac{1}{4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix}$</p> <p>$X = A^{-1}B$</p> <p>$= \frac{1}{4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix} = \begin{bmatrix} 90 \\ 65 \\ 25 \end{bmatrix}$</p> <p>$\therefore x = 90, y = 65, z = 25$</p> <p>Number of students allocated in sports, music and drama are 90 , 65 and 25 respectively .</p> | <p>1½</p> <p>½</p> <p>½</p> <p>1</p> <p>½</p> <p>1</p> |
|------------|---|--|

SECTION E

This section comprises 3 case study-based questions of 4 marks each.

Q36.




A technical company is designing a rectangular solar panel installation on a roof using 300 metres of boundary material. The design includes a partition running parallel to one of the sides dividing the area (roof) into two sections.

Let the length of the side perpendicular to the partition be x metres and with parallel to the partition be y metres.

| | | |
|-------------------|--|--|
| | <p>Based on this information, answer the following questions :</p> <p>(i) Write the equation for the total boundary material used in the boundary and parallel to the partition in terms of x and y. 1</p> <p>(ii) Write the area of the solar panel as a function of x. 1</p> <p>(iii) (a) Find the critical points of the area function. Use second derivative test to determine critical points at the maximum area. Also, find the maximum area. 2</p> <p style="text-align: center;">OR</p> <p>(iii) (b) Using first derivative test, calculate the maximum area the company can enclose with the 300 metres of boundary material, considering the parallel partition. 2</p> | |
| <p>Ans</p> | <p>(i) $2x + 3y = 300$ 1</p> <p>(ii) $A = xy = \frac{x}{3}(300 - 2x)$ 1</p> <p>(iii) (a) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$</p> <p>$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$ 1/2</p> <p>For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$ 1/2</p> <p>Also, $\frac{d^2A}{dx^2} = -\frac{4}{3} < 0$. So, A is maximum at $x = 75$ 1/2</p> <p>Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$ 1/2</p> <p style="text-align: center;">OR</p> <p>(iii) (b) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$</p> <p>$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$ 1/2</p> <p>For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$ 1/2</p> <p>As $\frac{dA}{dx}$ changes its sign from positive to negative as x passes through $x = 75$ from left to right, which means $x = 75$ is the point of maximum. 1/2</p> <p>Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$ 1/2</p> <p>Note : Full credit to be given if the student takes equation as $2x + 2y = 300$ or $2x + 4y = 300$ or $4x + 4y = 300$ or $4x + 3y = 300$</p> <p>The solutions of sub-parts will differ and marks may be given accordingly.</p> | |

| | | |
|--------------------|--|---|
| <p>Q37.</p> | <p>A class-room teacher is keen to assess the learning of her students the concept of “relations” taught to them. She writes the following five relations each defined on the set $A = \{1, 2, 3\}$:</p> <p>$R_1 = \{(2, 3), (3, 2)\}$</p> <p>$R_2 = \{(1, 2), (1, 3), (3, 2)\}$</p> <p>$R_3 = \{(1, 2), (2, 1), (1, 1)\}$</p> <p>$R_4 = \{(1, 1), (1, 2), (3, 3), (2, 2)\}$</p> <p>$R_5 = \{(1, 1), (1, 2), (3, 3), (2, 2), (2, 1), (2, 3), (3, 2)\}$</p> <p>The students are asked to answer the following questions about the above relations :</p> <p>(i) Identify the relation which is reflexive, transitive but not symmetric.</p> <p>(ii) Identify the relation which is reflexive and symmetric but not transitive.</p> <p>(iii) (a) Identify the relations which are symmetric but neither reflexive nor transitive.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) What pairs should be added to the relation R_2 to make it an equivalence relation ?</p> | |
| <p>Ans</p> | <p>(i) R_4</p> <p>(ii) R_5</p> <p>(iii)(a) R_1 and R_3</p> <p style="text-align: center;">OR</p> <p>(iii)(b) Required pairs to be added to make the relation R_2 as an equivalence relation are: $(1,1), (2,2), (3,3), (2,1), (3,1)$ and $(2,3)$</p> | <p>1</p> <p>1</p> <p>1+1</p> <p>2</p> |

Q38.



A bank offers loan to its customers on different types of interest namely, fixed rate, floating rate and variable rate. From the past data with the bank, it is known that a customer avails loan on fixed rate, floating rate or variable rate with probabilities 10%, 20% and 70% respectively. A customer after availing loan can pay the loan or default on loan repayment. The bank data suggests that the probability that a person defaults on loan after availing it at fixed rate, floating rate and variable rate is 5%, 3% and 1% respectively.

Based on the above information, answer the following :

- (i) What is the probability that a customer after availing the loan will default on the loan repayment ? 2
- (ii) A customer after availing the loan, defaults on loan repayment. What is the probability that he availed the loan at a variable rate of interest ? 2

| | | |
|-------------------|--|-------------------------------------|
| <p>Ans</p> | <p>E_1 :customer avails loan on fixed rate E_2 :customer avails loan on floating rate E_3 :customer avails loan on variable rate A:the person defaults on the loan</p> <p>$P(E_1) = \frac{1}{10}, P(E_2) = \frac{2}{10}, P(E_3) = \frac{7}{10}$</p> <p>$P(A E_1) = \frac{5}{100}, P(A E_2) = \frac{3}{100}, P(A E_3) = \frac{1}{100}$</p> <p>(i) $P(A) = P(E_1) \cdot P(A E_1) + P(E_2) \cdot P(A E_2) + P(E_3) \cdot P(A E_3)$</p> $= \frac{1}{10} \times \frac{5}{100} + \frac{2}{10} \times \frac{3}{100} + \frac{7}{10} \times \frac{1}{100}$ $= \frac{18}{1000} \text{ or } \frac{9}{500}$ <p>(ii) $P(E_3 A) = \frac{P(E_3) \cdot P(A E_3)}{P(E_1) \cdot P(A E_1) + P(E_2) \cdot P(A E_2) + P(E_3) \cdot P(A E_3)}$</p> $= \frac{\frac{7}{10} \times \frac{1}{100}}{\frac{18}{1000}}$ $= \frac{7}{18}$ | <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
|-------------------|--|-------------------------------------|

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025

SUBJECT NAME MATHEMATICS (Q.P. CODE – 65/1/2)

General Instructions: -

| | |
|----------|--|
| 1 | You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully. |
| 2 | “Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.” |
| 3 | Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded. |
| 4 | The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly. |
| 5 | The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators. |
| 6 | Evaluators will mark (\surd) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (\checkmark) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing. |
| 7 | If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly. |
| 8 | If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly. |
| 9 | If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” . |

| | |
|----|---|
| 10 | No marks to be deducted for the cumulative effect of an error. It should be penalized only once. |
| 11 | A full scale of marks _____ (example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it. |
| 12 | Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper. |
| 13 | <p>Ensure that you do not make the following common types of errors committed by the Examiner in the past: -</p> <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks |
| 14 | While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks. |
| 15 | Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously. |
| 16 | The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation. |
| 17 | Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words. |
| 18 | The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme. |

| | |
|-------------|---|
| Q7. | If $f(x) = x + x - 1 $, then which of the following is correct ? (A) $f(x)$ is both continuous and differentiable, at $x = 0$ and $x = 1$. (B) $f(x)$ is differentiable but not continuous, at $x = 0$ and $x = 1$. (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. (D) $f(x)$ is neither continuous nor differentiable, at $x = 0$ and $x = 1$. |
| Ans | (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. 1 |
| Q8. | Which of the following is <u>not</u> a homogeneous function of x and y ? (A) $y^2 - xy$ (B) $x - 3y$ (C) $\sin^2 \frac{y}{x} + \frac{y}{x}$ (D) $\tan x - \sec y$ |
| Ans | (D) $\tan x - \sec y$ 1 |
| Q9. | Let A be a matrix of order $m \times n$ and B is a matrix such that $A^T B$ and BA^T are defined. Then, the order of B is : (A) $m \times m$ (B) $n \times n$ (C) $m \times n$ (D) $n \times m$ |
| Ans | (C) $m \times n$ 1 |
| Q10. | If the feasible region of a linear programming problem with objective function $Z = ax + by$, is bounded, then which of the following is correct ? (A) It will only have a maximum value. (B) It will only have a minimum value. (C) It will have both maximum and minimum values. (D) It will have neither maximum nor minimum value. |
| Ans | (C) It will have both maximum and minimum values. 1 |

| | | |
|--------------------|--|-----------------|
| <p>Q11.</p> | <p>If $A = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then A^{-1} is</p> <p>(A) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ (B) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$</p> <p>(C) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (D) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$</p> | |
| <p>Ans</p> | <p>(D) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$</p> | <p>1</p> |
| <p>Q12.</p> | <p>The integrating factor of the differential equation $\frac{dy}{dx} + y = \frac{1+y}{x}$ is</p> <p>(A) xe^x (B) $\frac{e^x}{x}$</p> <p>(C) $\frac{x}{e^x}$ (D) $xe^{\frac{1}{x}}$</p> | |
| <p>Ans</p> | <p>(B) $\frac{e^x}{x}$</p> | <p>1</p> |
| <p>Q13.</p> | <p>Let $A = [a_{ij}]$ be a square matrix of order 3 such that $a_{ij} = \hat{j} - 2\hat{i}$. Then which of the following is true ?</p> <p>(A) $a_{12} > 0$ (B) all $a_{ij} < 0$</p> <p>(C) $a_{13} + a_{31} = -6$ (D) $a_{23} > a_{32}$</p> | |
| <p>Ans</p> | <p>(D) $a_{23} > a_{32}$</p> | <p>1</p> |
| <p>Q14.</p> | <p>The absolute maximum value of function $f(x) = x^3 - 3x + 2$ in $[0, 2]$ is :</p> <p>(A) 0 (B) 2</p> <p>(C) 4 (D) 5</p> | |
| <p>Ans</p> | <p>(C) 4</p> | <p>1</p> |

Assertion – Reason Based Questions

Direction : Question numbers 19 and 20 are Assertion (A) and Reason (R) based questions carrying 1 mark each. Two statements are given, one labelled Assertion (A) and other labelled Reason (R). Select the correct answer from the options (A), (B), (C) and (D) as given below.

- (A) Both Assertion (A) and Reason (R) are true and the Reason (R) is the correct explanation of the Assertion (A).
- (B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).
- (C) Assertion (A) is true, but Reason (R) is false.
- (D) Assertion (A) is false, but Reason (R) is true.

| | |
|-------------|--|
| Q19. | <p>Assertion (A) : Let Z be the set of integers. A function $f : Z \rightarrow Z$ defined as $f(x) = 3x - 5, \forall x \in Z$ is a bijective.</p> <p>Reason (R) : A function is a bijective if it is both surjective and injective.</p> |
|-------------|--|

| | | |
|------------|---|----------|
| Ans | (D) Assertion (A) is false, but Reason (R) is true. | 1 |
|------------|---|----------|

| | |
|-------------|--|
| Q20. | <p>Assertion (A) : $f(x) = \begin{cases} 3x - 8, & x \leq 5 \\ 2k, & x > 5 \end{cases}$</p> <p style="text-align: center;">is continuous at $x = 5$ for $k = \frac{5}{2}$.</p> <p>Reason (R) : For a function f to be continuous at $x = a$,</p> $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = f(a).$ |
|-------------|--|

| | | |
|------------|---|----------|
| Ans | (D) Assertion (A) is false, but Reason (R) is true. | 1 |
|------------|---|----------|

SECTION B

This section comprises very short answer (VSA) type questions of 2 marks each.

| | |
|-------------|---|
| Q21. | <p>(a) Two friends while flying kites from different locations, find the strings of their kites crossing each other. The strings can be represented by vectors $\vec{a} = 3\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 2\hat{i} - 2\hat{j} + 4\hat{k}$. Determine the angle formed between the kite strings. Assume there is no slack in the strings.</p> <p style="text-align: center;">OR</p> <p>(b) Find a vector of magnitude 21 units in the direction opposite to that of \vec{AB} where A and B are the points A(2, 1, 3) and B(8, -1, 0) respectively.</p> |
|-------------|---|

| | | |
|-----------|--|----------------------------|
| Ans (a) | <p>Let the required angle between the kite strings be θ.</p> <p>Then, $\cos \theta = \frac{\vec{a} \cdot \vec{b}}{ \vec{a} \vec{b} }$</p> $\Rightarrow \cos \theta = \frac{(3\hat{i} + \hat{j} + 2\hat{k})(2\hat{i} - 2\hat{j} + 4\hat{k})}{\sqrt{9+1+4} \sqrt{4+4+16}} = \frac{12}{\sqrt{336}} = \frac{3}{\sqrt{21}}$ $\Rightarrow \theta = \cos^{-1} \left(\frac{12}{\sqrt{336}} \right) \text{ or } \cos^{-1} \left(\frac{3}{\sqrt{21}} \right)$ | <p>1½</p> <p>½</p> |
| OR | | |
| Ans(b) | <p>$\vec{BA} = -6\hat{i} + 2\hat{j} + 3\hat{k}$</p> <p>Required unit vector of magnitude 21</p> $= 21 \times \left(\frac{-6\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{36+4+9}} \right)$ $= 3(-6\hat{i} + 2\hat{j} + 3\hat{k}) \text{ or } -18\hat{i} + 6\hat{j} + 9\hat{k}$ | <p>1</p> <p>½</p> <p>½</p> |
| Q22. | Find the values of 'a' for which $f(x) = x^2 - 2ax + b$ is an increasing function for $x > 0$. | |
| A22. | <p>$f'(x) = 2x - 2a$</p> <p>$0 < x < \infty \Rightarrow -2a < 2x - 2a < \infty \Rightarrow -2a < f'(x) < \infty$</p> <p>$f(x)$ is increasing iff $f'(x) \geq 0$</p> <p>$\Rightarrow -2a \in [0, \infty) \Rightarrow a \in (-\infty, 0] \text{ or } (-\infty, 0)$</p> | <p>1</p> <p>½</p> <p>½</p> |
| Q23. | <p>(a) Differentiate $2^{\cos^2 x}$ w.r.t $\cos^2 x$.</p> <p style="text-align: center;">OR</p> <p>(b) If $\tan^{-1}(x^2 + y^2) = a^2$, then find $\frac{dy}{dx}$.</p> | |
| Ans(a) | <p>Let $u = 2^{\cos^2 x} \Rightarrow \frac{du}{dx} = 2^{\cos^2 x} (-2 \cos x \sin x) \log 2$</p> <p>Let $v = \cos^2 x \Rightarrow \frac{dv}{dx} = -2 \cos x \sin x$</p> <p>Now $\frac{du}{dv} = \frac{\left(\frac{du}{dx} \right)}{\left(\frac{dv}{dx} \right)} = 2^{\cos^2 x} \log 2$</p> | <p>1</p> <p>½</p> <p>½</p> |

OR


| | | |
|---|--|---------------------------------------|
| Ans(b) | $\tan^{-1}(x^2 + y^2) = a^2 \Rightarrow x^2 + y^2 = \tan a^2$ <p>Differentiate both sides wrt x,</p> $2x + 2y \frac{dy}{dx} = 0$ $\Rightarrow \frac{dy}{dx} = -\frac{x}{y}$ | <p>1/2</p> <p>1</p> <p>1/2</p> |
| Q24. | Evaluate : $\sin^{-1}\left(\sin \frac{3\pi}{5}\right)$ | |
| Ans | $\sin^{-1}\left(\sin \frac{3\pi}{5}\right) = \sin^{-1}\left(\sin\left(\pi - \frac{2\pi}{5}\right)\right)$ $= \sin^{-1}\left(\sin\left(\frac{2\pi}{5}\right)\right)$ $= \frac{2\pi}{5}$ | <p>1</p> <p>1/2</p> <p>1/2</p> |
| Q25. | The diagonals of a parallelogram are given by $\vec{a} = 2\hat{i} - \hat{j} + \hat{k}$ and $\vec{b} = \hat{i} + 3\hat{j} - \hat{k}$. Find the area of the parallelogram. | |
| Ans | $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 2 & -1 & 1 \\ 1 & 3 & -1 \end{vmatrix} = -2\hat{i} + 3\hat{j} + 7\hat{k}$ <p>Area of parallelogram $= \frac{1}{2} \vec{a} \times \vec{b}$</p> $= \frac{1}{2} \sqrt{(-2)^2 + 3^2 + 7^2} = \frac{\sqrt{62}}{2}$ | <p>1</p> <p>1</p> |
| SECTION C | | |
| This section comprises short answer (SA) type questions of 3 marks each. | | |

| | |
|-------------|--|
| Q26. | <p>(a) Verify that lines given by $\vec{r} = (1 - \lambda)\hat{i} + (\lambda - 2)\hat{j} + (3 - 2\lambda)\hat{k}$ and $\vec{r} = (\mu + 1)\hat{i} + (2\mu - 1)\hat{j} - (2\mu + 1)\hat{k}$ are skew lines. Hence, find shortest distance between the lines.</p> <p style="text-align: center;">OR</p> <p>(b) During a cricket match, the position of the bowler, the wicket keeper and the leg slip fielder are in a line given by $\vec{B} = 2\hat{i} + 8\hat{j}$, $\vec{W} = 6\hat{i} + 12\hat{j}$ and $\vec{F} = 12\hat{i} + 18\hat{j}$ respectively. Calculate the ratio in which the wicketkeeper divides the line segment joining the bowler and the leg slip fielder.</p> |
|-------------|--|


| | | |
|---------------|--|---|
| Ans(a) | <p>Rewriting the lines, we get</p> $\vec{r} = (\hat{i} - 2\hat{j} + 3\hat{k}) + \lambda(-\hat{i} + \hat{j} - 2\hat{k}) \text{ and } \vec{r} = (\hat{i} - \hat{j} - \hat{k}) + \mu(\hat{i} + 2\hat{j} - 2\hat{k})$ <p>Let $\vec{a}_1 = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{a}_2 = \hat{i} - \hat{j} - \hat{k}$, $\vec{b}_1 = -\hat{i} + \hat{j} - 2\hat{k}$, $\vec{b}_2 = \hat{i} + 2\hat{j} - 2\hat{k}$</p> <p>Note that the dr's of given lines are not proportional so, they are not parallel lines. The lines will be skew if they do not intersect each other also.</p> <p>Here $\vec{a}_2 - \vec{a}_1 = \hat{j} - 4\hat{k}$, $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & -2 \\ 1 & 2 & -2 \end{vmatrix} = 2\hat{i} - 4\hat{j} - 3\hat{k}$</p> <p>Consider $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)$ $= (\hat{j} - 4\hat{k}) \cdot (2\hat{i} - 4\hat{j} - 3\hat{k}) = 8 \neq 0$</p> <p>Hence lines will not intersect. So the lines are skew.</p> <p>Shortest Distance = $\frac{ (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) }{ \vec{b}_1 \times \vec{b}_2 }$ $= \frac{8}{\sqrt{4 + 16 + 9}} = \frac{8}{\sqrt{29}}$</p> | <p>1/2</p> <p>1/2 + 1/2</p> <p>1/2</p> <p>1</p> |
|---------------|--|---|

OR

| | | |
|---------------|---|----------------------------|
| Ans(b) | <p>Let the wicket keeper divides the line segment in ratio $k : 1$</p> $\therefore \vec{W} = \frac{k\vec{F} + 1\vec{B}}{k + 1}$ $\Rightarrow 6\hat{i} + 12\hat{j} = \left(\frac{12k + 2}{k + 1}\right)\hat{i} + \left(\frac{18k + 8}{k + 1}\right)\hat{j}$ $\Rightarrow k = \frac{2}{3}$ <p>Hence, the required ratio is 2 : 3</p> | <p>1</p> <p>1</p> <p>1</p> |
|---------------|---|----------------------------|



Q27. Solve the following linear programming problem graphically :
 Maximise $Z = 20x + 30y$
 Subject to the constraints :
 $x + y \leq 80$
 $2x + 3y \geq 100$
 $x \geq 14$
 $y \geq 14$



For correct graph and shading $1\frac{1}{2}$

For correct table **1**

$\frac{1}{2}$

Q28. The area of an expanding rectangle is increasing at the rate of $48 \text{ cm}^2/\text{s}$. The length of the rectangle is always square of its breadth. At what rate the length of rectangle increasing at an instant, when breadth = 4.5 cm ?

Ans Let the length and breadth of the expanding rectangle at any time be 'x' and 'y' respectively.

Then, $x = y^2$, $A(\text{Area}) = xy = x^{\frac{3}{2}}$

$$\frac{dA}{dt} = \frac{3}{2} \sqrt{x} \frac{dx}{dt}$$


$$\Rightarrow 48 = \frac{3}{2} \sqrt{(4.5)^2} \frac{dx}{dt} \Rightarrow \frac{dx}{dt} = \frac{64}{9} \text{ cm/s}$$

1

1

1

| | | | | | | | | | | | | |
|-----------------------|---|---|----|---|---|---|-------------|---|----|----|---|--|
| <p>Q29.</p> | <p>(a) The probability distribution for the number of students being absent in a class on a Saturday is as follows :</p> <table border="1" data-bbox="304 232 815 333"> <tbody> <tr> <td>X</td> <td>0</td> <td>2</td> <td>4</td> <td>5</td> </tr> <tr> <td>P(X)</td> <td>p</td> <td>2p</td> <td>3p</td> <td>p</td> </tr> </tbody> </table> <p>Where X is the number of students absent.</p> <p>(i) Calculate p. 1</p> <p>(ii) Calculate the mean of the number of absent students on Saturday. 2</p> <p style="text-align: center;">OR</p> <p>(b) For the vacancy advertised in the newspaper, 3000 candidates submitted their applications. From the data it was revealed that two third of the total applicants were females and other were males. The selection for the job was done through a written test. The performance of the applicants indicates that the probability of a male getting a distinction in written test is 0.4 and that a female getting a distinction is 0.35. Find the probability that the candidate chosen at random will have a distinction in the written test.</p> | X | 0 | 2 | 4 | 5 | P(X) | p | 2p | 3p | p | |
| X | 0 | 2 | 4 | 5 | | | | | | | | |
| P(X) | p | 2p | 3p | p | | | | | | | | |
| <p>Ans (a)</p> | <p>(i) Since $\sum P(X) = 1 \Rightarrow p + 2p + 3p + p = 1$</p> $\Rightarrow p = \frac{1}{7}$ <p>(ii) Mean = $\sum X.P(X) = 0(p) + 2(2p) + 4(3p) + 5(p)$</p> $= 21p = 21\left(\frac{1}{7}\right) = 3$ | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> | | | | | | | | | | |
| OR | | | | | | | | | | | | |
| <p>Ans(b)</p> | <p>Let E_1 : The applicant is a male</p> <p>E_2 : The applicant is a female</p> <p>A : The candidate chosen will have distinction in the written test.</p> <p>$P(E_1) = \frac{1}{3}, P(E_2) = \frac{2}{3}, P(A E_1) = 0.4, P(A E_2) = 0.35$</p> <p>$\therefore P(A) = P(E_1)P(A E_1) + P(E_2)P(A E_2)$</p> $= \frac{1}{3} \times 0.4 + \frac{2}{3} \times 0.35$ $= \frac{11}{30}$ | <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> | | | | | | | | | | |


| | | |
|--|--|---|
| <p>Q30.</p> | <p>(a) Find : $\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx$</p> <p style="text-align: center;">OR</p> <p>(b) Evaluate : $\int_0^{\frac{\pi}{2}} \frac{5 \sin x + 3 \cos x}{\sin x + \cos x} dx$</p> | |
| <p>Ans (a)</p> | $\int \frac{\cos 2x}{(\sin x + \cos x)^2} dx = \int \frac{\cos^2 x - \sin^2 x}{(\sin x + \cos x)^2} dx$ $= \int \frac{\cos x - \sin x}{\sin x + \cos x} dx$ $= \log \sin x + \cos x + C$ | <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> |
| OR | | |
| <p>Ans(b)</p> | $I = \int_0^{\pi/2} \frac{5 \sin x + 3 \cos x}{\sin x + \cos x} dx \quad \text{-- (i)}$ $I = \int_0^{\pi/2} \frac{5 \sin\left(\frac{\pi}{2} - x\right) + 3 \cos\left(\frac{\pi}{2} - x\right)}{\sin\left(\frac{\pi}{2} - x\right) + \cos\left(\frac{\pi}{2} - x\right)} dx = \int_0^{\pi/2} \frac{5 \cos x + 3 \sin x}{\cos x + \sin x} dx \quad \text{-- (ii)}$ <p>Adding (i) and (ii), we get</p> $2I = \int_0^{\pi/2} 8 dx \Rightarrow I = 4x \Big _0^{\pi/2} = 2\pi$ | <p style="text-align: right;">1½</p> <p style="text-align: right;">1½</p> |
| <p>Q31.</p> | <p>Sketch the graph of $y = x + 3$ and find the area of the region enclosed by the curve, x-axis, between $x = -6$ and $x = 0$, using integration.</p> | |
| <p>A31.</p> <p>Required Area</p> $= \int_{-6}^0 y dx$ $= 2 \int_{-3}^0 (x + 3) dx$ $= 2 \left[\frac{(x + 3)^2}{2} \right]_{-3}^0$ $= 9$ |  | <p>For correct graph: 1 mark</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p> |

SECTION D

This section comprises long answer (LA) type questions of 5 marks each.

| | | |
|----------------|---|---|
| Q32. | <p>(a) Differentiate $\tan^{-1} \frac{\sqrt{1-x^2}}{x}$ w.r.t. $\cos^{-1} (2x\sqrt{1-x^2})$, $x \in \left(\frac{1}{\sqrt{2}}, 1\right)$</p> <p style="text-align: center;">OR</p> <p>(b) Find $\frac{dy}{dx}$, if $y = x^{\tan x} + \frac{\sqrt{x^2+1}}{2}$.</p> | |
| Ans (a) | <p>Put $x = \cos \theta \Rightarrow \theta = \cos^{-1} x$</p> <p>Let $u = \tan^{-1} \frac{\sqrt{1-x^2}}{x} = \tan^{-1} \left(\frac{\sin \theta}{\cos \theta} \right) = \tan^{-1} (\tan \theta) = \theta = \cos^{-1} x$</p> <p>$\Rightarrow \frac{du}{dx} = -\frac{1}{\sqrt{1-x^2}}$</p> <p>Let $v = \cos^{-1} (2x\sqrt{1-x^2}) = \cos^{-1} (\sin 2\theta) = \cos^{-1} \left(\cos \left(\frac{\pi}{2} - 2\theta \right) \right) = \frac{\pi}{2} - 2\cos^{-1} x$</p> <p>$\Rightarrow \frac{dv}{dx} = \frac{2}{\sqrt{1-x^2}}$</p> <p>$\therefore \frac{du}{dv} = \frac{du/dx}{dv/dx} = -\frac{1}{2}$</p> | <p>$\frac{1}{2}$</p> <p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |
| OR | | |
| Ans (b) | <p>Let $y = u + v \Rightarrow \frac{dy}{dx} = \frac{du}{dx} + \frac{dv}{dx}$, where $u = x^{\tan x}$, $v = \frac{\sqrt{x^2+1}}{2}$</p> <p>$u = x^{\tan x} \Rightarrow \log u = \tan x \log x$, differentiating with respect to 'x', we get</p> <p>$\Rightarrow \frac{1}{u} \frac{du}{dx} = \frac{\tan x}{x} + \sec^2 x \log x$</p> <p>$\Rightarrow \frac{du}{dx} = u \left(\frac{\tan x}{x} + \sec^2 x \log x \right) = x^{\tan x} \left(\frac{\tan x}{x} + \sec^2 x \log x \right)$</p> <p>$v = \frac{\sqrt{x^2+1}}{2} \Rightarrow \frac{dv}{dx} = \frac{2x}{4\sqrt{x^2+1}} = \frac{x}{2\sqrt{x^2+1}}$</p> <p>$\Rightarrow \frac{dy}{dx} = x^{\tan x} \left(\frac{\tan x}{x} + \sec^2 x \log x \right) + \frac{x}{2\sqrt{x^2+1}}$</p> | <p>1</p> <p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>$1\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> |


| | | |
|------|--|--|
| Q33. | Find the absolute maximum and absolute minimum of function $f(x) = 2x^3 - 15x^2 + 36x + 1$ on $[1, 5]$. | |
| A33. | $f(x) = 2x^3 - 15x^2 + 36x + 1$ $\Rightarrow f'(x) = 6(x^2 - 5x + 6) = 6(x - 2)(x - 3)$ $f'(x) = 0 \Rightarrow x = 2, 3 \in [1, 5]$ <p>Now $f(1) = 24, f(2) = 29, f(3) = 28, f(5) = 56$</p> <p>Hence, the absolute maximum value is 56 and the absolute minimum value is 24.</p> | <p>1</p> <p>1</p> <p>2</p> <p>1</p> |
| Q34. | <p>A school wants to allocate students into three clubs : Sports, Music and Drama, under following conditions :</p> <ul style="list-style-type: none"> The number of students in Sports club should be equal to the sum of the number of students in Music and Drama club. The number of students in Music club should be 20 more than half the number of students in Sports club. The total number of students to be allocated in all three clubs are 180. <p>Find the number of students allocated to different clubs, using matrix method.</p> | |
| Ans | <p>Let x, y and z be the no. of students allocated to Sports, Music and Drama clubs respectively.</p> <p>Here, $x = y + z, y = \frac{x}{2} + 20, x + y + z = 180$</p> $\Rightarrow x - y - z = 0, x - 2y = -40, x + y + z = 180$ <p>Given equations can be written as $AX = B$</p> <p>where, $A = \begin{bmatrix} 1 & -1 & -1 \\ 1 & -2 & 0 \\ 1 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$</p> <p>$A = -4 \neq 0 \Rightarrow A^{-1}$ exists.</p> $adjA = \begin{bmatrix} -2 & 0 & -2 \\ -1 & 2 & -1 \\ 3 & -2 & -1 \end{bmatrix}$ $A^{-1} = \frac{1}{ A } \times adjA = \frac{1}{-4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix}$ <p>$X = A^{-1}B$</p> $= \frac{1}{-4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix} = \begin{bmatrix} 90 \\ 65 \\ 25 \end{bmatrix}$ <p>$\therefore x = 90, y = 65, z = 25$</p> <p>Number of students allocated in sports, music and drama are 90, 65 and 25 respectively .</p> | <p>1½</p> <p>½</p> <p>½</p> <p>1</p> <p>½</p> <p>1</p> |

| | | |
|----------------------|---|---|
| <p>Q35.</p> | <p>(a) Find the image A' of the point A(1, 6, 3) in the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$. Also, find the equation of the line joining A and A'.</p> <p style="text-align: center;">OR</p> <p>(b) Find a point P on the line $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}$ such that its distance from point Q(2, 4, -1) is 7 units. Also, find the equation of line joining P and Q.</p> | |
| <p>Ans(a)</p> | <p>The equation of given line is $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda$</p> <p>Any arbitrary point on the line is $M(\lambda, 2\lambda + 1, 3\lambda + 2)$</p> <p>dir's of AM are $\langle \lambda - 1, 2\lambda - 5, 3\lambda - 1 \rangle$</p> <p>Here $1(\lambda - 1) + 2(2\lambda - 5) + 3(3\lambda - 1) = 0$</p> <p>$\Rightarrow \lambda = 1$</p> <p>$\therefore M(1, 3, 5)$ is the foot perpendicular of the point A to the given line.</p> <p>Let image of point A in the line be $A'(\alpha, \beta, \gamma)$</p> <p>Since M is the mid-point of AA', so $M\left(\frac{1+\alpha}{2}, \frac{6+\beta}{2}, \frac{3+\gamma}{2}\right) = M(1, 3, 5)$</p> <p>$\Rightarrow A'(1, 0, 7)$ is the image of A.</p> <p>Also, Equation of AA' is $\frac{x-1}{0} = \frac{y-6}{-3} = \frac{z-3}{2}$</p> | <div style="text-align: right;">  </div> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> |
| OR | | |
| <p>Ans(b)</p> | <p>The given line is $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9} = \lambda$ and $Q(2, 4, -1)$</p> <p>Any random point on the line will be given by $P(\lambda - 5, 4\lambda - 3, -9\lambda + 6)$</p> <p>Since $PQ = 7 \Rightarrow \sqrt{(\lambda - 7)^2 + (4\lambda - 7)^2 + (-9\lambda + 7)^2} = 7$</p> <p>$\Rightarrow 98(\lambda^2 - 2\lambda + 1) = 0 \Rightarrow \lambda = 1$</p> <p>Hence, the required point is $P(-4, 1, -3)$</p> <p>The equation of line PQ is $\frac{x+4}{6} = \frac{y-1}{3} = \frac{z+3}{2}$ or $\frac{x-2}{6} = \frac{y-4}{3} = \frac{z+1}{2}$</p> | <p style="text-align: right;">1</p> |

SECTION E

This section comprises 3 case study-based questions of 4 marks each.

Q36.



A bank offers loan to its customers on different types of interest namely, fixed rate, floating rate and variable rate. From the past data with the bank, it is known that a customer avails loan on fixed rate, floating rate or variable rate with probabilities 10%, 20% and 70% respectively. A customer after availing loan can pay the loan or default on loan repayment. The bank data suggests that the probability that a person defaults on loan after availing it at fixed rate, floating rate and variable rate is 5%, 3% and 1% respectively.

Based on the above information, answer the following :

- (i) What is the probability that a customer after availing the loan will default on the loan repayment ? 2
- (ii) A customer after availing the loan, defaults on loan repayment. What is the probability that he availed the loan at a variable rate of interest ? 2

Ans

E_1 :customer avails loan on fixed rate

E_2 :customer avails loan on floating rate

E_3 :customer avails loan on variable rate

A:the person defaults on the loan

$$P(E_1) = \frac{1}{10}, P(E_2) = \frac{2}{10}, P(E_3) = \frac{7}{10}$$

$$P(A|E_1) = \frac{5}{100}, P(A|E_2) = \frac{3}{100}, P(A|E_3) = \frac{1}{100}$$

$$\begin{aligned} (i) P(A) &= P(E_1) \cdot P(A|E_1) + P(E_2) \cdot P(A|E_2) + P(E_3) \cdot P(A|E_3) \\ &= \frac{1}{10} \times \frac{5}{100} + \frac{2}{10} \times \frac{3}{100} + \frac{7}{10} \times \frac{1}{100} \\ &= \frac{18}{1000} \text{ or } \frac{9}{500} \end{aligned}$$

$$\begin{aligned} (ii) P(E_3|A) &= \frac{P(E_3) \cdot P(A|E_3)}{P(E_1) \cdot P(A|E_1) + P(E_2) \cdot P(A|E_2) + P(E_3) \cdot P(A|E_3)} \\ &= \frac{\frac{7}{10} \times \frac{1}{100}}{\frac{18}{1000}} \\ &= \frac{7}{18} \end{aligned}$$


1

1

1

1

Q37.



A technical company is designing a rectangular solar panel installation on a roof using 300 metres of boundary material. The design includes a partition running parallel to one of the sides dividing the area (roof) into two sections.

Let the length of the side perpendicular to the partition be x metres and with parallel to the partition be y metres.

Based on this information, answer the following questions :

- (i) Write the equation for the total boundary material used in the boundary and parallel to the partition in terms of x and y . 1
- (ii) Write the area of the solar panel as a function of x . 1
- (iii) (a) Find the critical points of the area function. Use second derivative test to determine critical points at the maximum area. Also, find the maximum area. 2
- OR**
- (iii) (b) Using first derivative test, calculate the maximum area the company can enclose with the 300 metres of boundary material, considering the parallel partition. 2

Ans

(i) $2x + 3y = 300$

(ii) $A = xy = \frac{x}{3}(300 - 2x)$

(iii)(a) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$

$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$

For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$

Also, $\frac{d^2A}{dx^2} = -\frac{4}{3} < 0$. So, A is maximum at $x = 75$

Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$

OR

(iii)(b) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$

$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$

For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$

As $\frac{dA}{dx}$ changes its sign from positive to negative as x passes through

$x = 75$ from left to right, which means $x = 75$ is the point of maximum.

Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$

1

1

1/2

1/2

1/2

1/2

1/2

1/2

1/2

1/2

| | | |
|--------------------|--|---|
| | <p>Note : Full credit to be given if the student takes equation as $2x + 2y = 300$ or $2x + 4y = 300$ or $4x + 4y = 300$ or $4x + 3y = 300$ The solutions of sub-parts will differ and marks may be given accordingly.</p> | |
| <p>Q38.</p> | <p>A class-room teacher is keen to assess the learning of her students the concept of “relations” taught to them. She writes the following five relations each defined on the set $A = \{1, 2, 3\}$:</p> <p>$R_1 = \{(2, 3), (3, 2)\}$</p> <p>$R_2 = \{(1, 2), (1, 3), (3, 2)\}$</p> <p>$R_3 = \{(1, 2), (2, 1), (1, 1)\}$</p> <p>$R_4 = \{(1, 1), (1, 2), (3, 3), (2, 2)\}$</p> <p>$R_5 = \{(1, 1), (1, 2), (3, 3), (2, 2), (2, 1), (2, 3), (3, 2)\}$</p> <p>The students are asked to answer the following questions about the above relations :</p> <p>(i) Identify the relation which is reflexive, transitive but not symmetric.</p> <p>(ii) Identify the relation which is reflexive and symmetric but not transitive.</p> <p>(iii) (a) Identify the relations which are symmetric but neither reflexive nor transitive.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) What pairs should be added to the relation R_2 to make it an equivalence relation ?</p> | |
| <p>Ans</p> | <p>(i) R_4</p> <p>(ii) R_5</p> <p>(iii) (a) R_1 and R_3</p> <p style="text-align: center;">OR</p> <p>(iii) (b) Required pairs to be added to make the relation R_2 as an equivalence relation are: $(1,1), (2,2), (3,3), (2,1), (3,1)$ and $(2,3)$</p> | <p>1</p> <p>1</p> <p>1+1</p> <p>2</p> |

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025
SUBJECT NAME MATHEMATICS (Q.P. CODE – 65/1/3)

General Instructions: -


| | |
|----------|--|
| 1 | You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully. |
| 2 | “Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.” |
| 3 | Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded. |
| 4 | The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly. |
| 5 | The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators. |
| 6 | Evaluators will mark (√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing. |
| 7 | If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly. |
| 8 | If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly. |
| 9 | If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” . |

| | |
|----|---|
| 10 | No marks to be deducted for the cumulative effect of an error. It should be penalized only once. |
| 11 | A full scale of marks _____ (example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it. |
| 12 | Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper. |
| 13 | <p>Ensure that you do not make the following common types of errors committed by the Examiner in the past: -</p> <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks |
| 14 | While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks. |
| 15 | Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously. |
| 16 | The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation. |
| 17 | Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words. |
| 18 | The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme. |

| | | |
|------------|--|-----------------|
| Q4. | <p>If $\int \frac{2^x}{x^2} dx = k \cdot 2^{\frac{1}{x}} + C$, then k is equal to</p> <p>(A) $\frac{-1}{\log 2}$ (B) $-\log 2$</p> <p>(C) -1 (D) $\frac{1}{2}$</p> | |
| Ans | <p>(A) $\frac{-1}{\log 2}$</p> | <p>1</p> |
| Q5. | <p>If $A = \begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$, then A^{-1} is</p> <p>(A) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$ (B) $\begin{bmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & -1 \end{bmatrix}$</p> <p>(C) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$ (D) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$</p> | |
| Ans | <p>(D) $\begin{bmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$</p> | <p>1</p> |
| Q6. | <p>If $\begin{bmatrix} x+y & 3y \\ 3x & x+3 \end{bmatrix} = \begin{bmatrix} 9 & 4x+y \\ x+6 & y \end{bmatrix}$, then $(x-y) = ?$</p> <p>(A) -7 (B) -3</p> <p>(C) 3 (D) 7</p> | |
| A6. | <p>(B) -3</p> | <p>1</p> |
| Q7. | <p>Let M and N be two events such that $P(M) = 0.6$, $P(N) = 0.2$ and $P(M \cap N) = 0.5$, then $P(M'/N')$ is</p> <p>(A) $\frac{7}{8}$ (B) $\frac{2}{5}$</p> <p>(C) $\frac{1}{2}$ (D) $\frac{2}{3}$</p> | |
| Ans | <p>(A) $\frac{7}{8}$</p> | <p>1</p> |

| | | |
|-------------|---|----------|
| Q8. | Which of the following is <u>not</u> a homogeneous function of x and y ? (A) $y^2 - xy$ (B) $x - 3y$ (C) $\sin^2 \frac{y}{x} + \frac{y}{x}$ (D) $\tan x - \sec y$ | |
| Ans | (D) $\tan x - \sec y$ | 1 |
| Q9. | If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$, $ \vec{a} = \sqrt{37}$, $ \vec{b} = 3$ and $ \vec{c} = 4$, then angle between \vec{b} and \vec{c} is (A) $\frac{\pi}{6}$ (B) $\frac{\pi}{4}$ (C) $\frac{\pi}{3}$ (D) $\frac{\pi}{2}$ | |
| Ans | (C) $\frac{\pi}{3}$ | 1 |
| Q10. | If $f(x) = x + x - 1 $, then which of the following is correct ? (A) $f(x)$ is both continuous and differentiable, at $x = 0$ and $x = 1$. (B) $f(x)$ is differentiable but not continuous, at $x = 0$ and $x = 1$. (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. (D) $f(x)$ is neither continuous nor differentiable, at $x = 0$ and $x = 1$. | |
| Ans | (C) $f(x)$ is continuous but not differentiable, at $x = 0$ and $x = 1$. | 1 |
| Q11. | A system of linear equations is represented as $AX = B$, where A is coefficient matrix, X is variable matrix and B is the constant matrix. Then the system of equations is (A) Consistent, if $ A \neq 0$, solution is given by $X = BA^{-1}$. (B) Inconsistent if $ A = 0$ and $(\text{adj } A) B = 0$ (C) Inconsistent if $ A \neq 0$ (D) May or may not be consistent if $ A = 0$ and $(\text{adj } A) B = 0$ | |
| Ans | (D) May or may not be consistent if $ A = 0$ and $(\text{adj } A) B = 0$ | 1 |
| Q12. | The absolute maximum value of function $f(x) = x^3 - 3x + 2$ in $[0, 2]$ is : (A) 0 (B) 2 (C) 4 (D) 5 | |
| Ans | (C) 4 | 1 |

| | | |
|--------------------|---|-----------------|
| <p>Q13.</p> | <p>The order and degree of differential function $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^5 = \frac{d^2y}{dx^2}$ are</p> <p>(A) order 1, degree 1 (B) order 1, degree 2 (C) order 2, degree 1 (D) order 2, degree 2</p> | |
| <p>Ans</p> | <p>(C) order 2, degree 1</p> | <p>1</p> |
| <p>Q14.</p> | <p>The graph of a trigonometric function is as shown. Which of the following will represent graph of its inverse ?</p> <p>(A) </p> <p>(B) </p> <p>(C) </p> <p>(D) </p> | |
| <p>Ans</p> | <p>(C) </p> | <p>1</p> |
| <p>Q15.</p> | <p>The corner points of the feasible region in graphical representation of a L.P.P. are (2, 72), (15, 20) and (40, 15). If $Z = 18x + 9y$ be the objective function, then</p> <p>(A) Z is maximum at (2, 72), minimum at (15, 20) (B) Z is maximum at (15, 20) minimum at (40, 15) (C) Z is maximum at (40, 15), minimum at (15, 20) (D) Z is maximum at (40, 15), minimum at (2, 72)</p> | |
| <p>Ans</p> | <p>(C) Z is maximum at (40, 15), minimum at (15, 20)</p> | <p>1</p> |

| | | |
|-------------|---|----------|
| Q16. | <p>Let $A = \begin{bmatrix} 1 & -2 & -1 \\ 0 & 4 & -1 \\ -3 & 2 & 1 \end{bmatrix}$, $B = \begin{bmatrix} -2 \\ -5 \\ -7 \end{bmatrix}$, $C = [9 \ 8 \ 7]$, which of the following is defined ?</p> <p>(A) Only AB (B) Only AC (C) Only BA (D) All AB, AC and BA</p> | |
| Ans | (A) Only AB | 1 |
| Q17. | <p>If A and B are invertible matrices, then which of the following is <u>not</u> correct ?</p> <p>(A) $(A + B)^{-1} = B^{-1} + A^{-1}$ (B) $(AB)^{-1} = B^{-1}A^{-1}$ (C) $\text{adj}(A) = A A^{-1}$ (D) $A ^{-1} = A^{-1}$</p> | |
| Ans | (A) $(A + B)^{-1} = B^{-1} + A^{-1}$ | 1 |
| Q18. | <p>The area of the shaded region bounded by the curves $y^2 = x$, $x = 4$ and the x-axis is given by</p>  <p>(A) $\int_0^4 x \, dx$ (B) $\int_0^2 y^2 \, dy$ (C) $2 \int_0^4 \sqrt{x} \, dx$ (D) $\int_0^4 \sqrt{x} \, dx$</p> | |
| Ans | (D) $\int_0^4 \sqrt{x} \, dx$ | 1 |

Assertion – Reason Based Questions


Direction : Question numbers 19 and 20 are Assertion (A) and Reason (R) based questions carrying 1 mark each. Two statements are given, one labelled Assertion (A) and other labelled Reason (R). Select the correct answer from the options (A), (B), (C) and (D) as given below.

- (A) Both Assertion (A) and Reason (R) are true and the Reason (R) is the correct explanation of the Assertion (A).
 (B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).
 (C) Assertion (A) is true, but Reason (R) is false.
 (D) Assertion (A) is false, but Reason (R) is true.

| | | |
|----------------------|---|---|
| <p>Q23.</p> | <p>(a) Two friends while flying kites from different locations, find the strings of their kites crossing each other. The strings can be represented by vectors $\vec{a} = 3\hat{i} + \hat{j} + 2\hat{k}$ and $\vec{b} = 2\hat{i} - 2\hat{j} + 4\hat{k}$. Determine the angle formed between the kite strings. Assume there is no slack in the strings.</p> <p style="text-align: center;">OR</p> <p>(b) Find a vector of magnitude 21 units in the direction opposite to that of \vec{AB} where A and B are the points A(2, 1, 3) and B(8, -1, 0) respectively.</p> | |
| <p>Ans(a)</p> | <p>Let the required angle between the kite strings be θ.</p> <p>Then, $\cos\theta = \frac{\vec{a} \cdot \vec{b}}{ \vec{a} \vec{b} }$</p> $\Rightarrow \cos\theta = \frac{(3\hat{i} + \hat{j} + 2\hat{k})(2\hat{i} - 2\hat{j} + 4\hat{k})}{\sqrt{9+1+4}\sqrt{4+4+16}} = \frac{12}{\sqrt{336}} = \frac{3}{\sqrt{21}}$ $\Rightarrow \theta = \cos^{-1}\left(\frac{12}{\sqrt{336}}\right) \text{ or } \cos^{-1}\left(\frac{3}{\sqrt{21}}\right)$ | <p style="text-align: center;">1½</p> <p style="text-align: center;">½</p> |
| OR | | |
| <p>Ans(b)</p> | <p>$\vec{BA} = -6\hat{i} + 2\hat{j} + 3\hat{k}$</p> <p>Required unit vector of magnitude 21</p> $= 21 \times \left(\frac{-6\hat{i} + 2\hat{j} + 3\hat{k}}{\sqrt{36+4+9}} \right)$ $= 3(-6\hat{i} + 2\hat{j} + 3\hat{k}) \text{ or } -18\hat{i} + 6\hat{j} + 9\hat{k}$ | <p style="text-align: center;">1</p> <p style="text-align: center;">½</p> <p style="text-align: center;">½</p> |
| <p>Q24.</p> | <p>Solve for x,</p> $2 \tan^{-1} x + \sin^{-1} \left(\frac{2x}{1+x^2} \right) = 4\sqrt{3}$ | |
| <p>Ans</p> | $2 \tan^{-1} x + \sin^{-1} \left(\frac{2x}{1+x^2} \right) = 4\sqrt{3} \Rightarrow 2 \tan^{-1} x + 2 \tan^{-1} x = 4\sqrt{3}$ $\Rightarrow \tan^{-1} x = \sqrt{3} \notin \left(-\frac{\pi}{2}, \frac{\pi}{2} \right)$ <p>$\therefore x = \tan(\sqrt{3})$ which has no solution.</p> | <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> |

| | | |
|---|--|---|
| Q25. | <p>(a) Differentiate $2^{\cos^2 x}$ w.r.t $\cos^2 x$.</p> <p style="text-align: center;">OR</p> <p>(b) If $\tan^{-1}(x^2 + y^2) = a^2$, then find $\frac{dy}{dx}$.</p> | |
| Ans(a) | <p>Let $u = 2^{\cos^2 x} \Rightarrow \frac{du}{dx} = 2^{\cos^2 x} (-2 \cos x \sin x) \log 2$</p> <p>Let $v = \cos^2 x \Rightarrow \frac{dv}{dx} = -2 \cos x \sin x$</p> <p>Now $\frac{du}{dv} = \frac{\left(\frac{du}{dx}\right)}{\left(\frac{dv}{dx}\right)} = 2^{\cos^2 x} \log 2$</p> | <p style="text-align: center;">1</p> <p style="text-align: center;">1/2</p> <p style="text-align: center;">1/2</p> |
| OR | | |
| Ans(b) | <p>$\tan^{-1}(x^2 + y^2) = a^2 \Rightarrow x^2 + y^2 = \tan a^2$</p> <p>Differentiate both sides wrt x,</p> <p>$2x + 2y \frac{dy}{dx} = 0$</p> <p>$\Rightarrow \frac{dy}{dx} = -\frac{x}{y}$</p> | <p style="text-align: center;">1/2</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1/2</p> |
| SECTION C | | |
| This section comprises short answer (SA) type questions of 3 marks each. | | |
| Q26. | <p>Solve the following linear programming problem graphically :</p> <p>Maximize $Z = 8x + 9y$</p> <p>Subject to the constraints :</p> <p style="margin-left: 40px;">$2x + 3y \leq 6$</p> <p style="margin-left: 40px;">$3x - 2y \leq 6$</p> <p style="margin-left: 40px;">$y \leq 1$</p> <p style="margin-left: 40px;">$x \geq 0, y \geq 0$</p> | |

Ans



| Corner Point | Value of $Z = 8x + 9y$ |
|---|------------------------|
| $A(0,1)$ | 9 |
| $B\left(\frac{3}{2}, 1\right)$ | 21 |
| $C\left(\frac{30}{13}, \frac{6}{13}\right)$ | $\frac{294}{13}$ |
| $D(2,0)$ | 16 |
| $O(0,0)$ | 0 |

$$\text{Max}(Z) = \frac{294}{13}$$

For correct graph and shading
1½

For correct table
1

½

Q27.

(a) Find : $\int \frac{2x-1}{(x-1)(x+2)(x-3)} dx$

OR

(b) Evaluate : $\int_0^5 (|x-1| + |x-2| + |x-5|) dx$

Ans(a)

$$\int \frac{2x-1}{(x-1)(x+2)(x-3)} dx = -\frac{1}{6} \int \frac{1}{x-1} dx - \frac{1}{3} \int \frac{1}{x+2} dx + \frac{1}{2} \int \frac{1}{x-3} dx$$


(Using Partial Fraction)

$$= -\frac{1}{6} \log|x-1| - \frac{1}{3} \log|x+2| + \frac{1}{2} \log|x-3| + C$$

1½

1½

OR

| | | |
|-----------------------|--|---|
| <p>Q30.</p> | <p>(a) Verify that lines given by $\vec{r} = (1 - \lambda)\hat{i} + (\lambda - 2)\hat{j} + (3 - 2\lambda)\hat{k}$ and $\vec{r} = (\mu + 1)\hat{i} + (2\mu - 1)\hat{j} - (2\mu + 1)\hat{k}$ are skew lines. Hence, find shortest distance between the lines.</p> <p style="text-align: center;">OR</p> <p>(b) During a cricket match, the position of the bowler, the wicket keeper and the leg slip fielder are in a line given by $\vec{B} = 2\hat{i} + 8\hat{j}$, $\vec{W} = 6\hat{i} + 12\hat{j}$ and $\vec{F} = 12\hat{i} + 18\hat{j}$ respectively. Calculate the ratio in which the wicketkeeper divides the line segment joining the bowler and the leg slip fielder.</p> | |
| <p>Ans (a)</p> | <p>Rewriting the lines, we get</p> $\vec{r} = (\hat{i} - 2\hat{j} + 3\hat{k}) + \lambda(-\hat{i} + \hat{j} - 2\hat{k}) \text{ and } \vec{r} = (\hat{i} - \hat{j} - \hat{k}) + \mu(\hat{i} + 2\hat{j} - 2\hat{k})$ <p>Let $\vec{a}_1 = \hat{i} - 2\hat{j} + 3\hat{k}$, $\vec{a}_2 = \hat{i} - \hat{j} - \hat{k}$, $\vec{b}_1 = -\hat{i} + \hat{j} - 2\hat{k}$, $\vec{b}_2 = \hat{i} + 2\hat{j} - 2\hat{k}$</p> <p>Note that the dr's of given lines are not proportional so, they are not parallel lines. The lines will be skew if they do not intersect each other also.</p> <p>Here $\vec{a}_2 - \vec{a}_1 = \hat{j} - 4\hat{k}$, $\vec{b}_1 \times \vec{b}_2 = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ -1 & 1 & -2 \\ 1 & 2 & -2 \end{vmatrix} = 2\hat{i} - 4\hat{j} - 3\hat{k}$</p> <p>Consider $(\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2)$</p> $= (\hat{j} - 4\hat{k}) \cdot (2\hat{i} - 4\hat{j} - 3\hat{k}) = 8 \neq 0$ <p>Hence lines will not intersect. So the lines are skew.</p> <p>Shortest Distance = $\frac{ (\vec{a}_2 - \vec{a}_1) \cdot (\vec{b}_1 \times \vec{b}_2) }{ \vec{b}_1 \times \vec{b}_2 }$</p> $= \frac{8}{\sqrt{4 + 16 + 9}} = \frac{8}{\sqrt{29}}$ | <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2 + 1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> |
| OR | | |
| <p>Ans(b)</p> | <p>Let the wicket keeper divides the line segment in ratio $k : 1$</p> $\therefore \vec{W} = \frac{k\vec{F} + 1\vec{B}}{k + 1}$ $\Rightarrow 6\hat{i} + 12\hat{j} = \left(\frac{12k + 2}{k + 1}\right)\hat{i} + \left(\frac{18k + 8}{k + 1}\right)\hat{j}$ $\Rightarrow k = \frac{2}{3}$ <p>Hence, the required ratio is 2 : 3</p> <div style="text-align: center;">  </div> | <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> |

| | | | | | | | | | | | | |
|----------------------|--|---|----|---|---|---|-------------|---|----|----|---|--|
| <p>Q31.</p> | <p>(a) The probability distribution for the number of students being absent in a class on a Saturday is as follows :</p> <table border="1" data-bbox="300 226 802 331"> <tr> <td>X</td> <td>0</td> <td>2</td> <td>4</td> <td>5</td> </tr> <tr> <td>P(X)</td> <td>p</td> <td>2p</td> <td>3p</td> <td>p</td> </tr> </table> <p>Where X is the number of students absent.</p> <p>(i) Calculate p. 1</p> <p>(ii) Calculate the mean of the number of absent students on Saturday. 2</p> <p style="text-align: center;">OR</p> <p>(b) For the vacancy advertised in the newspaper, 3000 candidates submitted their applications. From the data it was revealed that two third of the total applicants were females and other were males. The selection for the job was done through a written test. The performance of the applicants indicates that the probability of a male getting a distinction in written test is 0.4 and that a female getting a distinction is 0.35. Find the probability that the candidate chosen at random will have a distinction in the written test.</p> | X | 0 | 2 | 4 | 5 | P(X) | p | 2p | 3p | p | |
| X | 0 | 2 | 4 | 5 | | | | | | | | |
| P(X) | p | 2p | 3p | p | | | | | | | | |
| <p>Ans(a)</p> | <p>(i) Since $\sum P(X) = 1 \Rightarrow p + 2p + 3p + p = 1$ $\Rightarrow p = \frac{1}{7}$</p> <p>(ii) Mean = $\sum X.P(X) = 0(p) + 2(2p) + 4(3p) + 5(p)$ $= 21p = 21\left(\frac{1}{7}\right) = 3$</p> | <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> | | | | | | | | | | |
| OR | | | | | | | | | | | | |
| <p>Ans(b)</p> | <p>Let E_1 : The applicant is a male E_2 : The applicant is a female A : The candidate chosen will have distinction in the written test.</p> <p>$P(E_1) = \frac{1}{3}, P(E_2) = \frac{2}{3}, P(A E_1) = 0.4, P(A E_2) = 0.35$</p> <p>$\therefore P(A) = P(E_1)P(A E_1) + P(E_2)P(A E_2)$ $= \frac{1}{3} \times 0.4 + \frac{2}{3} \times 0.35$ $= \frac{11}{30}$</p> | <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> | | | | | | | | | | |


SECTION D

This section comprises long answer (LA) type questions of 5 marks each.

| | | |
|-------------|---|--|
| Q32. | <p>A school wants to allocate students into three clubs : Sports, Music and Drama, under following conditions :</p> <ul style="list-style-type: none"> • The number of students in Sports club should be equal to the sum of the number of students in Music and Drama club. • The number of students in Music club should be 20 more than half the number of students in Sports club. • The total number of students to be allocated in all three clubs are 180. <p>Find the number of students allocated to different clubs, using matrix method.</p> | |
| Ans | <p>Let x, y and z be the no. of students allocated to Sports, Music and Drama clubs respectively.</p> <p>Here, $x = y + z, y = \frac{x}{2} + 20, x + y + z = 180$</p> <p>$\Rightarrow x - y - z = 0, x - 2y = -40, x + y + z = 180$</p> <p>Given equations can be written as $AX = B$</p> <p>where, $A = \begin{bmatrix} 1 & -1 & -1 \\ 1 & -2 & 0 \\ 1 & 1 & 1 \end{bmatrix}, B = \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix}, X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$</p> <p>$A = -4 \neq 0 \Rightarrow A^{-1}$ exists.</p> <p>$adjA = \begin{bmatrix} -2 & 0 & -2 \\ -1 & 2 & -1 \\ 3 & -2 & -1 \end{bmatrix}$</p> <p>$A^{-1} = \frac{1}{ A } \times adjA = \frac{1}{-4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix}$</p> <p>$X = A^{-1}B$</p> <p>$= \frac{1}{-4} \begin{bmatrix} 2 & 0 & 2 \\ 1 & -2 & 1 \\ -3 & 2 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ -40 \\ 180 \end{bmatrix} = \begin{bmatrix} 90 \\ 65 \\ 25 \end{bmatrix}$</p> <p>$\therefore x = 90, y = 65, z = 25$</p> <p>Number of students allocated in sports, music and drama are 90 , 65 and 25 respectively .</p> | <p>1½</p> <p>½</p> <p>½</p> <p>1</p> <p>½</p> <p>1</p> |

| | | |
|-----------|--|---|
| Q33. | Find : $\int \sin^{-1} \sqrt{\frac{x}{a+x}} dx.$ | |
| A33. | <p>Put $x = a \tan^2 \theta, dx = 2a \tan \theta \sec^2 \theta d\theta$</p> $\int \sin^{-1} \sqrt{\frac{x}{a+x}} dx = 2a \int \sin^{-1} \sqrt{\frac{\tan^2 \theta}{1+\tan^2 \theta}} (\tan \theta \cdot \sec^2 \theta) d\theta$ $= 2a \int \theta \cdot (\tan \theta \sec^2 \theta) d\theta$ $= 2a \left[\theta \frac{(\tan \theta)^2}{2} - \frac{1}{2} \int (\tan \theta)^2 d\theta \right]$ $= a [\theta \tan^2 \theta - (\tan \theta - \theta)] + C$ $= a \left[\tan^{-1} \left(\sqrt{\frac{x}{a}} \right) \cdot \frac{x}{a} - \sqrt{\frac{x}{a}} + \tan^{-1} \left(\sqrt{\frac{x}{a}} \right) \right] + C$ <p>or $(a+x) \tan^{-1} \left(\sqrt{\frac{x}{a}} \right) - \sqrt{ax} + C$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>½</p> <p>½</p> |
| Q34. | <p>(a) If $\sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$, then prove that $\frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$.</p> <p style="text-align: center;">OR</p> <p>(b) If $x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$ and $y = \sin \theta$, then find $\frac{d^2y}{dx^2}$ at $\theta = \frac{\pi}{4}$.</p> | |
| Ans(a) | <p>Let $x = \sin A, y = \sin B \Rightarrow A = \sin^{-1} x, B = \sin^{-1} y$</p> $\therefore \sqrt{1-x^2} + \sqrt{1-y^2} = a(x-y)$ $\Rightarrow \cos A + \cos B = a(\sin A - \sin B)$ $\Rightarrow 2 \cos \left(\frac{A+B}{2} \right) \cos \left(\frac{A-B}{2} \right) = 2a \cos \left(\frac{A+B}{2} \right) \sin \left(\frac{A-B}{2} \right)$ $\Rightarrow \cot \left(\frac{A-B}{2} \right) = a \Rightarrow A - B = 2 \cot^{-1} a$ $\Rightarrow \sin^{-1} x - \sin^{-1} y = 2 \cot^{-1} a$ <p>differentiate both sides wrt x,</p> $\frac{1}{\sqrt{1-x^2}} - \frac{1}{\sqrt{1-y^2}} \frac{dy}{dx} = 0$ $\Rightarrow \frac{dy}{dx} = \sqrt{\frac{1-y^2}{1-x^2}}$ | <p>1</p> <p>1</p> <p>1</p> <p>½</p> <p>1½</p> |
| OR | | |

| | | |
|----------------------|--|--|
| <p>Ans(b)</p> | $x = a \left(\cos \theta + \log \tan \frac{\theta}{2} \right)$ $\Rightarrow \frac{dx}{d\theta} = a \left(-\sin \theta + \frac{1}{\tan \frac{\theta}{2}} \times \sec^2 \frac{\theta}{2} \times \frac{1}{2} \right)$ $= a \left(-\sin \theta + \frac{1}{\sin \theta} \right) = a \left(\frac{1 - \sin^2 \theta}{\sin \theta} \right)$ $\frac{dx}{d\theta} = a \cot \theta \cos \theta$ <p>Also, $y = \sin \theta \Rightarrow \frac{dy}{d\theta} = \cos \theta$</p> $\therefore \frac{dy}{dx} = \frac{\tan \theta}{a}$ <p>Differentiating wrt x,</p> $\frac{d^2 y}{dx^2} = \frac{\sec^2 \theta}{a} \times \frac{d\theta}{dx}$ $= \frac{\sec^3 \theta \tan \theta}{a^2}$ $\left. \frac{d^2 y}{dx^2} \right]_{\text{at } \theta = \frac{\pi}{4}} = \frac{2\sqrt{2}}{a^2}$ | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> |
| <p>Q35.</p> | <p>(a) Find the image A' of the point A(1, 6, 3) in the line $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3}$. Also, find the equation of the line joining A and A'.</p> <p style="text-align: center;">OR</p> <p>(b) Find a point P on the line $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9}$ such that its distance from point Q(2, 4, -1) is 7 units. Also, find the equation of line joining P and Q.</p> | |

| | | |
|----------------------|---|--|
| <p>Ans(a)</p> | <p>The equation of given line is $\frac{x}{1} = \frac{y-1}{2} = \frac{z-2}{3} = \lambda$</p> <p>Any arbitrary point on the line is $M(\lambda, 2\lambda+1, 3\lambda+2)$</p> <p>dr's of AM are $\langle \lambda-1, 2\lambda-5, 3\lambda-1 \rangle$</p> <p>Here $1(\lambda-1) + 2(2\lambda-5) + 3(3\lambda-1) = 0$</p> <p>$\Rightarrow \lambda = 1$</p> <p>$\therefore M(1, 3, 5)$ is the foot perpendicular of the point A to the given line.</p> <p>Let image of point A in the line be $A'(\alpha, \beta, \gamma)$</p> <p>Since M is the mid-point of AA', so $M\left(\frac{1+\alpha}{2}, \frac{6+\beta}{2}, \frac{3+\gamma}{2}\right) = M(1, 3, 5)$</p> <p>$\Rightarrow A'(1, 0, 7)$ is the image of A.</p> <p>Also, Equation of AA' is $\frac{x-1}{0} = \frac{y-6}{-3} = \frac{z-3}{2}$</p> |  <p>1</p> <p>1</p> <p>1/2</p> <p>1</p> <p>1</p> |
| <p>OR</p> | | |
| <p>Ans(b)</p> | <p>The given line is $\frac{x+5}{1} = \frac{y+3}{4} = \frac{z-6}{-9} = \lambda$ and $Q(2, 4, -1)$</p> <p>Any random point on the line will be given by $P(\lambda-5, 4\lambda-3, -9\lambda+6)$</p> <p>Since $PQ = 7 \Rightarrow \sqrt{(\lambda-7)^2 + (4\lambda-7)^2 + (-9\lambda+7)^2} = 7$</p> <p>$\Rightarrow 98(\lambda^2 - 2\lambda + 1) = 0 \Rightarrow \lambda = 1$</p> <p>Hence, the required point is $P(-4, 1, -3)$</p> <p>The equation of line PQ is $\frac{x+4}{6} = \frac{y-1}{3} = \frac{z+3}{2}$ or $\frac{x-2}{6} = \frac{y-4}{3} = \frac{z+1}{2}$</p> | <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |

SECTION E

This section comprises 3 case study-based questions of 4 marks each.

Q36. A class-room teacher is keen to assess the learning of her students the concept of “relations” taught to them. She writes the following five relations each defined on the set $A = \{1, 2, 3\}$:

$R_1 = \{(2, 3), (3, 2)\}$
 $R_2 = \{(1, 2), (1, 3), (3, 2)\}$
 $R_3 = \{(1, 2), (2, 1), (1, 1)\}$
 $R_4 = \{(1, 1), (1, 2), (3, 3), (2, 2)\}$
 $R_5 = \{(1, 1), (1, 2), (3, 3), (2, 2), (2, 1), (2, 3), (3, 2)\}$

The students are asked to answer the following questions about the above relations :

(i) Identify the relation which is reflexive, transitive but not symmetric.
 (ii) Identify the relation which is reflexive and symmetric but not transitive.
 (iii) (a) Identify the relations which are symmetric but neither reflexive nor transitive.

OR

(iii) (b) What pairs should be added to the relation R_2 to make it an equivalence relation ?


| | | |
|------------|---|---|
| Ans | <p>(i) R_4 (ii) R_5 (iii)(a) R_1 and R_3</p> <p style="text-align: center;">OR</p> <p>(iii)(b) Required pairs to be added to make the relation R_2 as an equivalence relation are: $(1,1), (2,2), (3,3), (2,1), (3,1)$ and $(2,3)$</p> | <p>1 1 1+1 2</p> |
|------------|---|---|

Q37.



| | | |
|--|---|--|
| | <p>A bank offers loan to its customers on different types of interest namely, fixed rate, floating rate and variable rate. From the past data with the bank, it is known that a customer avails loan on fixed rate, floating rate or variable rate with probabilities 10%, 20% and 70% respectively. A customer after availing loan can pay the loan or default on loan repayment. The bank data suggests that the probability that a person defaults on loan after availing it at fixed rate, floating rate and variable rate is 5%, 3% and 1% respectively.</p> <p>Based on the above information, answer the following :</p> <p>(i) What is the probability that a customer after availing the loan will default on the loan repayment ? 2</p> <p>(ii) A customer after availing the loan, defaults on loan repayment. What is the probability that he availed the loan at a variable rate of interest ? 2</p> | |
|--|---|--|

| | | |
|-------------------|--|---|
| <p>Ans</p> | <p>E_1 :customer avails loan on fixed rate E_2 :customer avails loan on floating rate E_3 :customer avails loan on variable rate A:the person defaults on the loan</p> <p>$P(E_1) = \frac{1}{10}, P(E_2) = \frac{2}{10}, P(E_3) = \frac{7}{10}$</p> <p>$P(A E_1) = \frac{5}{100}, P(A E_2) = \frac{3}{100}, P(A E_3) = \frac{1}{100}$</p> <p>(i) $P(A) = P(E_1) \cdot P(A E_1) + P(E_2) \cdot P(A E_2) + P(E_3) \cdot P(A E_3)$</p> $= \frac{1}{10} \times \frac{5}{100} + \frac{2}{10} \times \frac{3}{100} + \frac{7}{10} \times \frac{1}{100}$ $= \frac{18}{1000} \text{ or } \frac{9}{500}$ <p>(ii) $P(E_3 A) = \frac{P(E_3) \cdot P(A E_3)}{P(E_1) \cdot P(A E_1) + P(E_2) \cdot P(A E_2) + P(E_3) \cdot P(A E_3)}$</p> $= \frac{\frac{7}{10} \times \frac{1}{100}}{\frac{18}{1000}}$ $= \frac{7}{18}$ | <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> |
|-------------------|--|---|

| | | |
|--------------------|--|--|
| <p>Q38.</p> |  <p>A technical company is designing a rectangular solar panel installation on a roof using 300 metres of boundary material. The design includes a partition running parallel to one of the sides dividing the area (roof) into two sections.</p> | |
|--------------------|--|--|

| | | |
|-------------------|--|--|
| | <p>Let the length of the side perpendicular to the partition be x metres and with parallel to the partition be y metres.</p> <p>Based on this information, answer the following questions :</p> <p>(i) Write the equation for the total boundary material used in the boundary and parallel to the partition in terms of x and y. 1</p> <p>(ii) Write the area of the solar panel as a function of x. 1</p> <p>(iii) (a) Find the critical points of the area function. Use second derivative test to determine critical points at the maximum area. Also, find the maximum area. 2</p> <p style="text-align: center;">OR</p> <p>(iii) (b) Using first derivative test, calculate the maximum area the company can enclose with the 300 metres of boundary material, considering the parallel partition. 2</p> | |
| <p>Ans</p> | <p>(i) $2x + 3y = 300$</p> <p>(ii) $A = xy = \frac{x}{3}(300 - 2x)$</p> <p>(iii) (a) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$</p> <p>$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$ 1/2</p> <p>For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$ 1/2</p> <p>Also, $\frac{d^2A}{dx^2} = -\frac{4}{3} < 0$. So, A is maximum at $x = 75$ 1/2</p> <p>Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$ 1/2</p> <p style="text-align: center;">OR</p> <p>(iii) (b) $A = \frac{x}{3}(300 - 2x) = \frac{1}{3}(300x - 2x^2)$</p> <p>$\Rightarrow \frac{dA}{dx} = \frac{1}{3}(300 - 4x)$ 1/2</p> <p>For critical points, put $\frac{dA}{dx} = 0 \Rightarrow x = 75$ 1/2</p> <p>As $\frac{dA}{dx}$ changes its sign from positive to negative as x passes through $x = 75$ from left to right, which means $x = 75$ is the point of maximum. 1/2</p> <p>Also, maximum area is $A = \frac{75}{3}(300 - 150) = 3750 \text{ m}^2$ 1/2</p> <p>Note : Full credit to be given if the student takes equation as $2x + 2y = 300$ or $2x + 4y = 300$ or $4x + 4y = 300$ or $4x + 3y = 300$</p> <p>The solutions of sub-parts will differ and marks may be given accordingly.</p> | |