

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025
SUBJECT: MATHEMATICS (Q.P. CODE – 65/5/1)


General Instructions: -

1	You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2	“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.”
3	Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.
4	The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.
5	The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
6	Evaluators will mark (√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing.
7	If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.
8	If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
9	If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” .

10	No marks to be deducted for the cumulative effect of an error. It should be penalized only once.
11	A full scale of marks_(example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.
12	Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.
13	Ensure that you do not make the following common types of errors committed by the Examiner in the past: - <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks awarded.
14	While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.
15	Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
16	The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation.
17	Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.
18	The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

Q.No.	EXPECTED ANSWER / VALUE POINTS	Marks
SECTION-A <i>This section comprises multiple choice questions (MCQs) of 1 mark each.</i>		
1.	<p>If $A = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$, then A^3 is :</p> <p>(A) $3 \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$ (B) $\begin{bmatrix} 125 & 0 & 0 \\ 0 & 125 & 0 \\ 0 & 0 & 125 \end{bmatrix}$</p> <p>(C) $\begin{bmatrix} 15 & 0 & 0 \\ 0 & 15 & 0 \\ 0 & 0 & 15 \end{bmatrix}$ (D) $\begin{bmatrix} 5^3 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$</p>	
Ans	(B) $\begin{bmatrix} 125 & 0 & 0 \\ 0 & 125 & 0 \\ 0 & 0 & 125 \end{bmatrix}$	1
2.	<p>If $P(A \cup B) = 0.9$ and $P(A \cap B) = 0.4$, then $P(\bar{A}) + P(\bar{B})$ is :</p> <p>(A) 0.3 (B) 1</p> <p>(C) 1.3 (D) 0.7</p>	
Ans	(D) 0.7	1
3.	<p>If $A = \begin{bmatrix} 1 & 2 & 3 \\ -4 & 3 & 7 \end{bmatrix}$ and $B = \begin{bmatrix} 4 & 3 \\ -1 & 2 \\ 0 & 5 \end{bmatrix}$, then the correct statement is :</p> <p>(A) Only AB is defined.</p> <p>(B) Only BA is defined.</p> <p>(C) AB and BA, both are defined.</p> <p>(D) AB and BA, both are not defined.</p>	
Ans	(C) AB and BA, both are defined.	1
4.	<p>If $\begin{vmatrix} 2x & 5 \\ 12 & x \end{vmatrix} = \begin{vmatrix} 6 & -5 \\ 4 & 3 \end{vmatrix}$, then the value of x is :</p> <p>(A) 3 (B) 7</p> <p>(C) ± 7 (D) ± 3</p>	
Ans	(C) ± 7	1

5.	<p>If $f(x) = \begin{cases} \frac{\sin^2 ax}{x^2}, & x \neq 0 \\ 1, & x = 0 \end{cases}$</p> <p>is continuous at $x = 0$, then the value of a is :</p> <p>(A) 1 (B) -1 (C) ± 1 (D) 0</p>	
Ans	(C) ± 1	1
6.	<p>If $A = [a_{ij}]$ is a 3×3 diagonal matrix such that $a_{11} = 1$, $a_{22} = 5$ and $a_{33} = -2$, then A is :</p> <p>(A) 0 (B) -10 (C) 10 (D) 1</p>	
Ans	(B) -10	1
7.	<p>The principal value of $\cot^{-1}\left(-\frac{1}{\sqrt{3}}\right)$ is :</p> <p>(A) $-\frac{\pi}{3}$ (B) $-\frac{2\pi}{3}$ (C) $\frac{\pi}{3}$ (D) $\frac{2\pi}{3}$</p>	
Ans	(D) $\frac{2\pi}{3}$	1
8.	<p>If $\begin{bmatrix} 4+x & x-1 \\ -2 & 3 \end{bmatrix}$ is a singular matrix, then the value of x is :</p> <p>(A) 0 (B) 1 (C) -2 (D) -4</p>	
Ans	(C) -2	1
9.	<p>If $f(x) = \{[x], x \in \mathbb{R}\}$ is the greatest integer function, then the correct statement is :</p> <p>(A) f is continuous but not differentiable at $x = 2$. (B) f is neither continuous nor differentiable at $x = 2$. (C) f is continuous as well as differentiable at $x = 2$. (D) f is not continuous but differentiable at $x = 2$.</p>	
Ans	(B) f is neither continuous nor differentiable at $x=2$.	1


15.	<p>The sum of the order and degree of the differential equation</p> $\left[1 + \left(\frac{dy}{dx} \right)^2 \right]^3 = \frac{d^2y}{dx^2}$ <p>is :</p> <p>(A) 2 (B) $\frac{5}{2}$ (C) 3 (D) 4</p>	
Ans	(C) 3	1
16.	<p>For a Linear Programming Problem (LPP), the given objective function $Z = 3x + 2y$ is subject to constraints :</p> $x + 2y \leq 10$ $3x + y \leq 15$ $x, y \geq 0$  <p>The correct feasible region is :</p> <p>(A) ABC (B) AOEC (C) CED (D) Open unbounded region BCD</p>	
Ans	(B) AOEC	1
17.	<p>Let \vec{a} be a position vector whose tip is the point $(2, -3)$. If $\vec{AB} = \vec{a}$, where coordinates of A are $(-4, 5)$, then the coordinates of B are :</p> <p>(A) $(-2, -2)$ (B) $(2, -2)$ (C) $(-2, 2)$ (D) $(2, 2)$</p>	
Ans	(C) $(-2, 2)$	1
18.	<p>The respective values of \vec{a} and \vec{b}, if given</p> $(\vec{a} - \vec{b}) \cdot (\vec{a} + \vec{b}) = 512$ <p>and $\vec{a} = 3 \vec{b}$, are :</p> <p>(A) 48 and 16 (B) 3 and 1 (C) 24 and 8 (D) 6 and 2</p>	
Ans	(C) 24 and 8	1

Questions number 19 and 20 are Assertion and Reason based questions. Two statements are given, one labelled Assertion (A) and the other labelled Reason (R). Select the correct answer from the codes (A), (B), (C) and (D) as given below.

- (A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).
- (B) Both Assertion (A) and Reason (R) are true, but Reason (R) is **not** the correct explanation of the Assertion (A).
- (C) Assertion (A) is true, but Reason (R) is false.
- (D) Assertion (A) is false, but Reason (R) is true.

19.

Assertion (A) : The shaded portion of the graph represents the feasible region for the given Linear Programming Problem (LPP).



$$\text{Min } Z = 50x + 70y$$

subject to constraints

$$2x + y \geq 8, \quad x + 2y \geq 10, \quad x, y \geq 0$$

$Z = 50x + 70y$ has a minimum value = 380 at B(2, 4).

Reason (R) : The region representing $50x + 70y < 380$ does not have any point common with the feasible region.

Ans

(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).

1

20.

Assertion (A) : Let $A = \{x \in \mathbb{R} : -1 \leq x \leq 1\}$. If $f : A \rightarrow A$ be defined as $f(x) = x^2$, then f is not an onto function.

Reason (R) : If $y = -1 \in A$, then $x = \pm \sqrt{-1} \notin A$.

Ans

(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).

1

SECTION-B

This section comprises 5 Very Short Answer (VSA) type questions of 2 marks each.

21.	Find the domain of the function $f(x) = \cos^{-1}(x^2 - 4)$.	
Ans	Domain of $\cos^{-1}x$ is $[-1, 1]$ $\Rightarrow -1 \leq x^2 - 4 \leq 1 \Rightarrow 3 \leq x^2 \leq 5$ $\Rightarrow x \in [-\sqrt{5}, -\sqrt{3}] \cup [\sqrt{3}, \sqrt{5}]$	1 $\frac{1}{2}$ $\frac{1}{2}$
22.	Surface area of a balloon (spherical), when air is blown into it, increases at a rate of $5 \text{ mm}^2/\text{s}$. When the radius of the balloon is 8 mm , find the rate at which the volume of the balloon is increasing.	
Ans	$\frac{dS}{dt} = 5 \text{ mm}^2/\text{s}, \quad \left(\frac{dV}{dt}\right)_{r=8} = ?$ $S = 4\pi r^2 \Rightarrow \frac{dS}{dt} = 8\pi r \cdot \frac{dr}{dt} \Rightarrow \frac{dr}{dt} = \frac{5}{8\pi r}$ $V = \frac{4}{3}\pi r^3 \Rightarrow \frac{dV}{dt} = 4\pi r^2 \cdot \frac{dr}{dt} \Rightarrow \frac{dV}{dt} = \frac{5}{2}r$ $\Rightarrow \left(\frac{dV}{dt}\right)_{r=8} = 20 \text{ mm}^3/\text{s}$	$\frac{1}{2}$ 1 $\frac{1}{2}$
23.	<p>(a) Differentiate $\frac{\sin x}{\sqrt{\cos x}}$ with respect to x.</p> <p style="text-align: center;">OR</p> <p>(b) If $y = 5 \cos x - 3 \sin x$, prove that $\frac{d^2y}{dx^2} + y = 0$.</p>	
Ans	<p>(a) Let $y = \frac{\sin x}{\sqrt{\cos x}}$</p> $\frac{dy}{dx} = \frac{\sqrt{\cos x} \cdot \cos x - \sin x \cdot \left(\frac{-\sin x}{2\sqrt{\cos x}}\right)}{\cos x}$ $\Rightarrow \frac{dy}{dx} = \frac{2\cos^2 x + \sin^2 x}{2(\cos x)^{3/2}} \text{ or } \frac{1 + \cos^2 x}{2(\cos x)^{3/2}}$ <p style="text-align: center;">OR</p> <p>(b) $y = 5\cos x - 3\sin x$, then $\frac{dy}{dx} = -5 \cdot \sin x - 3 \cdot \cos x$</p> $\Rightarrow \frac{d^2y}{dx^2} = -5 \cdot \cos x + 3 \cdot \sin x = -y$ $\Rightarrow \frac{d^2y}{dx^2} + y = 0$	$1\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$


24.	<p>(a) Find a vector of magnitude 5 which is perpendicular to both the vectors $3\hat{i} - 2\hat{j} + \hat{k}$ and $4\hat{i} + 3\hat{j} - 2\hat{k}$.</p> <p style="text-align: center;">OR</p> <p>(b) Let \vec{a}, \vec{b} and \vec{c} be three vectors such that $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ and $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$, $\vec{a} \neq 0$. Show that $\vec{b} = \vec{c}$.</p>	
Ans	<p>Let $\vec{a} = 3\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = 4\hat{i} + 3\hat{j} - 2\hat{k}$</p> $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -2 & 1 \\ 4 & 3 & -2 \end{vmatrix} = \hat{i} + 10\hat{j} + 17\hat{k}$ $ \vec{a} \times \vec{b} = \sqrt{1^2 + 10^2 + 17^2} = \sqrt{390}$ <p>Unit vector $\hat{n} = \frac{\vec{a} \times \vec{b}}{ \vec{a} \times \vec{b} } = \frac{1}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$</p> <p>$\therefore$ Required vector = $\frac{5}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$</p> <p style="text-align: center;">OR</p> <p>(b) $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} \Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0$ \Rightarrow either $\vec{b} = \vec{c}$ or $\vec{a} \perp (\vec{b} - \vec{c})$, since $\vec{a} \neq 0$</p> <p>Also, $\vec{a} \times \vec{b} = \vec{a} \times \vec{c} \Rightarrow \vec{a} \times (\vec{b} - \vec{c}) = 0$ \Rightarrow either $\vec{b} = \vec{c}$ or $\vec{a} \parallel (\vec{b} - \vec{c})$, since $\vec{a} \neq 0$</p> <p>Since vectors \vec{a} and $(\vec{b} - \vec{c})$ cannot be \parallel and \perp simultaneously</p> <p>Hence $\vec{b} = \vec{c}$</p>	<p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p>
25.	<p>A man needs to hang two lanterns on a straight wire whose end points have coordinates A (4, 1, -2) and B (6, 2, -3). Find the coordinates of the points where he hangs the lanterns such that these points trisect the wire AB.</p>	
Ans	<div style="text-align: center;">  </div> <p>Let P and Q trisect the wire AB.</p> <p>P divides AB in the ratio 1:2 then, coordinate of point P = $(\frac{14}{3}, \frac{4}{3}, -\frac{7}{3})$</p> <p>Q divides AB in the ratio 2:1 then, coordinate of point Q = $(\frac{16}{3}, \frac{5}{3}, -\frac{8}{3})$</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p>

SECTION-C

This section comprises 6 Short Answer (SA) type questions of 3 marks each.

26.	Find the value of 'a' for which $f(x) = \sqrt{3} \sin x - \cos x - 2ax + 6$ is decreasing in R.	
Ans	<p>Since $f(x)$ is a decreasing function $\Rightarrow f'(x) \leq 0$</p> <p>$\Rightarrow \sqrt{3} \cdot \cos x + \sin x - 2a \leq 0$</p> <p>$\Rightarrow 2 \left(\frac{\sqrt{3}}{2} \cdot \cos x + \frac{1}{2} \sin x \right) - 2a \leq 0$</p> <p>$\Rightarrow \cos \left(x - \frac{\pi}{6} \right) \leq a$</p> <p>Since, $-1 \leq \cos \left(x - \frac{\pi}{6} \right) \leq 1 \Rightarrow a \geq 1$ i.e. $a \in [1, \infty)$ or $(1, \infty)$</p>	<p>1</p> <p>1</p> <p>1</p>
27.	<p>(a) Find :</p> $\int \frac{2x}{(x^2 + 3)(x^2 - 5)} dx$ <p style="text-align: center;">OR</p> <p>(b) Evaluate :</p> $\int_1^4 (x-2 + x-4) dx$	
Ans	<p>(a) Let $I = \int \frac{2x}{(x^2+3)(x^2-5)} dx$</p> <p>Put $x^2 = t \Rightarrow 2x \cdot dx = dt$</p> <p>$\Rightarrow I = \int \frac{dt}{(t+3)(t-5)}$</p> <p>$= \int \left(-\frac{1}{8(t+3)} + \frac{1}{8(t-5)} \right) dt$</p> <p>$= \frac{1}{8} [\log t-5 - \log t+3] + c$</p> <p>$= \frac{1}{8} \log \left \frac{x^2-5}{x^2+3} \right + c$</p> <p style="text-align: center;">OR</p> <p>(b) $\int_1^4 (x-2 + x-4) dx$</p> <p>$= \int_1^2 (2-x) dx + \int_2^4 (x-2) dx - \int_1^4 (x-4) dx$</p> <p>$= \left[\frac{(2-x)^2}{-2} \right]_1^2 + \left[\frac{(x-2)^2}{2} \right]_2^4 - \left[\frac{(x-4)^2}{2} \right]_1^4$</p> <p>$= \frac{1}{2} + 2 + \frac{9}{2} = 7$</p>	<p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>OR</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p>

28.	<p>Find the particular solution of the differential equation</p> $\left[x \sin^2\left(\frac{y}{x}\right) - y \right] dx + x dy = 0$ <p>given that $y = \frac{\pi}{4}$, when $x = 1$.</p>	
Ans	$\left[x \sin^2 \frac{y}{x} - y \right] \cdot dx + x \cdot dy = 0$ $\Rightarrow \frac{dy}{dx} = \frac{y}{x} - \sin^2 \frac{y}{x}$ <p>Put $y = vx \Rightarrow \frac{dy}{dx} = x \frac{dv}{dx} + v$</p> $\therefore x \frac{dv}{dx} + v = v - \sin^2 v$ $\Rightarrow - \int \operatorname{cosec}^2 v \, dv = \int \frac{dx}{x}$ $\Rightarrow \cot v = \log x + c$ $\Rightarrow \cot \frac{y}{x} = \log x + c$ <p>$x = 1, y = \frac{\pi}{4} \Rightarrow c = 1$</p> $\Rightarrow \cot \frac{y}{x} = \log x + 1$	<p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
29.	<p>In the Linear Programming Problem (LPP), find the point/points giving maximum value for $Z = 5x + 10y$ subject to constraints</p> $x + 2y \leq 120$ $x + y \geq 60$ $x - 2y \geq 0$ $x, y \geq 0$	
Ans		



1½ for correct graph and correct feasible region

Corner Points	Value of Z
A (60, 30)	600
B (120, 0)	600
C (60, 0)	300
D (40, 20)	400

1

Since Z is maximum on points A and B

Hence all points lying on segment AB give maximum Z.

½

30.

(a) If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ such that $|\vec{a}| = 3$, $|\vec{b}| = 5$, $|\vec{c}| = 7$, then find the angle between \vec{a} and \vec{b} .

OR

(b) If \vec{a} and \vec{b} are unit vectors inclined with each other at an angle θ , then prove that $\frac{1}{2} |\vec{a} - \vec{b}| = \sin \frac{\theta}{2}$.

Ans

$$\text{Given } \vec{a} + \vec{b} + \vec{c} = \vec{0} \Rightarrow |\vec{a} + \vec{b}| = |-\vec{c}|$$

$$\Rightarrow |\vec{a} + \vec{b}|^2 = |\vec{c}|^2 \Rightarrow |\vec{a}|^2 + |\vec{b}|^2 + 2\vec{a} \cdot \vec{b} = |\vec{c}|^2$$

$$\Rightarrow 9 + 25 + 2\vec{a} \cdot \vec{b} = 49$$

1

1

	$\Rightarrow 2 \vec{a} \vec{b} \cos\theta = 15$ $\Rightarrow \cos\theta = \frac{1}{2} \therefore \theta = \frac{\pi}{3}$ <p style="text-align: center;">OR</p> <p>(b) $\vec{a} = \vec{b} = 1$</p> $ \vec{a} - \vec{b} ^2 = \vec{a} ^2 + \vec{b} ^2 - 2\vec{a} \cdot \vec{b}$ $= 1 + 1 - 2 \vec{a} \vec{b} \cos\theta$ $= 2 - 2\cos\theta$ $= 2 \left(2\sin^2\frac{\theta}{2} \right) = 4\sin^2\frac{\theta}{2}$ $\Rightarrow \sin\frac{\theta}{2} = \frac{1}{2} \vec{a} - \vec{b} $	<p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">$\frac{1}{2}$</p>
<p>31.</p>	<p>(a) The probability that a student buys a colouring book is 0.7 and that she buys a box of colours is 0.2. The probability that she buys a colouring book, given that she buys a box of colours, is 0.3. Find the probability that the student :</p> <p>(i) Buys both the colouring book and the box of colours.</p> <p>(ii) Buys a box of colours given that she buys the colouring book.</p> <p style="text-align: center;">OR</p> <p>(b) A person has a fruit box that contains 6 apples and 4 oranges. He picks out a fruit three times, one after the other, after replacing the previous one in the box. Find :</p> <p>(i) The probability distribution of the number of oranges he draws.</p> <p>(ii) The expectation of the random variable (number of oranges).</p>	
<p>Ans</p>	<p>(a) Let A be the event of buying colouring book and B be the event of buying coloured box.</p> $P(A) = 0.7, \quad P(B) = 0.2, \quad P(A/B) = 0.3$ <p>(i) $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} \Rightarrow 0.3 = \frac{P(A \cap B)}{0.2}$</p> $\Rightarrow P(A \cap B) = 0.06 \text{ or } \frac{3}{50}$	<p style="text-align: center;">1</p> <p style="text-align: center;">$\frac{1}{2}$</p> <p style="text-align: center;">1</p>

$$(ii) \quad P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)}$$

$$= \frac{0.06}{0.7} = \frac{3}{35} \text{ or } 0.086$$

1

OR

(b) Let X be random variable for number of oranges.

$$X = 0, 1, 2, 3$$

1/2

Let A be the event that orange is drawn.

$$P(A) = \frac{4}{10} = \frac{2}{5}, \quad P(\bar{A}) = 1 - \frac{2}{5} = \frac{3}{5}$$

1/2

(i)

X	0	1	2	3
P(X)	$\frac{27}{125}$	$\frac{54}{125}$	$\frac{36}{125}$	$\frac{8}{125}$

1

$$(ii) \quad E(X) = \sum p_i x_i = 0 \times \frac{27}{125} + 1 \times \frac{54}{125} + 2 \times \frac{36}{125} + 3 \times \frac{8}{125}$$

$$= \frac{150}{125} \text{ or } \frac{6}{5}$$

1/2

1/2


SECTION-D

This section comprises 4 Long Answer (LA) type questions of 5 marks each.

32.

Sketch a graph of $y = x^2$. Using integration, find the area of the region bounded by $y = 9$, $x = 0$ and $y = x^2$.


Ans



1 1/2 for correct figure and shading

	<p>Required area = $\int_0^9 \sqrt{y} \, dy$</p> $= \frac{2}{3} [y^{3/2}]_0^9$ $= 18$ <p>Note: If area is found in second quadrant, may be considered.</p>	<p>1½</p> <p>1</p> <p>1</p>
33.	<p>A furniture workshop produces three types of furniture – chairs, tables and beds each day. On a particular day the total number of furniture pieces produced is 45. It was also found that production of beds exceeds that of chairs by 8, while the total production of beds and chairs together is twice the production of tables. Determine the units produced of each type of furniture, using matrix method.</p>	
Ans	<p>Let the numbers of chairs, tables and beds produced be x, y and z respectively.</p> $\therefore x + y + z = 45; \quad -x + 0.y + z = 8; \quad x - 2y + z = 0$ <p>Let $A = \begin{bmatrix} 1 & 1 & 1 \\ -1 & 0 & 1 \\ 1 & -2 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 45 \\ 8 \\ 0 \end{bmatrix}$</p> $ A = 1(0 + 2) - 1(-1 - 1) + 1(2 - 0) = 6 \neq 0$ <p>$\therefore A^{-1}$ exists</p> $AX = B \Rightarrow X = A^{-1}B$ $\text{adj}(A) = \begin{bmatrix} 2 & -3 & 1 \\ 2 & 0 & -2 \\ 2 & 3 & 1 \end{bmatrix}$ $A^{-1} = \frac{1}{6} \begin{bmatrix} 2 & -3 & 1 \\ 2 & 0 & -2 \\ 2 & 3 & 1 \end{bmatrix}$ $\therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{6} \begin{bmatrix} 2 & -3 & 1 \\ 2 & 0 & -2 \\ 2 & 3 & 1 \end{bmatrix} \begin{bmatrix} 45 \\ 8 \\ 0 \end{bmatrix} = \begin{bmatrix} 11 \\ 15 \\ 19 \end{bmatrix}$ <p>So, $x = 11, y = 15, z = 19$</p> <p>Hence the numbers of chairs, tables and beds produced are 11, 15 and 19 respectively.</p>	<p>1 ½</p> <p>½</p> <p>½</p> <p>1½</p> <p>1</p>
34.	<p>(a) For a positive constant 'a', differentiate $a^{t+\frac{1}{t}}$ with respect to $\left(t + \frac{1}{t}\right)^a$, where t is a non-zero real number.</p> <p style="text-align: center;">OR</p> <p>(b) Find $\frac{dy}{dx}$ if $y^x + x^y + x^x = a^b$, where a and b are constants.</p>	

<p>Ans</p>	<p>(a) Let $u = a^{t+\frac{1}{t}} \Rightarrow \frac{du}{dt} = a^{t+\frac{1}{t}} \cdot \log a \cdot \left(1 - \frac{1}{t^2}\right)$</p> <p>$v = \left(t + \frac{1}{t}\right)^a \Rightarrow \frac{dv}{dt} = a \left(t + \frac{1}{t}\right)^{a-1} \cdot \left(1 - \frac{1}{t^2}\right)$</p> <p>$\frac{du}{dv} = \frac{du/dt}{dv/dt} = \frac{a^{t+\frac{1}{t}} \cdot \log a}{a \left(t + \frac{1}{t}\right)^{a-1}}$</p> <p style="text-align: center;">OR</p> <p>(b) Let $u = y^x$, $v = x^y$ and $w = x^x$</p> <p>$\Rightarrow \frac{du}{dx} + \frac{dv}{dx} + \frac{dw}{dx} = 0$(i)</p> <p>$u = y^x \Rightarrow \log u = x \cdot \log y \Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \frac{x}{y} \cdot \frac{dy}{dx} + \log y$</p> <p>$\Rightarrow \frac{du}{dx} = y^x \left(\frac{x}{y} \cdot \frac{dy}{dx} + \log y\right) = xy^{x-1} \frac{dy}{dx} + y^x \log y$</p> <p>$v = x^y \Rightarrow \log v = y \cdot \log x \Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \frac{y}{x} + \log x \cdot \frac{dy}{dx}$</p> <p>$\Rightarrow \frac{dv}{dx} = x^y \left(\frac{y}{x} + \log x \cdot \frac{dy}{dx}\right) = yx^{y-1} + x^y \log x \cdot \frac{dy}{dx}$</p> <p>$w = x^x \Rightarrow \log w = x \cdot \log x \Rightarrow \frac{1}{w} \cdot \frac{dw}{dx} = 1 + \log x$</p> <p>$\Rightarrow \frac{dw}{dx} = x^x \cdot (1 + \log x)$</p> <p>$\therefore$ From (i), we get</p> <p>$xy^{x-1} \cdot \frac{dy}{dx} + y^x \cdot \log y + yx^{y-1} + x^y \cdot \log x \cdot \frac{dy}{dx} + x^x \cdot (1 + \log x) = 0$</p> <p>$\Rightarrow \frac{dy}{dx} = -\frac{x^x \cdot (1 + \log x) + y^x \cdot \log y + yx^{y-1}}{x \cdot y^{x-1} + x^y \cdot \log x}$</p>	<p>2</p> <p>2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
<p>35.</p>	<p>(a) Find the foot of the perpendicular drawn from the point (1, 1, 4) on the line $\frac{x+2}{5} = \frac{y+1}{2} = \frac{-z+4}{-3}$.</p> <p style="text-align: center;">OR</p> <p>(b) Find the point on the line $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3}$ at a distance of $2\sqrt{2}$ units from the point (-1, -1, 2).</p>	
<p>Ans</p>	<p>(a) Let $\frac{x+2}{5} = \frac{y+1}{2} = \frac{z-4}{3} = \lambda$</p> <p>Coordinate of general point on the given line are $M (5\lambda - 2, 2\lambda - 1, 3\lambda + 4)$</p>	<p>1</p>



Direction Ratios of PM vector are $\langle 5\lambda - 3, 2\lambda - 2, 3\lambda \rangle$

Since, $PM \perp l$

$$\Rightarrow 5(5\lambda - 3) + 2(2\lambda - 2) + 3(3\lambda) = 0$$

$$\Rightarrow \lambda = \frac{1}{2}$$

Hence, coordinates of M are $(\frac{1}{2}, 0, \frac{11}{2})$

OR

(b) Equation of given line be $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3} = \lambda$ (say)

Coordinate of any general point on the line are $P(3\lambda + 1, 2\lambda - 1, 3\lambda + 4)$.

Let distance of point P from $(-1, -1, 2)$ is $2\sqrt{2}$.

$$\Rightarrow \sqrt{(3\lambda + 2)^2 + (2\lambda)^2 + (3\lambda + 2)^2} = 2\sqrt{2}$$

$$\Rightarrow 22\lambda^2 + 24\lambda = 0$$

$$\Rightarrow \lambda = 0 \text{ or } \lambda = -\frac{12}{11}$$

Hence, coordinates of point P are $(1, -1, 4)$ or $(-\frac{25}{11}, -\frac{35}{11}, \frac{8}{11})$

1

1

1

1

1

1½

1

1½

SECTION-E

This section comprises 3 case study-based questions of 4 marks each

36.

Case Study – 1

A carpenter needs to make a wooden cuboidal box, closed from all sides, which has a square base and fixed volume. Since he is short of the paint required to paint the box on completion, he wants the surface area to be minimum.

On the basis of the above information, answer the following questions :

(i) Taking length = breadth = x m and height = y m, express the surface area (S) of the box in terms of x and its volume (V), which is constant.


(ii) Find $\frac{dS}{dx}$.

(iii) (a) Find a relation between x and y such that the surface area (S) is minimum.

OR

(iii) (b) If surface area (S) is constant, the volume (V) = $\frac{1}{4}(Sx - 2x^3)$, x being the edge of base. Show that volume (V) is maximum for $x = \sqrt{\frac{S}{6}}$.

Ans



(i) $V = x^2y \Rightarrow y = \frac{V}{x^2} \dots \dots \dots$ (i)

Hence, $S = 2x^2 + 4xy = 2x^2 + \frac{4V}{x}$

(ii) $\frac{dS}{dx} = 4\left(x - \frac{V}{x^2}\right)$

(iii) (a) $\frac{dS}{dx} = 0 \Rightarrow V = x^3 \Rightarrow x^2y = x^3 \Rightarrow y = x$

$\frac{d^2S}{dx^2} = 4\left(1 + \frac{2V}{x^3}\right) > 0 \Rightarrow S$ is minimum if $y = x$.

OR

(iii) (b) $V = \frac{1}{4}(Sx - 2x^3) \Rightarrow \frac{dV}{dx} = \frac{1}{4}(S - 6x^2)$

1

1

1

1

1

	<p>Put $\frac{dV}{dx} = 0 \Rightarrow x = \sqrt{\frac{5}{6}}$</p> <p>$\left(\frac{d^2V}{dx^2}\right)_{x=\sqrt{\frac{5}{6}}} = -3\sqrt{\frac{5}{6}} < 0 \Rightarrow$ Volume is maximum for $x = \sqrt{\frac{5}{6}}$.</p>	<p>1/2</p> <p>1/2</p>
<p>37.</p>	<p style="text-align: center;">Case Study - 2</p> <p>Let A be the set of 30 students of class XII in a school. Let $f : A \rightarrow N$, N is a set of natural numbers such that function $f(x) =$ Roll Number of student x.</p> <p>On the basis of the given information, answer the following :</p> <p>(i) Is f a bijective function ?</p> <p>(ii) Give reasons to support your answer to (i).</p> <p>(iii) (a) Let R be a relation defined by the teacher to plan the seating arrangement of students in pairs, where $R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = 3x\}$. List the elements of R. Is the relation R reflexive, symmetric and transitive ? Justify your answer.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) Let R be a relation defined by $R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = x^3\}$. List the elements of R. Is R a function ? Justify your answer.</p>	
<p>Ans</p>	<p>(i) No, f is not bijective function</p> <p>(ii) Range = {1, 2, 3, 4,, 30} and codomain = N Since, Range \neq codomain \Rightarrow f is not onto and hence f is not bijective.</p> <p>(iii) (a) $R = \{(1, 3), (2, 6), (3, 9), (4, 12), (5, 15), (6, 18), (7, 21), (8, 24), (9, 27), (10, 30)\}$ Since $(1, 1) \notin R \Rightarrow R$ is not reflexive. $(1, 3) \in R$ but $(3, 1) \notin R \Rightarrow R$ is not symmetric $(1, 3) \in R, (3, 9) \in R$ but $(1, 9) \notin R \Rightarrow R$ is not transitive.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) $R = \{(1, 1), (2, 8), (3, 27)\}$ \therefore elements 4, 5, 6 ... 30 do not have an image. Hence the above relation is not a function.</p>	<p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>

38.

Case Study – 3


A gardener wanted to plant vegetables in his garden. Hence he bought 10 seeds of brinjal plant, 12 seeds of cabbage plant and 8 seeds of radish plant. The shopkeeper assured him of germination probabilities of brinjal, cabbage and radish to be 25%, 35% and 40% respectively. But before he could plant the seeds, they got mixed up in the bag and he had to sow them randomly.



Radish



Cabbage



Brinjal

Based upon the above information, answer the following questions :

- Calculate the probability of a randomly chosen seed to germinate.
- What is the probability that it is a cabbage seed, given that the chosen seed germinates ?

Ans

Let A: Event that chosen seed germinates.

B: Event that Brinjal seed is chosen.

C: Event that Cabbage seed is chosen.

R: Event that Radish seed is chosen.

$$P(B) = \frac{10}{30}; P(C) = \frac{12}{30}; P(R) = \frac{8}{30};$$

$$P\left(\frac{A}{B}\right) = \frac{25}{100}; P\left(\frac{A}{C}\right) = \frac{35}{100}; P\left(\frac{A}{R}\right) = \frac{40}{100}$$

$$(i) P(A) = P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)$$

$$= \frac{10}{30} \times \frac{25}{100} + \frac{12}{30} \times \frac{35}{100} + \frac{8}{30} \times \frac{40}{100}$$

$$= \frac{990}{3000} \text{ or } \frac{33}{100}$$

$$(ii) (a) P\left(\frac{C}{A}\right) = \frac{P(C) \cdot P\left(\frac{A}{C}\right)}{P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)}$$

$$= \frac{\frac{12}{30} \times \frac{35}{100}}{\frac{990}{3000}}$$

$$= \frac{42}{99} \text{ or } \frac{14}{33}$$

1

1

1

1


Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025
SUBJECT: MATHEMATICS (Q.P. CODE – 65/5/2)

General Instructions: -

1	You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2	“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.”
3	Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.
4	The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.
5	The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
6	Evaluators will mark (√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing.
7	If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.
8	If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
9	If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” .

10	No marks to be deducted for the cumulative effect of an error. It should be penalized only once.
11	A full scale of marks_(example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.
12	Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.
13	Ensure that you do not make the following common types of errors committed by the Examiner in the past: - <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks awarded.
14	While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.
15	Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
16	The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation.
17	Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.
18	The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

5.	<p>If $A = \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$, then A^3 is :</p> <p>(A) $3 \begin{bmatrix} 5 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$ (B) $\begin{bmatrix} 125 & 0 & 0 \\ 0 & 125 & 0 \\ 0 & 0 & 125 \end{bmatrix}$</p> <p>(C) $\begin{bmatrix} 15 & 0 & 0 \\ 0 & 15 & 0 \\ 0 & 0 & 15 \end{bmatrix}$ (D) $\begin{bmatrix} 5^3 & 0 & 0 \\ 0 & 5 & 0 \\ 0 & 0 & 5 \end{bmatrix}$</p>	
Ans	(B) $\begin{bmatrix} 125 & 0 & 0 \\ 0 & 125 & 0 \\ 0 & 0 & 125 \end{bmatrix}$	1
6.	<p>If $\begin{vmatrix} 2x & 5 \\ 12 & x \end{vmatrix} = \begin{vmatrix} 6 & -5 \\ 4 & 3 \end{vmatrix}$, then the value of x is :</p> <p>(A) 3 (B) 7</p> <p>(C) ± 7 (D) ± 3</p>	
Ans	(C) ± 7	1
7.	<p>If $P(A \cup B) = 0.9$ and $P(A \cap B) = 0.4$, then $P(\bar{A}) + P(\bar{B})$ is :</p> <p>(A) 0.3 (B) 1</p> <p>(C) 1.3 (D) 0.7</p>	
Ans	(D) 0.7	1
8.	<p>If a matrix A is both symmetric and skew-symmetric, then A is a :</p> <p>(A) diagonal matrix (B) zero matrix</p> <p>(C) non-singular matrix (D) scalar matrix</p>	
Ans	(B) zero matrix	1
9.	<p>The slope of the curve $y = -x^3 + 3x^2 + 8x - 20$ is maximum at :</p> <p>(A) (1, -10) (B) (1, 10)</p> <p>(C) (10, 1) (D) (-10, 1)</p>	
Ans	(A) (1, -10)	1

14.	<p>The integrating factor of the differential equation</p> $\frac{dx}{dy} = \frac{x \log x}{\frac{2}{x} \log x - y}$ <p>is :</p> <p>(A) $\frac{1}{8x}$ (B) e</p> <p>(C) $e^{\log x}$ (D) $\log x$</p>	
Ans	(D) $\log x$	1
15.	<p>Let \vec{a} be a position vector whose tip is the point $(2, -3)$. If $\vec{AB} = \vec{a}$, where coordinates of A are $(-4, 5)$, then the coordinates of B are :</p> <p>(A) $(-2, -2)$ (B) $(2, -2)$ (C) $(-2, 2)$ (D) $(2, 2)$</p>	
Ans	(C) $(-2, 2)$	1
16.	<p>The respective values of \vec{a} and \vec{b}, if given</p> $(\vec{a} - \vec{b}) \cdot (\vec{a} + \vec{b}) = 512 \text{ and } \vec{a} = 3 \vec{b} ,$ <p>are :</p> <p>(A) 48 and 16 (B) 3 and 1</p> <p>(C) 24 and 8 (D) 6 and 2</p>	
Ans	(C) 24 and 8	1
17.	<p>For a Linear Programming Problem (LPP), the given objective function $Z = 3x + 2y$ is subject to constraints :</p> $x + 2y \leq 10$ $3x + y \leq 15$ $x, y \geq 0$  <p>The correct feasible region is :</p> <p>(A) ABC (B) AOEC</p> <p>(C) CED (D) Open unbounded region BCD</p>	

Ans	(B) AOEC	1
18.	<p>The sum of the order and degree of the differential equation</p> $\left[1 + \left(\frac{dy}{dx}\right)^2\right]^3 = \frac{d^2y}{dx^2} \text{ is :}$ <p>(A) 2 (B) $\frac{5}{2}$ (C) 3 (D) 4</p>	
Ans	(C) 3	1
<p><i>Questions number 19 and 20 are Assertion and Reason based questions. Two statements are given, one labelled Assertion (A) and the other labelled Reason (R). Select the correct answer from the codes (A), (B), (C) and (D) as given below.</i></p> <p>(A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).</p> <p>(B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).</p> <p>(C) Assertion (A) is true, but Reason (R) is false.</p> <p>(D) Assertion (A) is false, but Reason (R) is true.</p>		
19.	<p><i>Assertion (A) :</i> The shaded portion of the graph represents the feasible region for the given Linear Programming Problem (LPP).</p> <p>Min $Z = 50x + 70y$ subject to constraints $2x + y \geq 8$, $x + 2y \geq 10$, $x, y \geq 0$ $Z = 50x + 70y$ has a minimum value = 380 at B(2, 4).</p> <p><i>Reason (R) :</i> The region representing $50x + 70y < 380$ does not have any point common with the feasible region.</p>	

Ans	(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).	1
20.	<p><i>Assertion (A)</i> : Let $A = \{x \in \mathbb{R} : -1 \leq x \leq 1\}$. If $f : A \rightarrow A$ be defined as $f(x) = x^2$, then f is not an onto function.</p> <p><i>Reason (R)</i> : If $y = -1 \in A$, then $x = \pm \sqrt{-1} \notin A$.</p>	
Ans	(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).	1
SECTION-B <i>This section comprises 5 Very Short Answer (VSA) type questions of 2 marks each.</i>		
21.	Find the domain of $\sec^{-1}(2x + 1)$.	
Ans	<p>Domain of $\sec^{-1} x$ is $(-\infty, -1] \cup [1, \infty)$</p> <p>$\Rightarrow 2x + 1 \leq -1$ or $2x + 1 \geq 1 \Rightarrow x \leq -1$ or $x \geq 0$</p> <p>\Rightarrow Domain = $(-\infty, -1] \cup [0, \infty)$</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
22.	The radius of a cylinder is decreasing at a rate of 2 cm/s and the altitude is increasing at the rate of 3 cm/s. Find the rate of change of volume of this cylinder when its radius is 4 cm and altitude is 6 cm.	
Ans	<p>$\frac{dr}{dt} = -2 \text{ cm/s}, \frac{dh}{dt} = 3 \text{ cm/s}, \left(\frac{dV}{dt}\right)_{r=4, h=6} = ?$</p> <p>$V = \pi r^2 h \Rightarrow \frac{dV}{dt} = 2\pi r \cdot \frac{dr}{dt} \cdot h + \pi r^2 \frac{dh}{dt}$</p> <p>When $r = 4 \text{ cm}$ and $h = 6 \text{ cm}$,</p> <p>$\frac{dV}{dt} = 2\pi(4)(-2)(6) + \pi(4)^2(3) = -48\pi \text{ cm}^3/\text{s}$</p> <p>Volume is decreasing at the rate of $48\pi \text{ cm}^3/\text{s}$</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p>

23.	<p>(a) Find a vector of magnitude 5 which is perpendicular to both the vectors $3\hat{i} - 2\hat{j} + \hat{k}$ and $4\hat{i} + 3\hat{j} - 2\hat{k}$.</p> <p style="text-align: center;">OR</p> <p>(b) Let \vec{a}, \vec{b} and \vec{c} be three vectors such that $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ and $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$, $\vec{a} \neq 0$. Show that $\vec{b} = \vec{c}$.</p>	
Ans	<p>Let $\vec{a} = 3\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = 4\hat{i} + 3\hat{j} - 2\hat{k}$</p> $\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -2 & 1 \\ 4 & 3 & -2 \end{vmatrix} = \hat{i} + 10\hat{j} + 17\hat{k}$ $ \vec{a} \times \vec{b} = \sqrt{1^2 + 10^2 + 17^2} = \sqrt{390}$ <p>Unit vector $\hat{n} = \frac{\vec{a} \times \vec{b}}{ \vec{a} \times \vec{b} } = \frac{1}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$</p> <p>$\therefore$ Required vector = $\frac{5}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$</p> <p style="text-align: center;">OR</p> <p>(b) $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} \Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0$ \Rightarrow either $\vec{b} = \vec{c}$ or $\vec{a} \perp (\vec{b} - \vec{c})$, since $\vec{a} \neq 0$</p> <p>Also, $\vec{a} \times \vec{b} = \vec{a} \times \vec{c} \Rightarrow \vec{a} \times (\vec{b} - \vec{c}) = 0$ \Rightarrow either $\vec{b} = \vec{c}$ or $\vec{a} \parallel (\vec{b} - \vec{c})$, since $\vec{a} \neq 0$</p> <p>Since vectors \vec{a} and $(\vec{b} - \vec{c})$ cannot be \parallel and \perp simultaneously</p> <p>Hence $\vec{b} = \vec{c}$</p>	<p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1/2</p>
24.	<p>A man needs to hang two lanterns on a straight wire whose end points have coordinates A (4, 1, -2) and B (6, 2, -3). Find the coordinates of the points where he hangs the lanterns such that these points trisect the wire AB.</p>	
Ans	<div style="text-align: center;"> <p style="text-align: center;">A (4, 1, -2) — P — Q — B (6, 2, -3)</p> </div> <p>Let P and Q trisect the wire AB.</p> <p>P divides AB in the ratio 1:2 then, coordinate of point P = $(\frac{14}{3}, \frac{4}{3}, -\frac{7}{3})$</p> <p>Q divides AB in the ratio 2:1 then, coordinate of point Q = $(\frac{16}{3}, \frac{5}{3}, -\frac{8}{3})$</p>	<p style="text-align: right;">1</p> <p style="text-align: right;">1</p>

25.	<p>(a) Differentiate $\frac{\sin x}{\sqrt{\cos x}}$ with respect to x.</p> <p style="text-align: center;">OR</p> <p>(b) If $y = 5 \cos x - 3 \sin x$, prove that $\frac{d^2y}{dx^2} + y = 0$.</p>	
Ans	<p>(a) Let $y = \frac{\sin x}{\sqrt{\cos x}}$</p> $\frac{dy}{dx} = \frac{\sqrt{\cos x} \cdot \cos x - \sin x \cdot \left(\frac{-\sin x}{2\sqrt{\cos x}}\right)}{\cos x}$ $\Rightarrow \frac{dy}{dx} = \frac{2\cos^2 x + \sin^2 x}{2(\cos x)^{3/2}} \text{ or } \frac{1 + \cos^2 x}{2(\cos x)^{3/2}}$ <p style="text-align: center;">OR</p> <p>(b) $y = 5\cos x - 3\sin x$, then $\frac{dy}{dx} = -5 \cdot \sin x - 3 \cdot \cos x$</p> $\Rightarrow \frac{d^2y}{dx^2} = -5 \cdot \cos x + 3 \cdot \sin x = -y$ $\Rightarrow \frac{d^2y}{dx^2} + y = 0$	<p style="text-align: right;">1½</p> <p style="text-align: right;">½</p> <p style="text-align: right;">1</p> <p style="text-align: right;">½</p> <p style="text-align: right;">½</p>
<p>SECTION-C</p> <p><i>This section comprises 6 Short Answer (SA) type questions of 3 marks each.</i></p>		
26.	<p>Show that $f(x) = \tan^{-1}(\sin x + \cos x)$ is an increasing function in $\left[0, \frac{\pi}{4}\right]$.</p>	
Ans	$f'(x) = \frac{1}{1 + (\sin x + \cos x)^2} (\cos x - \sin x)$ <p>For $x \in \left[0, \frac{\pi}{4}\right]$, $\cos x \geq \sin x$</p> $\Rightarrow f'(x) \geq 0, f \text{ is an increasing function in } \left[0, \frac{\pi}{4}\right]$	<p style="text-align: right;">1½</p> <p style="text-align: right;">1</p> <p style="text-align: right;">½</p>

27.	<p>(a) The probability that a student buys a colouring book is 0.7 and that she buys a box of colours is 0.2. The probability that she buys a colouring book, given that she buys a box of colours, is 0.3. Find the probability that the student :</p> <p>(i) Buys both the colouring book and the box of colours.</p> <p>(ii) Buys a box of colours given that she buys the colouring book.</p> <p style="text-align: center;">OR</p> <p>(b) A person has a fruit box that contains 6 apples and 4 oranges. He picks out a fruit three times, one after the other, after replacing the previous one in the box. Find :</p> <p>(i) The probability distribution of the number of oranges he draws.</p> <p>(ii) The expectation of the random variable (number of oranges).</p>	
Ans	<p>(a) Let A be the event of buying colouring book and B be the event of buying coloured box. } $P(A) = 0.7, \quad P(B) = 0.2, \quad P(A/B) = 0.3$</p> <p>(i) $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} \Rightarrow 0.3 = \frac{P(A \cap B)}{0.2}$ $\Rightarrow P(A \cap B) = 0.06$ or $\frac{3}{50}$</p> <p>(ii) $P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)}$ $= \frac{0.06}{0.7} = \frac{3}{35}$ or 0.086</p> <p style="text-align: center;">OR</p> <p>(b) Let X be random variable for number of oranges. $X = 0, 1, 2, 3$ Let A be the event that orange is drawn. $P(A) = \frac{4}{10} = \frac{2}{5}, \quad P(\bar{A}) = 1 - \frac{2}{5} = \frac{3}{5}$</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>

	<p>(i)</p> <table border="1" data-bbox="336 259 1217 416"> <tr> <td>X</td> <td>0</td> <td>1</td> <td>2</td> <td>3</td> </tr> <tr> <td>P(X)</td> <td>$\frac{27}{125}$</td> <td>$\frac{54}{125}$</td> <td>$\frac{36}{125}$</td> <td>$\frac{8}{125}$</td> </tr> </table> <p>(ii) $E(X) = \sum p_i x_i = 0 \times \frac{27}{125} + 1 \times \frac{54}{125} + 2 \times \frac{36}{125} + 3 \times \frac{8}{125}$ $= \frac{150}{125}$ or $\frac{6}{5}$</p>	X	0	1	2	3	P(X)	$\frac{27}{125}$	$\frac{54}{125}$	$\frac{36}{125}$	$\frac{8}{125}$	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
X	0	1	2	3								
P(X)	$\frac{27}{125}$	$\frac{54}{125}$	$\frac{36}{125}$	$\frac{8}{125}$								
<p>28.</p>	<p>Find the particular solution of the differential equation $\frac{dy}{dx} - \frac{y}{x} + \operatorname{cosec}\left(\frac{y}{x}\right) = 0$; given that $y = 0$, when $x = 1$.</p>											
<p>Ans</p>	<p>$\frac{dy}{dx} = \frac{y}{x} - \operatorname{cosec} \frac{y}{x}$</p> <p>Put $y = vx$</p> <p>$\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$</p> <p>$\Rightarrow v + x \frac{dv}{dx} = v - \operatorname{cosec} v$</p> <p>$\Rightarrow - \int \sin v \, dv = \int \frac{dx}{x}$</p> <p>$\Rightarrow \cos v = \log x + c$</p> <p>$\Rightarrow \cos \frac{y}{x} = \log x + c$</p> <p>$x = 1, y = 0 \Rightarrow c = 1$</p> <p>$\Rightarrow \cos \frac{y}{x} = \log x + 1$</p>	<p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>										

<p>29.</p>	<p>(a) Find :</p> $\int \frac{2x}{(x^2 + 3)(x^2 - 5)} dx$ <p style="text-align: center;">OR</p> <p>(b) Evaluate :</p> $\int_1^4 (x - 2 + x - 4) dx$	
<p>Ans</p>	<p>(a) Let $I = \int \frac{2x}{(x^2+3)(x^2-5)} dx$</p> <p>Put $x^2 = t \Rightarrow 2x \cdot dx = dt$</p> $\Rightarrow I = \int \frac{dt}{(t+3)(t-5)}$ $= \int \left(-\frac{1}{8(t+3)} + \frac{1}{8(t-5)} \right) dt$ $= \frac{1}{8} [\log t - 5 - \log t + 3] + c$ $= \frac{1}{8} \log \left \frac{x^2-5}{x^2+3} \right + c$ <p style="text-align: center;">OR</p> <p>(b) $\int_1^4 (x - 2 + x - 4) dx$</p> $= \int_1^2 (2 - x) dx + \int_2^4 (x - 2) dx - \int_1^4 (x - 4) dx$ $= \left[\frac{(2-x)^2}{-2} \right]_1^2 + \left[\frac{(x-2)^2}{2} \right]_2^4 - \left[\frac{(x-4)^2}{2} \right]_1^4$ $= \frac{1}{2} + 2 + \frac{9}{2} = 7$	<p style="text-align: right;">1/2</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1/2</p> <p style="text-align: right;">1 1/2</p> <p style="text-align: right;">1</p> <p style="text-align: right;">1/2</p>

30.

In the Linear Programming Problem (LPP), find the point/points giving maximum value for $Z = 5x + 10y$ subject to constraints


$$x + 2y \leq 120$$

$$x + y \geq 60$$

$$x - 2y \geq 0$$

$$x, y \geq 0$$

Ans



Corner Points	Value of Z
A (60, 30)	600
B (120, 0)	600
C (60, 0)	300
D (40, 20)	400

Since Z is maximum on points A and B


Hence all points lying on segment AB give maximum Z .


1½ for correct graph and correct feasible region

1



½

31.	<p>(a) If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ such that $\vec{a} = 3$, $\vec{b} = 5$, $\vec{c} = 7$, then find the angle between \vec{a} and \vec{b}.</p> <p style="text-align: center;">OR</p> <p>(b) If \vec{a} and \vec{b} are unit vectors inclined with each other at an angle θ, then prove that $\frac{1}{2} \vec{a} - \vec{b} = \sin \frac{\theta}{2}$.</p>	
Ans	<p>Given $\vec{a} + \vec{b} + \vec{c} = \vec{0} \Rightarrow \vec{a} + \vec{b} = -\vec{c}$</p> <p>$\Rightarrow \vec{a} + \vec{b} ^2 = \vec{c} ^2 \Rightarrow \vec{a} ^2 + \vec{b} ^2 + 2\vec{a} \cdot \vec{b} = \vec{c} ^2$</p> <p>$\Rightarrow 9 + 25 + 2\vec{a} \cdot \vec{b} = 49$</p> <p>$\Rightarrow 2 \vec{a} \vec{b} \cos \theta = 15$</p> <p>$\Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$</p> <p style="text-align: center;">OR</p> <p>(b) $\vec{a} = \vec{b} = 1$</p> <p>$\vec{a} - \vec{b} ^2 = \vec{a} ^2 + \vec{b} ^2 - 2\vec{a} \cdot \vec{b}$</p> <p>$= 1 + 1 - 2 \vec{a} \vec{b} \cos \theta$</p> <p>$= 2 - 2 \cos \theta$</p> <p>$= 2 \left(2 \sin^2 \frac{\theta}{2} \right) = 4 \sin^2 \frac{\theta}{2}$</p> <p>$\Rightarrow \sin \frac{\theta}{2} = \frac{1}{2} \vec{a} - \vec{b}$</p>	<p style="text-align: center;">1</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1/2</p> <p style="text-align: center;">1</p> <p style="text-align: center;">1/2</p> <p style="text-align: center;">1/2</p> <p style="text-align: center;">1/2</p>
<p>SECTION-D</p> <p><i>This section comprises 4 Long Answer (LA) type questions of 5 marks each.</i></p>		
32.	<p>Draw a rough sketch of the curve $y = \sqrt{x}$. Using integration, find the area of the region bounded by the curve $y = \sqrt{x}$, $x = 4$ and x-axis, in the first quadrant.</p>	


<p>Ans</p>	 <p>Required area = $\int_0^4 \sqrt{x} \, dx$</p> $= \frac{2}{3} [x^{3/2}]_0^4$ $= \frac{2}{3} \times 8 = \frac{16}{3}$	<p>2 marks for correct figure and shading</p> <p>1</p> <p>1</p> <p>1</p>
<p>33.</p>	<p>An amount of ₹ 10,000 is put into three investments at the rate of 10%, 12% and 15% per annum. The combined annual income of all three investments is ₹ 1,310, however the combined annual income of the first and the second investments is ₹ 190 short of the income from the third. Use matrix method and find the investment amount in each at the beginning of the year.</p>	
<p>Ans</p>	<p>Let x, y and z (in ₹) be three investment amounts.</p> <p>Then $x + y + z = 10,000$</p> $\frac{10}{100}x + \frac{12}{100}y + \frac{15}{100}z = 1310$ $- \frac{10}{100}x - \frac{12}{100}y + \frac{15}{100}z = 190$ <p>Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 10 & 12 & 15 \\ -10 & -12 & 15 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 10,000 \\ 1,310,000 \\ 19,000 \end{bmatrix}$</p> <p>$A = 60 \neq 0$</p> <p>$\therefore A^{-1}$ exists</p>	<p>1½</p> <p>½</p> <p>½</p>

	$AX = B \Rightarrow X = A^{-1}B$ $\text{adj}(A) = \begin{bmatrix} 360 & -27 & 3 \\ -300 & 25 & -5 \\ 0 & 2 & 2 \end{bmatrix}$ $A^{-1} = \frac{1}{60} \begin{bmatrix} 360 & -27 & 3 \\ -300 & 25 & -5 \\ 0 & 2 & 2 \end{bmatrix}$ $\therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{60} \begin{bmatrix} 360 & -27 & 3 \\ -300 & 25 & -5 \\ 0 & 2 & 2 \end{bmatrix} \begin{bmatrix} 10,000 \\ 1,31,000 \\ 19,000 \end{bmatrix} = \begin{bmatrix} 2000 \\ 3000 \\ 5000 \end{bmatrix}$ <p>Hence the investments are ₹ 2000, ₹ 3000 and ₹ 5000 respectively.</p>	<p>1½</p> <p>1</p>
<p>34.</p>	<p>(a) Find the foot of the perpendicular drawn from the point (1, 1, 4) on the line $\frac{x+2}{5} = \frac{y+1}{2} = \frac{-z+4}{-3}$.</p> <p style="text-align: center;">OR</p> <p>(b) Find the point on the line $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3}$ at a distance of $2\sqrt{2}$ units from the point (-1, -1, 2).</p>	
<p>Ans</p>	<p>(a) Let $\frac{x+2}{5} = \frac{y+1}{2} = \frac{z-4}{-3} = \lambda$</p> <p>Coordinate of general point on the given line are M (5λ - 2, 2λ - 1, 3λ + 4)</p> <div style="text-align: center;">  </div> <p>Direction ratios of PM vector are $\langle 5\lambda - 3, 2\lambda - 2, 3\lambda \rangle$</p> <p>Since, $PM \perp l$</p> $\Rightarrow 5(5\lambda - 3) + 2(2\lambda - 2) + 3(3\lambda) = 0$ $\Rightarrow \lambda = \frac{1}{2}$ <p>Hence, coordinates of M are $(\frac{1}{2}, 0, \frac{11}{2})$</p> <p style="text-align: center;">OR</p>	<p>1</p> <p>1</p> <p>1</p> <p>1</p>

	<p>(b) Equation of given line be $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3} = \lambda$ (say)</p> <p>Coordinate of any general point on the line are $P(3\lambda + 1, 2\lambda - 1, 3\lambda + 4)$.</p> <p>Let distance of point P from $(-1, -1, 2)$ is $2\sqrt{2}$.</p> $\Rightarrow \sqrt{(3\lambda + 2)^2 + (2\lambda)^2 + (3\lambda + 2)^2} = 2\sqrt{2}$ $\Rightarrow 22\lambda^2 + 24\lambda = 0$ $\Rightarrow \lambda = 0 \text{ or } \lambda = -\frac{12}{11}$ <p>Hence, coordinates of point P are $(1, -1, 4)$ or $(-\frac{25}{11}, -\frac{35}{11}, \frac{8}{11})$</p>	<p>1</p> <p>1½</p> <p>1</p> <p>1½</p>
35.	<p>(a) For a positive constant 'a', differentiate $a^{t+\frac{1}{t}}$ with respect to $(t+\frac{1}{t})^a$, where t is a non-zero real number.</p> <p style="text-align: center;">OR</p> <p>(b) Find $\frac{dy}{dx}$ if $y^x + x^y + x^x = a^b$, where a and b are constants.</p>	
Ans	<p>(a) Let $u = a^{t+\frac{1}{t}} \Rightarrow \frac{du}{dt} = a^{t+\frac{1}{t}} \cdot \log a \cdot (1 - \frac{1}{t^2})$</p> $v = (t + \frac{1}{t})^a \Rightarrow \frac{dv}{dt} = a(t + \frac{1}{t})^{a-1} \cdot (1 - \frac{1}{t^2})$ $\frac{du}{dv} = \frac{du/dt}{dv/dt} = \frac{a^{t+\frac{1}{t}} \cdot \log a}{a(t+\frac{1}{t})^{a-1}}$ <p style="text-align: center;">OR</p> <p>(b) Let $u = y^x$, $v = x^y$ and $w = x^x$</p> $\Rightarrow \frac{du}{dx} + \frac{dv}{dx} + \frac{dw}{dx} = 0 \dots\dots\dots(i)$ $u = y^x \Rightarrow \log u = x \cdot \log y \Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \frac{x}{y} \cdot \frac{dy}{dx} + \log y$ $\Rightarrow \frac{du}{dx} = y^x \left(\frac{x}{y} \cdot \frac{dy}{dx} + \log y \right) = xy^{x-1} \frac{dy}{dx} + y^x \log y$ $v = x^y \Rightarrow \log v = y \cdot \log x \Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \frac{y}{x} + \log x \cdot \frac{dy}{dx}$	<p>2</p> <p>2</p> <p>1</p> <p>1</p> <p>1</p>

	$\Rightarrow \frac{dv}{dx} = x^y \left(\frac{y}{x} + \log x \cdot \frac{dy}{dx} \right) = yx^{y-1} + x^y \log x \frac{dy}{dx}$ $w = x^x \Rightarrow \log w = x \cdot \log x \Rightarrow \frac{1}{w} \cdot \frac{dw}{dx} = 1 + \log x$ $\Rightarrow \frac{dw}{dx} = x^x \cdot (1 + \log x)$ <p>\therefore From (i), we get</p> $xy^{x-1} \cdot \frac{dy}{dx} + y^x \cdot \log y + yx^{y-1} + x^y \cdot \log x \cdot \frac{dy}{dx} + x^x \cdot (1 + \log x) = 0$ $\Rightarrow \frac{dy}{dx} = - \frac{x^x \cdot (1 + \log x) + y^x \cdot \log y + yx^{y-1}}{x \cdot y^{x-1} + x^y \cdot \log x}$	<p>1</p> <p>1</p> <p>1</p>
<p>SECTION-E</p> <p><i>This section comprises 3 case study-based questions of 4 marks each</i></p>		
<p>36.</p>	<p style="text-align: center;">Case Study – 1</p> <p>A gardener wanted to plant vegetables in his garden. Hence he bought 10 seeds of brinjal plant, 12 seeds of cabbage plant and 8 seeds of radish plant. The shopkeeper assured him of germination probabilities of brinjal, cabbage and radish to be 25%, 35% and 40% respectively. But before he could plant the seeds, they got mixed up in the bag and he had to sow them randomly.</p> <div style="display: flex; justify-content: space-around; align-items: center;"> <div style="text-align: center;">  <p>Radish</p> </div> <div style="text-align: center;">  <p>Cabbage</p> </div> <div style="text-align: center;">  <p>Brinjal</p> </div> </div> <p>Based upon the above information, answer the following questions :</p> <p>(i) Calculate the probability of a randomly chosen seed to germinate. 2</p> <p>(ii) What is the probability that it is a cabbage seed, given that the chosen seed germinates ? 2</p>	
<p>Ans</p>	<p>Let A: Event that chosen seed germinates.</p> <p>B: Event that Brinjal seed is chosen.</p> <p>C: Event that Cabbage seed is chosen.</p> <p>R: Event that Radish seed is chosen.</p>	

	$P(B) = \frac{10}{30}; P(C) = \frac{12}{30}; P(R) = \frac{8}{30};$ $P\left(\frac{A}{B}\right) = \frac{25}{100}; P\left(\frac{A}{C}\right) = \frac{35}{100}; P\left(\frac{A}{R}\right) = \frac{40}{100}$ <p>(i) $P(A) = P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)$</p> $= \frac{10}{30} \times \frac{25}{100} + \frac{12}{30} \times \frac{35}{100} + \frac{8}{30} \times \frac{40}{100}$ $= \frac{990}{3000} \text{ or } \frac{33}{100}$ <p>(ii) (a) $P\left(\frac{C}{A}\right) = \frac{P(C) \cdot P\left(\frac{A}{C}\right)}{P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)}$</p> $= \frac{\frac{12}{30} \times \frac{35}{100}}{\frac{990}{3000}}$ $= \frac{42}{99} \text{ or } \frac{14}{33}$	<p>1</p> <p>1</p> <p>1</p> <p>1</p>
37.	<p style="text-align: center;">Case Study – 2</p> <p>A carpenter needs to make a wooden cuboidal box, closed from all sides, which has a square base and fixed volume. Since he is short of the paint required to paint the box on completion, he wants the surface area to be minimum.</p> <p>On the basis of the above information, answer the following questions :</p> <p>(i) Taking length = breadth = x m and height = y m, express the surface area (S) of the box in terms of x and its volume (V), which is constant.</p> <p>(ii) Find $\frac{dS}{dx}$.</p> <p>(iii) (a) Find a relation between x and y such that the surface area (S) is minimum.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) If surface area (S) is constant, the volume (V) = $\frac{1}{4}(Sx - 2x^3)$, x being the edge of base. Show that volume (V) is maximum for $x = \sqrt{\frac{S}{6}}$.</p>	<p>1</p> <p>1</p> <p>2</p> <p>2</p>
Ans		



(i) $V = x^2y \Rightarrow y = \frac{V}{x^2} \dots \dots \dots$ (i)

Hence, $S = 2x^2 + 4xy = 2x^2 + \frac{4V}{x}$

(ii) $\frac{dS}{dx} = 4 \left(x - \frac{V}{x^2} \right)$

(iii) (a) $\frac{dS}{dx} = 0 \Rightarrow V = x^3 \Rightarrow x^2y = x^3 \Rightarrow y = x$

$\frac{d^2S}{dx^2} = 4 \left(1 + \frac{2V}{x^3} \right) = 12 > 0 \Rightarrow S$ is minimum if $y = x$.

OR

(iii) (b) $V = \frac{1}{4}(Sx - 2x^3) \Rightarrow \frac{dV}{dx} = \frac{1}{4}(S - 6x^2)$

Put $\frac{dV}{dx} = 0 \Rightarrow x = \sqrt{\frac{S}{6}}$

$\left(\frac{d^2V}{dx^2} \right)_{x=\sqrt{\frac{S}{6}}} = -3\sqrt{\frac{S}{6}} < 0 \Rightarrow$ Volume is maximum for $x = \sqrt{\frac{S}{6}}$.

1
1
1
1
1
1/2
1/2

Case Study – 3

Let A be the set of 30 students of class XII in a school. Let $f : A \rightarrow N$, N is a set of natural numbers such that function $f(x) =$ Roll Number of student x. On the basis of the given information, answer the following :

38.

(i) Is f a bijective function ?

1

(ii) Give reasons to support your answer to (i).

1

(iii) (a) Let R be a relation defined by the teacher to plan the seating arrangement of students in pairs, where $R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = 3x\}$. List the elements of R. Is the relation R reflexive, symmetric and transitive ? Justify your answer.

2

OR

(iii) (b) Let R be a relation defined by

$R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = x^3\}$. List the elements of R. Is R a function ? Justify your answer.

2

Ans (i) No, f is not bijective function

1

	<p>(ii) Range = {1, 2, 3, 4,, 30} and codomain= N Since, Range \neq codomain \Rightarrow f is not onto and hence f is not bijective.</p> <p>(iii) (a) $R = \{(1, 3), (2, 6), (3, 9), (4, 12), (5, 15), (6, 18), (7, 21), (8, 24), (9, 27), (10, 30)\}$ Since $(1, 1) \notin R \Rightarrow R$ is not reflexive. $(1, 3) \in R$ but $(3, 1) \notin R \Rightarrow R$ is not symmetric $(1, 3) \in R, (3, 9) \in R$ but $(1, 9) \notin R \Rightarrow R$ is not transitive.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) $R = \{(1, 1), (2, 8), (3, 27)\}$ \because elements 4, 5, 6 ... 30 do not have an image. Hence the above relation is not a function.</p>	<p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>
--	---	---

Marking Scheme
Strictly Confidential
(For Internal and Restricted use only)
Senior Secondary Examination, 2025
SUBJECT: MATHEMATICS (Q.P. CODE – 65/5/3)


General Instructions: -

1	You are aware that evaluation is the most important process in the actual and correct assessment of the candidates. A small mistake in evaluation may lead to serious problems which may affect the future of the candidates, education system and teaching profession. To avoid mistakes, it is requested that before starting evaluation, you must read and understand the spot evaluation guidelines carefully.
2	“Evaluation policy is a confidential policy as it is related to the confidentiality of the examinations conducted, Evaluation done and several other aspects. Its leakage to the public in any manner could lead to derailment of the examination system and affect the life and future of millions of candidates. Sharing this policy/document to anyone, publishing in any magazine and printing in Newspaper/Website, etc. may invite action under various rules of the Board and IPC.”
3	Evaluation is to be done as per instructions provided in the Marking Scheme. It should not be done according to one’s own interpretation or any other consideration. The Marking Scheme should be strictly adhered to and religiously followed. However, while evaluating, answers which are based on latest information or knowledge and/or are innovative, they may be assessed for their correctness otherwise and due marks be awarded to them. In class-XII, while evaluating the competency-based questions, please try to understand the given answer and even if reply is not from a marking scheme but correct competency is enumerated by the candidate, due marks should be awarded.
4	The Marking Scheme carries only suggested value points for the answers. These are Guidelines only and do not constitute the complete answer. The students can have their own expression and if the expression is correct, the due marks should be awarded accordingly.
5	The Head-Examiner must go through the first five answer books evaluated by each evaluator on the first day, to ensure that evaluation has been carried out as per the instructions given in the Marking Scheme. If there is any variation, the same should be zero after deliberation and discussion. The remaining answer books meant for evaluation shall be given only after ensuring that there is no significant variation in the marking of individual evaluators.
6	Evaluators will mark (√) wherever answer is correct. For wrong answer CROSS ‘X’ be marked. Evaluators will not put right (✓) while evaluating which gives the impression that the answer is correct, and no marks are awarded. This is the most common mistake which evaluators are committing.
7	If a question has parts, please award marks on the right-hand side for each part. Marks awarded for different parts of the question should then be totaled up and written in the left-hand margin and encircled. This may be followed strictly.
8	If a question does not have any parts, marks must be awarded in the left-hand margin and encircled. This may also be followed strictly.
9	If a student has attempted an extra question, answer to the question deserving more marks should be retained and the other answer scored out with a note “Extra Question” .

10	No marks to be deducted for the cumulative effect of an error. It should be penalized only once.
11	A full scale of marks (example 0 to 80/70/60/50/40/30 marks as given in Question Paper) has to be used. Please do not hesitate to award full marks if the answer deserves it.
12	Every examiner must necessarily do evaluation work for full working hours, i.e., 8 hours every day and evaluate 20 answer books per day in main subjects and 25 answer books per day in other subjects (Details are given in Spot Guidelines). This is in view of the reduced syllabus and number of questions in question paper.
13	Ensure that you do not make the following common types of errors committed by the Examiner in the past: - <ul style="list-style-type: none"> ● Leaving answer or part thereof unassessed in an answer book. ● Giving more marks for an answer than assigned to it. ● Wrong totaling of marks awarded on an answer. ● Wrong transfer of marks from the inside pages of the answer book to the title page. ● Wrong question wise totaling on the title page. ● Wrong totaling of marks of the two columns on the title page. ● Wrong grand total. ● Marks in words and figures not tallying/not same. ● Wrong transfer of marks from the answer book to online award list. ● Answers marked as correct, but marks not awarded. (Ensure that the right tick mark is correctly and clearly indicated. It should merely be a line. Same is with the X for incorrect answer.) ● Half or a part of the answer marked correct and the rest as wrong, but no marks awarded.
14	While evaluating the answer books if the answer is found to be totally incorrect, it should be marked as cross (X) and awarded zero (0) Marks.
15	Any unassessed portion, non-carrying over of marks to the title page, or total error detected by the candidate shall damage the prestige of all the personnel engaged in the evaluation work as also of the Board. Hence, to uphold the prestige of all concerned, it is again reiterated that the instructions be followed meticulously and judiciously.
16	The Examiners should acquaint themselves with the guidelines given in the “ Guidelines for Spot Evaluation ” before starting the actual evaluation.
17	Every Examiner shall also ensure that all the answers are evaluated, marks carried over to the title page, correctly totaled and written in figures and words.
18	The candidates are entitled to obtain a photocopy of the Answer Book on request on payment of the prescribed processing fee. All Examiners/Additional Head Examiners/Head Examiners are once again reminded that they must ensure that evaluation is carried out strictly as per value points for each answer as given in the Marking Scheme.

Ans	(B) $\frac{16}{3}$	1
10.	<p>If $f(x) = \{[x], x \in \mathbb{R}\}$ is the greatest integer function, then the correct statement is :</p> <p>(A) f is continuous but not differentiable at $x = 2$. (B) f is neither continuous nor differentiable at $x = 2$. (C) f is continuous as well as differentiable at $x = 2$. (D) f is not continuous but differentiable at $x = 2$.</p>	
Ans	(B) f is neither continuous nor differentiable at $x=2$.	1
11.	<p>$\int \frac{\cos 2x - \cos 2\theta}{\cos x - \cos \theta} dx$ is equal to :</p> <p>(A) $2(\sin x + x \cos \theta) + C$ (B) $2(\sin x - x \cos \theta) + C$ (C) $2(\sin x + \sin \theta) + C$ (D) $2(\sin x - x \sin \theta) + C$</p>	
Ans	(A) $2(\sin x + x \cos \theta) + C$	1
12.	<p>$\int_0^1 \frac{2x}{5x^2 + 1} dx$ is equal to :</p> <p>(A) $\frac{1}{5} \log 6$ (B) $\frac{1}{5} \log 5$ (C) $\frac{1}{2} \log 6$ (D) $\frac{1}{2} \log 5$</p>	
Ans	(A) $\frac{1}{5} \log 6$	1
13.	<p>The slope of the curve $y = -x^3 + 3x^2 + 8x - 20$ is maximum at :</p> <p>(A) $(1, -10)$ (B) $(1, 10)$ (C) $(10, 1)$ (D) $(-10, 1)$</p>	
Ans	(A) $(1, -10)$	1

17.	<p>The respective values of \vec{a} and \vec{b}, if given $(\vec{a} - \vec{b}) \cdot (\vec{a} + \vec{b}) = 512$ and $\vec{a} = 3 \vec{b}$, are :</p> <p>(A) 48 and 16 (B) 3 and 1 (C) 24 and 8 (D) 6 and 2</p>	
Ans	(C) 24 and 8	1
18.	<p>Let \vec{a} be a position vector whose tip is the point $(2, -3)$. If $\vec{AB} = \vec{a}$, where coordinates of A are $(-4, 5)$, then the coordinates of B are :</p> <p>(A) $(-2, -2)$ (B) $(2, -2)$ (C) $(-2, 2)$ (D) $(2, 2)$</p>	
Ans	(C) $(-2, 2)$	1
	<p><i>Questions number 19 and 20 are Assertion and Reason based questions. Two statements are given, one labelled Assertion (A) and the other labelled Reason (R). Select the correct answer from the codes (A), (B), (C) and (D) as given below.</i></p> <p>(A) Both Assertion (A) and Reason (R) are true and Reason (R) is the correct explanation of the Assertion (A).</p> <p>(B) Both Assertion (A) and Reason (R) are true, but Reason (R) is not the correct explanation of the Assertion (A).</p> <p>(C) Assertion (A) is true, but Reason (R) is false.</p> <p>(D) Assertion (A) is false, but Reason (R) is true.</p>	

19.	<p><i>Assertion (A) :</i> The shaded portion of the graph represents the feasible region for the given Linear Programming Problem (LPP).</p>  <p>Min $Z = 50x + 70y$ subject to constraints $2x + y \geq 8$, $x + 2y \geq 10$, $x, y \geq 0$ $Z = 50x + 70y$ has a minimum value = 380 at $B(2, 4)$.</p> <p><i>Reason (R) :</i> The region representing $50x + 70y < 380$ does not have any point common with the feasible region.</p>	
Ans	(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).	1
20.	<p><i>Assertion (A) :</i> Let $A = \{x \in \mathbb{R} : -1 \leq x \leq 1\}$. If $f : A \rightarrow A$ be defined as $f(x) = x^2$, then f is not an onto function.</p> <p><i>Reason (R) :</i> If $y = -1 \in A$, then $x = \pm \sqrt{-1} \notin A$.</p>	
Ans	(A) Both Assertion (A) and Reason (R) are true, and Reason (R) is the correct explanation of the Assertion (A).	1
<p>SECTION-B</p> <p><i>This section comprises 5 Very Short Answer (VSA) type questions of 2 marks each.</i></p>		
21.	Find the domain of $\sin^{-1}(x^2 - 3)$.	
Ans	<p>Domain of $\sin^{-1} x$ is $[-1, 1]$</p> <p>$-1 \leq x^2 - 3 \leq 1 \Rightarrow 2 \leq x^2 \leq 4$</p> <p>$\Rightarrow$ Domain = $[-2, -\sqrt{2}] \cup [\sqrt{2}, 2]$</p>	<p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>

22.	Let the volume of a metallic hollow sphere be constant. If the inner radius increases at the rate of 2 cm/s, find the rate of increase of the outer radius when the radii are 2 cm and 4 cm respectively.	
Ans	$\frac{dr}{dt} = 2 \text{ cm/s}, \left(\frac{dR}{dt}\right)_{R=4, r=2} = ?$ $V = \frac{4}{3}\pi(R^3 - r^3) \Rightarrow \frac{dV}{dt} = \frac{4}{3}\pi(3R^2 \cdot \frac{dR}{dt} - 3r^2 \frac{dr}{dt})$ <p>When R = 4 cm and r = 2 cm,</p> $0 = \frac{4}{3}\pi[3(4)^2 \cdot \frac{dR}{dt} - 3(2)^2(2)]$ $\Rightarrow \frac{dR}{dt} = \frac{1}{2} \text{ cm/s}$	<p>1</p> <p>1/2</p> <p>1/2</p>
23.	A man needs to hang two lanterns on a straight wire whose end points have coordinates A (4, 1, -2) and B (6, 2, -3). Find the coordinates of the points where he hangs the lanterns such that these points trisect the wire AB.	
Ans	 <p>Let P and Q trisect the wire AB.</p> <p>P divides AB in the ratio 1:2 then, coordinate of point P = $\left(\frac{14}{3}, \frac{4}{3}, -\frac{7}{3}\right)$</p> <p>Q divides AB in the ratio 2:1 then, coordinate of point Q = $\left(\frac{16}{3}, \frac{5}{3}, -\frac{8}{3}\right)$</p>	<p>1</p> <p>1</p>
24.	<p>(a) Differentiate $\frac{\sin x}{\sqrt{\cos x}}$ with respect to x.</p> <p style="text-align: center;">OR</p> <p>(b) If $y = 5 \cos x - 3 \sin x$, prove that $\frac{d^2y}{dx^2} + y = 0$.</p>	
Ans	<p>(a) Let $y = \frac{\sin x}{\sqrt{\cos x}}$</p> $\frac{dy}{dx} = \frac{\sqrt{\cos x} \cdot \cos x - \sin x \cdot \left(\frac{-\sin x}{2\sqrt{\cos x}}\right)}{\cos x}$ $\Rightarrow \frac{dy}{dx} = \frac{2\cos^2 x + \sin^2 x}{2(\cos x)^{3/2}} \text{ OR } \frac{1 + \cos^2 x}{2(\cos x)^{3/2}}$	<p>1 1/2</p> <p>1/2</p>

OR

(b) $y = 5\cos x - 3\sin x$, then $\frac{dy}{dx} = -5\sin x - 3\cos x$

$$\Rightarrow \frac{d^2y}{dx^2} = -5\cos x + 3\sin x = -y$$

$$\Rightarrow \frac{d^2y}{dx^2} + y = 0$$

1

1/2

1/2

25.

- (a) Find a vector of magnitude 5 which is perpendicular to both the vectors $3\hat{i} - 2\hat{j} + \hat{k}$ and $4\hat{i} + 3\hat{j} - 2\hat{k}$.

OR

- (b) Let \vec{a} , \vec{b} and \vec{c} be three vectors such that $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c}$ and $\vec{a} \times \vec{b} = \vec{a} \times \vec{c}$, $\vec{a} \neq 0$. Show that $\vec{b} = \vec{c}$.

Ans

Let $\vec{a} = 3\hat{i} - 2\hat{j} + \hat{k}$, $\vec{b} = 4\hat{i} + 3\hat{j} - 2\hat{k}$

$$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ 3 & -2 & 1 \\ 4 & 3 & -2 \end{vmatrix} = \hat{i} + 10\hat{j} + 17\hat{k}$$

$$|\vec{a} \times \vec{b}| = \sqrt{1^2 + 10^2 + 17^2} = \sqrt{390}$$

$$\text{Unit vector } \hat{n} = \frac{\vec{a} \times \vec{b}}{|\vec{a} \times \vec{b}|} = \frac{1}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$$

$$\therefore \text{Required vector} = \frac{5}{\sqrt{390}}(\hat{i} + 10\hat{j} + 17\hat{k})$$

OR

(b) $\vec{a} \cdot \vec{b} = \vec{a} \cdot \vec{c} \Rightarrow \vec{a} \cdot (\vec{b} - \vec{c}) = 0$

$$\Rightarrow \text{either } \vec{b} = \vec{c} \text{ or } \vec{a} \perp (\vec{b} - \vec{c}), \text{ since } \vec{a} \neq 0$$

$$\text{Also, } \vec{a} \times \vec{b} = \vec{a} \times \vec{c} \Rightarrow \vec{a} \times (\vec{b} - \vec{c}) = 0$$

$$\Rightarrow \text{either } \vec{b} = \vec{c} \text{ or } \vec{a} \parallel (\vec{b} - \vec{c}), \text{ since } \vec{a} \neq 0$$

Since vectors \vec{a} and $(\vec{b} - \vec{c})$ cannot be \parallel and \perp simultaneously

$$\text{Hence } \vec{b} = \vec{c}$$

1/2

1/2

1/2

1/2

1

1/2

1/2

SECTION-C

This section comprises 6 Short Answer (SA) type questions of 3 marks each.

26.

Find the interval/intervals in which the function $f(x) = \sin 3x - \cos 3x$, $0 < x < \frac{\pi}{2}$ is strictly increasing.

Ans	$f'(x) = 3 \cos 3x + 3 \sin 3x$ $f'(x) = 0 \Rightarrow \sin 3x = -\cos 3x \Rightarrow x = \frac{\pi}{4}$ For $x \in \left(0, \frac{\pi}{4}\right)$, $3 \cos 3x + 3 \sin 3x > 0$ $\Rightarrow f'(x) > 0$, f is strictly increasing function in $\left(0, \frac{\pi}{4}\right)$ or $\left(0, \frac{\pi}{4}\right]$	1 $\frac{1}{2}$ 1 $\frac{1}{2}$
27.	<p>(a) If $\vec{a} + \vec{b} + \vec{c} = \vec{0}$ such that $\vec{a} = 3$, $\vec{b} = 5$, $\vec{c} = 7$, then find the angle between \vec{a} and \vec{b}.</p> <p style="text-align: center;">OR</p> <p>(b) If \vec{a} and \vec{b} are unit vectors inclined with each other at an angle θ, then prove that $\frac{1}{2} \vec{a} - \vec{b} = \sin \frac{\theta}{2}$.</p>	
Ans	Given $\vec{a} + \vec{b} + \vec{c} = \vec{0} \Rightarrow \vec{a} + \vec{b} = -\vec{c} $ $\Rightarrow \vec{a} + \vec{b} ^2 = \vec{c} ^2 \Rightarrow \vec{a} ^2 + \vec{b} ^2 + 2\vec{a} \cdot \vec{b} = \vec{c} ^2$ $\Rightarrow 9 + 25 + 2\vec{a} \cdot \vec{b} = 49$ $\Rightarrow 2 \vec{a} \vec{b} \cos \theta = 15$ $\Rightarrow \cos \theta = \frac{1}{2} \Rightarrow \theta = \frac{\pi}{3}$ <p style="text-align: center;">OR</p> <p>(b) $\vec{a} = \vec{b} = 1$ $\vec{a} - \vec{b} ^2 = \vec{a} ^2 + \vec{b} ^2 - 2\vec{a} \cdot \vec{b}$ $= 1 + 1 - 2 \vec{a} \vec{b} \cos \theta$ $= 2 - 2\cos \theta$ $= 2 \left(2\sin^2 \frac{\theta}{2}\right) = 4\sin^2 \frac{\theta}{2}$ $\Rightarrow \sin \frac{\theta}{2} = \frac{1}{2} \vec{a} - \vec{b}$ </p>	1 1 1 $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$ $\frac{1}{2}$

28.	Solve the differential equation $x \cos\left(\frac{y}{x}\right) \frac{dy}{dx} = y \cos\left(\frac{y}{x}\right) + x.$	
Ans	$\Rightarrow \frac{dy}{dx} = \frac{y}{x} + \sec \frac{y}{x}$ <p style="text-align: center;">Put $y = vx$</p> $\Rightarrow \frac{dy}{dx} = v + x \frac{dv}{dx}$ $\Rightarrow v + x \frac{dv}{dx} = v + \sec v$ $\Rightarrow \int \cos v \, dv = \int \frac{dx}{x}$ $\Rightarrow \sin v = \log x + c$ $\Rightarrow \sin \frac{y}{x} = \log x + c$	$\frac{1}{2}$ $\frac{1}{2}$ 1 $\frac{1}{2}$ $\frac{1}{2}$
29.	<p>(a) The probability that a student buys a colouring book is 0.7 and that she buys a box of colours is 0.2. The probability that she buys a colouring book, given that she buys a box of colours, is 0.3. Find the probability that the student :</p> <p>(i) Buys both the colouring book and the box of colours.</p> <p>(ii) Buys a box of colours given that she buys the colouring book.</p> <p style="text-align: center;">OR</p> <p>(b) A person has a fruit box that contains 6 apples and 4 oranges. He picks out a fruit three times, one after the other, after replacing the previous one in the box. Find :</p> <p>(i) The probability distribution of the number of oranges he draws.</p> <p>(ii) The expectation of the random variable (number of oranges).</p>	
Ans	<p>(a) Let A be the event of buying colouring book and B be the event of buying coloured box.</p>	$\left. \vphantom{\begin{matrix} \\ \\ \end{matrix}} \right\}$ $\frac{1}{2}$

	<p>$P(A) = 0.7, \quad P(B) = 0.2, \quad P(A/B) = 0.3$</p> <p>(i) $P\left(\frac{A}{B}\right) = \frac{P(A \cap B)}{P(B)} \Rightarrow 0.3 = \frac{P(A \cap B)}{0.2}$</p> <p>$\Rightarrow P(A \cap B) = 0.06$ or $\frac{3}{50}$</p> <p>(ii) $P\left(\frac{B}{A}\right) = \frac{P(A \cap B)}{P(A)}$</p> <p>$= \frac{0.06}{0.7} = \frac{3}{35}$ or 0.086</p> <p style="text-align: center;">OR</p> <p>(b) Let X be random variable for number of oranges.</p> <p>$X = 0, 1, 2, 3$</p> <p>Let A be the event that orange is drawn.</p> <p>$P(A) = \frac{4}{10} = \frac{2}{5}, \quad P(\bar{A}) = 1 - \frac{2}{5} = \frac{3}{5}$</p> <p>(i)</p> <table border="1" style="margin-left: auto; margin-right: auto;"> <thead> <tr> <th>X</th> <th>0</th> <th>1</th> <th>2</th> <th>3</th> </tr> </thead> <tbody> <tr> <td>$P(X)$</td> <td>$\frac{27}{125}$</td> <td>$\frac{54}{125}$</td> <td>$\frac{36}{125}$</td> <td>$\frac{8}{125}$</td> </tr> </tbody> </table> <p>(ii) $E(X) = \sum p_i x_i = 0 \times \frac{27}{125} + 1 \times \frac{54}{125} + 2 \times \frac{36}{125} + 3 \times \frac{8}{125}$</p> <p>$= \frac{150}{125}$ or $\frac{6}{5}$</p>	X	0	1	2	3	$P(X)$	$\frac{27}{125}$	$\frac{54}{125}$	$\frac{36}{125}$	$\frac{8}{125}$	<p>$\frac{1}{2}$</p> <p>1</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p> <p>1</p> <p>$\frac{1}{2}$</p> <p>$\frac{1}{2}$</p>
X	0	1	2	3								
$P(X)$	$\frac{27}{125}$	$\frac{54}{125}$	$\frac{36}{125}$	$\frac{8}{125}$								
30.	<p>(a) Find :</p> $\int \frac{2x}{(x^2 + 3)(x^2 - 5)} dx$ <p style="text-align: center;">OR</p> <p>(b) Evaluate :</p> $\int_1^4 (x-2 + x-4) dx$											
Ans	<p>(a) Let $I = \int \frac{2x}{(x^2+3)(x^2-5)} dx$</p> <p>Put $x^2 = t \Rightarrow 2x \cdot dx = dt$</p>	$\frac{1}{2}$										

$$\Rightarrow I = \int \frac{dt}{(t+3)(t-5)}$$

$$= \int \left(-\frac{1}{8(t+3)} + \frac{1}{8(t-5)} \right) dt$$

$$= \frac{1}{8} [\log |t-5| - \log |t+3|] + c$$

$$= \frac{1}{8} \log \left| \frac{x^2-5}{x^2+3} \right| + c$$

OR

(b) $\int_1^4 (|x-2| + |x-4|) dx$

$$= \int_1^2 (2-x) dx + \int_2^4 (x-2) dx - \int_1^4 (x-4) dx$$

$$= \left[\frac{(2-x)^2}{-2} \right]_1^2 + \left[\frac{(x-2)^2}{2} \right]_2^4 - \left[\frac{(x-4)^2}{2} \right]_1^4$$

$$= \frac{1}{2} + 2 + \frac{9}{2} = 7$$

1

1

1/2

1 1/2

1

1/2

31. In the Linear Programming Problem (LPP), find the point/points giving maximum value for $Z = 5x + 10y$ subject to constraints


$$x + 2y \leq 120$$

$$x + y \geq 60$$

$$x - 2y \geq 0$$

$$x, y \geq 0$$

Ans



1/2 for correct graph and correct feasible region

Corner Points	Value of Z
A (60, 30)	600
B (120, 0)	600
C (60, 0)	300
D (40, 20)	400

Since Z is maximum on points A and B

Hence all points lying on segment AB give maximum Z.

1


1/2

SECTION-D

This section comprises 4 Long Answer (LA) type questions of 5 marks each.

32. In a rough sketch, mark the region bounded by $y = 1 + |x + 1|$, $x = -2$, $x = 2$ and $y = 0$. Using integration, find the area of the marked region.

Ans




s

2 marks
for
correct
graph
and
shading

	<p>Required area = $\int_{-2}^{-1}(-x) dx + \int_{-1}^2(x+2) dx$</p> $= -\frac{1}{2} [x^2]_{-2}^{-1} + \left[\frac{1}{2}x^2 + 2x\right]_{-1}^2$ $= 9$	<p>1½</p> <p>1</p> <p>½</p>
33.	<p>Three students run on a racing track such that their speeds add up to 6 km/h. However, double the speed of the third runner added to the speed of the first results in 7 km/h. If thrice the speed of the first runner is added to the original speeds of the other two, the result is 12 km/h. Using matrix method, find the original speed of each runner.</p>	
Ans	<p>Let original speed of three runners be x, y and z respectively.</p> <p>Then $x + y + z = 6$; $x + 2z = 7$; $3x + y + z = 12$</p> <p>Let $A = \begin{bmatrix} 1 & 1 & 1 \\ 1 & 0 & 2 \\ 3 & 1 & 1 \end{bmatrix}$, $X = \begin{bmatrix} x \\ y \\ z \end{bmatrix}$, $B = \begin{bmatrix} 6 \\ 7 \\ 12 \end{bmatrix}$</p> <p>$A = 4 \neq 0 \Rightarrow A^{-1}$ exists</p> <p>$AX = B \Rightarrow X = A^{-1}B$</p> <p>$\text{adj}(A) = \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$</p> <p>$A^{-1} = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix}$</p> <p>$\therefore \begin{bmatrix} x \\ y \\ z \end{bmatrix} = \frac{1}{4} \begin{bmatrix} -2 & 0 & 2 \\ 5 & -2 & -1 \\ 1 & 2 & -1 \end{bmatrix} \begin{bmatrix} 6 \\ 7 \\ 12 \end{bmatrix} = \begin{bmatrix} 3 \\ 1 \\ 2 \end{bmatrix}$</p> <p>Hence the original speed of three runners are 3km/h, 1km/h and 2 km/h respectively.</p>	<p>1½</p> <p>½</p> <p>½</p> <p>1½</p> <p>1</p>

34.	<p>(a) For a positive constant 'a', differentiate $a^{t+\frac{1}{t}}$ with respect to $\left(t+\frac{1}{t}\right)^a$, where t is a non-zero real number.</p> <p style="text-align: center;">OR</p> <p>(b) Find $\frac{dy}{dx}$ if $y^x + x^y + x^x = a^b$, where a and b are constants.</p>	
Ans	<p>(a) Let $u = a^{t+\frac{1}{t}} \Rightarrow \frac{du}{dt} = a^{t+\frac{1}{t}} \cdot \log a \cdot \left(1 - \frac{1}{t^2}\right)$</p> <p style="text-align: center;">$v = \left(t + \frac{1}{t}\right)^a \Rightarrow \frac{dv}{dt} = a \left(t + \frac{1}{t}\right)^{a-1} \cdot \left(1 - \frac{1}{t^2}\right)$</p> <p style="text-align: center;">$\frac{du}{dv} = \frac{du/dt}{dv/dt} = \frac{a^{t+\frac{1}{t}} \cdot \log a}{a \left(t + \frac{1}{t}\right)^{a-1}}$</p> <p style="text-align: center;">OR</p> <p>(b) Let $u = y^x$, $v = x^y$ and $w = x^x$</p> <p>$\Rightarrow \frac{du}{dx} + \frac{dv}{dx} + \frac{dw}{dx} = 0$(i)</p> <p>$u = y^x \Rightarrow \log u = x \cdot \log y \Rightarrow \frac{1}{u} \cdot \frac{du}{dx} = \frac{x}{y} \cdot \frac{dy}{dx} + \log y$</p> <p>$\Rightarrow \frac{du}{dx} = y^x \left(\frac{x}{y} \cdot \frac{dy}{dx} + \log y\right) = xy^{x-1} \frac{dy}{dx} + y^x \log y$</p> <p>$v = x^y \Rightarrow \log v = y \cdot \log x \Rightarrow \frac{1}{v} \cdot \frac{dv}{dx} = \frac{y}{x} + \log x \cdot \frac{dy}{dx}$</p> <p>$\Rightarrow \frac{dv}{dx} = x^y \left(\frac{y}{x} + \log x \cdot \frac{dy}{dx}\right) = yx^{y-1} + x^y \log x \frac{dy}{dx}$</p> <p>$w = x^x \Rightarrow \log w = x \cdot \log x \Rightarrow \frac{1}{w} \cdot \frac{dw}{dx} = 1 + \log x$</p> <p>$\Rightarrow \frac{dw}{dx} = x^x \cdot (1 + \log x)$</p> <p>$\therefore$ From (i), we get</p> <p style="text-align: center;">$xy^{x-1} \cdot \frac{dy}{dx} + y^x \cdot \log y + yx^{y-1} + x^y \cdot \log x \cdot \frac{dy}{dx} + x^x \cdot (1 + \log x) = 0$</p> <p style="text-align: center;">$\Rightarrow \frac{dy}{dx} = -\frac{x^x \cdot (1 + \log x) + y^x \cdot \log y + yx^{y-1}}{x \cdot y^{x-1} + x^y \cdot \log x}$</p>	<p style="text-align: right;">2</p> <p style="text-align: right;">2</p> <p style="text-align: right;">1</p>


35.	<p>(a) Find the foot of the perpendicular drawn from the point (1, 1, 4) on the line $\frac{x+2}{5} = \frac{y+1}{2} = \frac{-z+4}{-3}$.</p> <p style="text-align: center;">OR</p> <p>(b) Find the point on the line $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3}$ at a distance of $2\sqrt{2}$ units from the point (-1, -1, 2).</p>	
Ans	<p>(a) Let $\frac{x+2}{5} = \frac{y+1}{2} = \frac{z-4}{3} = \lambda$</p> <p>Coordinate of general point on the given line are M (5λ - 2, 2λ - 1, 3λ + 4)</p> <div style="text-align: center;">  </div> <p>Direction Ratios of PM vector are $\langle 5\lambda - 3, 2\lambda - 2, 3\lambda \rangle$</p> <p>Since, $PM \perp l$</p> $\Rightarrow 5(5\lambda - 3) + 2(2\lambda - 2) + 3(3\lambda) = 0$ $\Rightarrow \lambda = \frac{1}{2}$ <p>Hence, coordinates of M are $(\frac{1}{2}, 0, \frac{11}{2})$</p> <p style="text-align: center;">OR</p> <p>(b) Equation of given line be $\frac{x-1}{3} = \frac{y+1}{2} = \frac{z-4}{3} = \lambda$ (say)</p> <p>Coordinate of any general point on the line are P (3λ + 1, 2λ - 1, 3λ + 4).</p> <p>Let distance of point P from (-1, -1, 2) is $2\sqrt{2}$.</p> $\Rightarrow \sqrt{(3\lambda + 2)^2 + (2\lambda)^2 + (3\lambda + 2)^2} = 2\sqrt{2}$ $\Rightarrow 22\lambda^2 + 24\lambda = 0$ $\Rightarrow \lambda = 0 \text{ or } \lambda = -\frac{12}{11}$	<p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> <p>1½</p> <p>1</p>

	Hence, coordinates of point P are $(1, -1, 4)$ or $\left(-\frac{25}{11}, -\frac{35}{11}, \frac{8}{11}\right)$	1½
	SECTION-E	
	<i>This section comprises 3 case study-based questions of 4 marks each</i>	
	Case Study – 1	
36.	<p>Let A be the set of 30 students of class XII in a school. Let $f : A \rightarrow N$, N is a set of natural numbers such that function $f(x) = \text{Roll Number of student } x$.</p> <p>On the basis of the given information, answer the following :</p> <p>(i) Is f a bijective function ? 1</p> <p>(ii) Give reasons to support your answer to (i). 1</p> <p>(iii) (a) Let R be a relation defined by the teacher to plan the seating arrangement of students in pairs, where $R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = 3x\}$. List the elements of R. Is the relation R reflexive, symmetric and transitive ? Justify your answer. 2</p> <p style="text-align: center;">OR</p> <p>(iii) (b) Let R be a relation defined by $R = \{(x, y) : x, y \text{ are Roll Numbers of students such that } y = x^3\}$. List the elements of R. Is R a function ? Justify your answer. 2</p>	
Ans	<p>(i) No, f is not bijective function</p> <p>(ii) Range = $\{1, 2, 3, 4, \dots, 30\}$ and codomain = N Since, Range \neq codomain \Rightarrow f is not onto and hence f is not bijective.</p> <p>(iii) (a) $R = \{(1, 3), (2, 6), (3, 9), (4, 12), (5, 15), (6, 18), (7, 21), (8, 24), (9, 27), (10, 30)\}$</p> <p>Since $(1, 1) \notin R \Rightarrow R$ is not reflexive.</p> <p>$(1, 3) \in R$ but $(3, 1) \notin R \Rightarrow R$ is not symmetric</p> <p>$(1, 3) \in R, (3, 9) \in R$ but $(1, 9) \notin R \Rightarrow R$ is not transitive.</p> <p style="text-align: center;">OR</p> <p>(iii) (b) $R = \{(1, 1), (2, 8), (3, 27)\}$</p> <p>$\because$ elements 4, 5, 6 ... 30 do not have an image. Hence the above relation is not a function.</p>	<p>1</p> <p>½</p> <p>½</p> <p>1</p> <p>1</p> <p>1</p> <p>1</p>

Case Study – 2

37.


A gardener wanted to plant vegetables in his garden. Hence he bought 10 seeds of brinjal plant, 12 seeds of cabbage plant and 8 seeds of radish plant. The shopkeeper assured him of germination probabilities of brinjal, cabbage and radish to be 25%, 35% and 40% respectively. But before he could plant the seeds, they got mixed up in the bag and he had to sow them randomly.



Radish



Cabbage



Brinjal

Based upon the above information, answer the following questions :

- (i) Calculate the probability of a randomly chosen seed to germinate. 2
- (ii) What is the probability that it is a cabbage seed, given that the chosen seed germinates? 2

Ans

Let A: Event that chosen seed germinates.

B: Event that Brinjal seed is chosen.

C: Event that Cabbage seed is chosen.

R: Event that Radish seed is chosen.

$$P(B) = \frac{10}{30}; \quad P(C) = \frac{12}{30}; \quad P(R) = \frac{8}{30};$$

$$P\left(\frac{A}{B}\right) = \frac{25}{100}; \quad P\left(\frac{A}{C}\right) = \frac{35}{100}; \quad P\left(\frac{A}{R}\right) = \frac{40}{100}$$

(i) $P(A) = P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)$

$$= \frac{10}{30} \times \frac{25}{100} + \frac{12}{30} \times \frac{35}{100} + \frac{8}{30} \times \frac{40}{100}$$

$$= \frac{990}{3000} \text{ or } \frac{33}{100}$$

(ii) (a) $P\left(\frac{C}{A}\right) = \frac{P(C) \cdot P\left(\frac{A}{C}\right)}{P(B) \cdot P\left(\frac{A}{B}\right) + P(C) \cdot P\left(\frac{A}{C}\right) + P(R) \cdot P\left(\frac{A}{R}\right)}$

$$= \frac{\frac{12}{30} \times \frac{35}{100}}{\frac{990}{3000}}$$

$$= \frac{42}{99} \text{ or } \frac{14}{33}$$

1

1

1

1

Case Study - 3


38.

A carpenter needs to make a wooden cuboidal box, closed from all sides, which has a square base and fixed volume. Since he is short of the paint required to paint the box on completion, he wants the surface area to be minimum.

On the basis of the above information, answer the following questions :

- (i) Taking length = breadth = x m and height = y m, express the surface area (S) of the box in terms of x and its volume (V), which is constant. 1
- (ii) Find $\frac{dS}{dx}$. 1
- (iii) (a) Find a relation between x and y such that the surface area (S) is minimum. 2
- OR**
- (iii) (b) If surface area (S) is constant, the volume (V) = $\frac{1}{4}(Sx - 2x^3)$, x being the edge of base. Show that volume (V) is maximum for $x = \sqrt{\frac{S}{6}}$. 2

Ans



(i) $V = x^2y \Rightarrow y = \frac{V}{x^2} \dots \dots \dots$ (i)

Hence, $S = 2x^2 + 4xy = 2x^2 + \frac{4V}{x}$ 1

(ii) $\frac{dS}{dx} = 4\left(x - \frac{V}{x^2}\right)$ 1

(iii) (a) $\frac{dS}{dx} = 0 \Rightarrow V = x^3 \Rightarrow x^2y = x^3 \Rightarrow y = x$ 1

$\frac{d^2S}{dx^2} = 4\left(1 + \frac{2V}{x^3}\right) = 12 > 0 \Rightarrow S$ is minimum if $y = x$. 1

OR

(iii) (b) $V = \frac{1}{4}(Sx - 2x^3) \Rightarrow \frac{dV}{dx} = \frac{1}{4}(S - 6x^2)$ 1

Put $\frac{dV}{dx} = 0 \Rightarrow x = \sqrt{\frac{S}{6}}$ 1/2

$\left(\frac{d^2V}{dx^2}\right)_{x=\sqrt{\frac{S}{6}}} = -3\sqrt{\frac{S}{6}} < 0 \Rightarrow$ Volume is maximum for $x = \sqrt{\frac{S}{6}}$. 1/2